



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/20	A1	(11) International Publication Number: WO 90/14436 (43) International Publication Date: 29 November 1990 (29.11.90)								
<p>(21) International Application Number: PCT/US90/02853</p> <p>(22) International Filing Date: 18 May 1990 (18.05.90)</p> <p>(30) Priority data:</p> <table border="0"> <tr> <td>355,961</td> <td>18 May 1989 (18.05.89)</td> <td>US</td> </tr> <tr> <td>456,637</td> <td>21 December 1989 (21.12.89)</td> <td>US</td> </tr> <tr> <td>505,435</td> <td>4 April 1990 (04.04.90)</td> <td>US</td> </tr> </table> <p>(71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US).</p> <p>(72) Inventors: HOUGHTON, Michael ; 53 Rosemead Court, Danville, CA 94526 (US). CHOO, Qui-Lim ; 5700 Fern Street, El Cerrito, CA 94530 (US). KUO, George ; 1370 Sixth Avenue, San Francisco, CA 94112 (US). HAN, Jang ; 3238 Del Mar, Lafayette, CA 94549 (US). URDEA, Michael, S. ; 100 Bunce Meadow Road, Alamo, CA 94501 (US). WEINER, Amy, J. ; 1766 Indian Way, Oakland, CA 94611 (US).</p>	355,961	18 May 1989 (18.05.89)	US	456,637	21 December 1989 (21.12.89)	US	505,435	4 April 1990 (04.04.90)	US	<p>(74) Agents: MONROY, Gladys, H. et al.; Irell & Manella, 545 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).</p> <p>(81) Designated States: AT, AU, BB, BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CM (OAPI patent), DE*, DK, ES, FI, GA (OAPI patent), GB, HU, JP, KP, KR, LK, LU, MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NO, RO, SD, SE, SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
355,961	18 May 1989 (18.05.89)	US								
456,637	21 December 1989 (21.12.89)	US								
505,435	4 April 1990 (04.04.90)	US								

(54) Title: NANBV DIAGNOSTICS: POLYNUCLEOTIDES USEFUL FOR SCREENING FOR HEPATITIS C VIRUS

(57) Abstract

A new virus, Hepatitis C virus (HCV), which has proven to be the major etiologic agent of blood-borne NANBH, was discovered by Applicant. Reagents for isolating, amplifying, and detecting HCV polynucleotides are provided. These reagents are oligomers comprised of polynucleotide sequences which are capable of forming hybrid structures with HCV target polynucleotide sequences.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark			US	United States of America

-1-

5

NANBV DIAGNOSTICS: POLYNUCLEOTIDES USEFUL
FOR SCREENING FOR HEPATITIS C VIRUS

10

Technical Field

The invention relates to materials and methodologies for managing the spread of non-A, non-B hepatitis virus (NANBV) infection. More specifically, it relates to an etiologic agent of non-A, non-B hepatitis (NANBH), hepatitis C virus (HCV), and to polynucleotides and analogs thereof, which are useful in assays for the detection of HCV in biological samples.

20

References Cited in the Application

- Barr et al. (1986), *Biotechniques* 4:428.
- Beaucage et al. (1981), *Tetrahedron Letters* 22:1859.
- Botstein (1979), *Gene* 8:17.
- 25 Brinton, M.A. (1986) in *THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE* (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.327-374.
- Broach (1981) in: *Molecular Biology of the Yeast Saccharomyces*, Vol. 1, p.445, Cold Spring Harbor Press.
- 30 Broach et al. (1983), *Meth. Enz.* 101:307.
- Brown et al. (1979), *Methods in Enzymology* 68:109.
- Byrne et al. (1988), *Nucleic Acids Res.* 16:4165.
- Castle et al. (1986), *Virology* 119:10.
- 35 Chang et al. (1977), *Nature* 198:1056.
- Chirgwin et al. (1979), *Biochemistry* 18:5294.

- Choo et al. (1989), Science 244:359.
- Chomczynski and Sacchi (1987), Analytical Biochemistry 162:156.
- Clewell et al. (1969), Proc. Natl. Acad. Sci. USA 62:1159.
- 5 Clewell (1972), J. Bacteriol. 110:667.
- Cohen (1972), Proc. Natl. Acad. Sci. USA 69:2110.
- Cousens et al. (1987), Gene 61:265.
- De Boer et al. (1983), Proc. Natl. Acad. Sci. USA 292:128.
- Dreesman et al. (1985), J. Infect. Disease 151:761.
- 10 Feinstone et al. (1981), J. Inf. Dis. 144: 588.
- Feinstone et al. (1983), Infection and Immunology 41:816.
- Feinstone, S.M. and Hoofnagle, J.H. (1984), New Engl. J. Med. 311:185.
- Fields & Knipe (1986), FUNDAMENTAL VIROLOGY (Raven Press, N.Y.).
- 15 Fiers et al. (1978), Nature 273:113.
- Gerety, R.J. et al., in VIRAL HEPATITIS AND LIVER DISEASE (Vyas, B.N., Dienstag, J.L., and Hoofnagle, J.H., eds, Grune and Stratton, Inc., 1984) pp 23-47.
- 20 Goeddel et al. (1980), Nucleic Acids Res. 8:4057.
- Graham and Van der Eb (1978), Virology 52:546.
- Grunstein and Hogness (1975), Proc. Natl. Acad. Sci. USA 73:3961.
- Grych et al. (1985), Nature 316:74.
- 25 Gubler and Hoffman (1983), Gene 25:263.
- Hahn et al. (1988), Virology 162:167.
- Han (1987), Biochemistry 26:1617.
- Hammerling et al. (1981), MONOCLONAL ANTIBODIES AND T-CELL HYBRIDOMAS.
- 30 Hess et al. (1968), J. Adv. Enzyme Reg 7:149.
- Hinnen et al. (1978), Proc. Natl. Acad. Sci. 75:1929.
- Hitzeman et al. (1980), J. Biol. Chem. 255:2073.
- Holland et al. (1978), Biochemistry 17:4900.
- Holland (1981), J. Biol. Chem. 256: 1385.
- 35 Houghton et al. (1981), Nucleic Acids Res. 9:247
- Hunyh, T.V. et al. (1985) in DNA CLONING TECHNIQUES; A

- PRACTICAL APPROACH (D. Glover, Ed., IRL Press, Oxford, U.K.) pp. 49-78.
- Immun. Rev. (1982) 62:185.
- Iwarson (1987), British Medical J. 295:946.
- 5 Kennett et al. (1980) MONOCLONAL ANTIBODIES.
- Kuo et al. (1989), Science 244:362.
- Kyte and Doolittle (1982), J. Mol. Biol. 157:105-132.
- Landegren et al. (1988), Science 242:229.
- Maniatis, T., et al. (1982) MOLECULAR CLONING; A
- 10 LABORATORY MANUAL (Cold Spring Harbor Press, Cold Spring Harbor, N.Y.).
- Matthews and Kricka (1988), Analytical Biochemistry 169:1.
- METHODS IN ENZYMOLOGY (Academic Press).
- Mittlin (1989), Clinical chem. 35:1819.
- 15 Laemmli (1970), Nature 227, 680.
- Lee et al. (1988), Science 239:1288.
- Loh et al. (1989), Science 243:217.
- Mackow et al. (1987), Virology 159:217.
- Mayer and Walker, eds. (1987), IMMUNOCHEMICAL METHODS IN
- 20 CELL AND MOLECULAR BIOLOGY (Academic Press, London).
- Maxam et al. (1980), Methods in Enzymology 65:499.
- MacNamara et al. (1984), Science 226:1325.
- Messing et al. (1981), Nucleic Acids Res. 9:309.
- Messing (1983), Methods in Enzymology 101:20-37.
- 25 METHODS IN ENZYMOLOGY (Academic Press).
- Michelle et al., Int. Symposium on Viral Hepatitis.
- Monath (1986) in THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press),
- 30 p.375-440.
- Murakawa et al. (1988), DNA 7:287.
- Nagahuma et al. (1984), Anal. Biochem. 141:74.
- Narang et al (1979), Methods in Enzymology 68:90.
- Neurath et al. (1984), Science 224:392.
- 35 Nisonoff et al. (1981), Clin. Immunol. Immunopathol. 21:397-406.

- Overby, L.R. (1985), *Curr. Hepatol.* 5:49.
- Overby, L.R. (1986), *Curr. Hepatol.* 6:65.
- Overby, L.R. (1987), *Curr. Hepatol.* 7:35.
- Peleg (1969), *Nature* 221:193.
- 5 Pfefferkorn and Shapiro (1974), in *COMPREHENSIVE VIROLOGY*, Vol. 2 (Fraenkel-Conrat & Wagner, eds., Plenum, N.Y.) pp. 171-230.
- Prince, A.M. (1983), *Annu. Rev. Microbiol.* 37:217.
- Rice et al. (1985), *Science* 229:726.
- 10 Rice et al. (1986) in *THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE* (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.279-328.
- Roehrig (1986) in *THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE* (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press)
- 15 Rosenberg et al. (1984), *Nature* 312:7.
- Sadler et al. (1980), *Gene* 8, 279.
- Saiki et al. (1985), *Science* 230:1350.
- 20 Saiki et al. (1986), *Nature* 324: 163.
- Saiki et al. (1988), *Science* 239:487.
- Sanger et al. (1977), *Proc. Natl. Acad. Sci. USA* 74:5463.
- Scharf et al. (1986), *Science* 233:1076.
- Schlesinger et al. (1986), *J. Virol.* 60:1153.
- 25 Schreier, M., et al. (1980) *HYBRIDOMA TECHNIQUES*
- Scopes (1984), *PROTEIN PURIFICATION, PRINCIPLES AND PRACTICE, SECOND EDITION* (Springer-Verlag, N.Y.).
- Shimatake et al. (1981), *Nature* 292:128.
- Shigekawa and Dower (1988), *BioTechniques* 6:742.
- 30 Steimer et al. (1986), *J. Virol.* 58:9.
- Stollar (1980), in *THE TOGAVIRUSES* (R.W. Schlesinger, ed., Academic Press, N.Y.), pp. 584-622.
- Sumiyoshi et al. (1987), *Virology* 161:497.
- Taylor et al. (1976), *Biochem. Biophys. Acta* 442:324.
- 35 Towbin et al. (1979), *Proc. Natl. Acad. Sci. USA* 76, 4350.
- Tsu and Herzenberg (1980), in *SELECTED METHODS IN CELLULAR*

- IMMUNOLOGY (W.H. Freeman and Co.) pp. 373-391.
- Vytdehaag et al. (1985), J. Immunol. 134:1225.
- Valenzuela, P., et al. (1982), Nature 298:344.
- Valenzuela, P., et al. (1984), in HEPATITIS B (Millman,
5 I., et al., ed, Plenum Press) pp. 225-236.
- Warner (1984), DNA 3:401.
- Wu and Grossman (1987), Methods in Enzymology Vol. 154,
RECOMBINANT DNA, Part E.
- Wu (1987), Methods in Enzymology vol 155, RECOMBINANT DNA,
10 part F.
- Zoller (1982), Nucleic Acids Res. 10:6487.

Cited Patents

- U.S. Patent No. 4,341,761
- 15 U.S. Patent No. 4,399,121
- U.S. Patent No. 4,427,783
- U.S. Patent No. 4,444,887
- U.S. Patent No. 4,466,917
- U.S. Patent No. 4,472,500
- 20 U.S. Patent No. 4,491,632
- U.S. Patent No. 4,493,890
- U.S. Patent No. 4,683,202
- U.S. Patent No. 4,458,066
- U.S. Patent No. 4,868,105

25

Background Art

- Non-A, Non-B hepatitis (NANBH) is a
transmissible disease or family of diseases that are
believed to be viral-induced, and that are distinguishable
30 from other forms of viral-associated liver diseases,
including that caused by the known hepatitis viruses,
i.e., hepatitis A virus (HAV), hepatitis B virus (HBV),
and delta hepatitis virus (HDV), as well as the hepatitis
induced by cytomegalovirus (CMV) or Epstein-Barr virus
35 (EBV). NANBH was first identified in transfused
individuals. Transmission from man to chimpanzee and se-

rial passage in chimpanzees provided evidence that NANBH is due to a transmissible infectious agent or agents.

Epidemiologic evidence is suggestive that there may be three types of NANBH: the water-borne epidemic
5 type; the blood or needle associated type; and the sporadically occurring (community acquired) type. However, the number of agents which may be the causative of NANBH are unknown.

There have been a number of candidate NANBV.
10 See, for example the reviews by Prince (1983), Feinstone and Hoofnagle (1984), and Overby (1985, 1986, 1987) and the article by Iwarson (1987). However, there is no proof that any of these candidates represent the etiological agent of NANBH.

15 The demand for sensitive, specific methods for screening and identifying carriers of NANBV and NANBV contaminated blood or blood products is significant. Post-transfusion hepatitis (PTH) occurs in approximately 10% of transfused patients, and NANBH accounts for up to
20 90% of these cases. The major problem in this disease is the frequent progression to chronic liver damage (25-55%).

Patient care as well as the prevention of transmission of NANBH by blood and blood products or by close personal contact require reliable screening,
25 diagnostic and prognostic tools to detect nucleic acids, antigens and antibodies related to NANBV.

Methods for detecting specific polynucleotides by hybridization assays are known in the art. See, for example, Matthews and Kricka (1988), Analytical Bio-
30 chemistry 169:1; Landegren et al. (1988), Science 242:229; and Mittlin (1989), Clinical chem. 35:1819. U.S. Patent No. 4,868,105, issued Sept. 9, 1989, and in E.P.O. Publication No. 225,807 (published June 16, 1987).

Applicant discovered a new virus, the Hepatitis
35 C virus (HCV), which has proven to be the major etiologic agent of blood-borne NANBH (BB-NANBH). Applicant's

initial work, including a partial genomic sequence of the prototype HCV isolate, CDC/HCV1 (also called HCV1), is described in E.P.O. Publication No. 318,216 (published 31 May 1989) and PCT Pub. No. WO 89/04669 (published 1 June 1989). The disclosures of these patent applications, as well as any corresponding national patent applications, are incorporated herein by reference. These applications teach, inter alia, recombinant DNA methods of cloning HCV sequences, HCV probe diagnostic techniques, anti-HCV antibodies, and methods of isolating new HCV sequences.

Disclosure of the Invention

The present invention is based on HCV sequences described in E.P.O. Publication No. 318,216 and in PCT Pub. No. WO 89/04669, as well as other HCV sequences that are described herein. Methods for isolating and/or detecting specific polynucleotides by hybridization could not be used for screening for HCV until Applicants' discovery of HCV. Accordingly, one aspect of the invention is an oligomer capable of hybridizing to an HCV sequence in an analyte polynucleotide strand, wherein the oligomer is comprised of an HCV targeting sequence complementary to at least 4 contiguous nucleotides of HCV cDNA shown in Fig. 18.

Another aspect of the invention is a process for detecting an HCV sequence in an analyte strand suspected of containing an HCV polynucleotide, wherein the HCV polynucleotide comprises a selected target region, said process comprising:

(a) providing an oligomer capable of hybridizing to an HCV sequence in an analyte polynucleotide strand, wherein the oligomer is comprised of an HCV targeting sequence complementary to at least 4 contiguous nucleotides of HCV cDNA shown in Fig. 18

(b) incubating the analyte strand with the oligomer of (a) which allow specific hybrid duplexes to

form between the targeting sequence and the target sequence; and

(d) detecting hybrids formed between target region, if any, and the oligomer.

5 Yet another aspect of the invention is a method for preparing blood free of HCV comprising:

(a) providing analyte nucleic acids from a sample of blood suspected of containing an HCV target sequence;

10 (b) providing an oligomer capable of hybridizing to the HCV sequence in an analyte polynucleotide strand, if any, wherein the oligomer is comprised of an HCV targeting sequence complementary to a sequence of at least 8 nucleotides present in a conserved HCV nucleotide
15 sequence in HCV RNA;

(c) reacting (a) with (b) under conditions which allow the formation of a polynucleotide duplex between the targeting sequence and the target sequence, if any;

20 (d) detecting a duplex formed in (c), if any; and

(e) saving the blood from which complexes were not detected in (d).

25 Brief Description of the Drawings

Fig. 1 shows the sequence of the HCV cDNA in clone 12f, and the amino acids encoded therein.

Fig. 2 shows the HCV cDNA sequence in clone k9-1, and the amino acids encoded therein.

30 Fig. 3 shows the sequence of clone 15e, and the amino acids encoded therein.

Fig. 4 shows the nucleotide sequence of HCV cDNA in clone 13i, the amino acids encoded therein, and the sequences which overlap with clone 12f.

35

Fig. 5 shows the nucleotide sequence of HCV cDNA in clone 26j, the amino acids encoded therein, and the sequences which overlap clone 13i.

Fig. 6 shows the nucleotide sequence of HCV cDNA in clone CA59a, the amino acids encoded therein, and the sequences which overlap with clones 26j and K9-1.

Fig. 7 shows the nucleotide sequence of HCV cDNA in clone CA84a, the amino acids encoded therein, and the sequences which overlap with clone CA59a.

Fig. 8 shows the nucleotide sequence of HCV cDNA in clone CA156e, the amino acids encoded therein, and the sequences which overlap with CA84a.

Fig. 9 shows the nucleotide sequence of HCV cDNA in clone CA167b, the amino acids encoded therein, and the sequences which overlap CA156e.

Fig. 10 shows the nucleotide sequence of HCV cDNA in clone CA216a, the amino acids encoded therein, and the overlap with clone CA167b.

Fig. 11 shows the nucleotide sequence of HCV cDNA in clone CA290a, the amino acids encoded therein, and the overlap with clone CA216a.

Fig. 12 shows the nucleotide sequence of HCV cDNA in clone ag30a and the overlap with clone CA290a.

Fig. 13 shows the nucleotide sequence of HCV cDNA in clone CA205a, and the overlap with the HCV cDNA sequence in clone CA290a.

Fig. 14 shows the nucleotide sequence of HCV cDNA in clone 18g, and the overlap with the HCV cDNA sequence in clone ag30a.

Fig. 15 shows the nucleotide sequence of HCV cDNA in clone 16jh, the amino acids encoded therein, and the overlap of nucleotides with the HCV cDNA sequence in clone 15e.

Fig. 16 shows the nucleotide sequence of HCV cDNA in clone 6k, the amino acids encoded therein, and the

overlap of nucleotides with the HCV cDNA sequence in clone 16jh.

Fig. 17 shows the nucleotide sequence of HCV cDNA in clone p131jh, the amino acids encoded therein, and
5 the overlap of nucleotides with the HCV cDNA sequence in clone 6k.

Fig. 18 shows the the compiled HCV cDNA sequence derived from the clones described herein and from the compiled HCV cDNA sequence presented in E.P.O. Publication
10 No. 318,216. The clones from which the sequence was derived are 5'-clone32, b114a, 18g, ag30a, CA205a, CA290a, CA216a, p114a, CA167b, CA156e, CA84a, CA59a, K9-1 (also called k9-1), 26j, 13i, 12f, 14i, 11b, 7f, 7e, 8h, 33c, 40b, 37b, 35, 36, 81, 32, 33b, 25c, 14c, 8f, 33f, 33g,
15 39c, 35f, 19g, 26g, 15e, b5a, 16jh, 6k, and p131jh. In the figure the three horizontal dashes above the sequence indicate the position of the putative initiator methionine codon. Also shown in the figure is the amino acid sequence of the putative polyprotein encoded in the HCV
20 cDNA. Heterogeneities in cloned DNAs of HCV1 are indicated by the amino acids indicated above the putatively encoded sequence of the large ORF; the parentheses indicate that the heterogeneity was detected at or near to the 5'- or 3'- end of the HCV cDNA in the
25 clone.

Fig. 19 shows the sequences of capture and label probes for the detection of HCV RNA in biological samples.

Fig. 20 shows schematic alignment of a flaviviral polyprotein and a putative HCV polyprotein
30 encoded in the major ORF of the HCV genome. Also indicated in the figure are the possible functions of the flaviviral polypeptides cleaved from the flaviral polyprotein. In addition, the relative placements of the HCV polypeptides, NANB₅₋₁₋₁ and C100, with respect to the
35 putative HCV polyprotein are indicated.

Fig. 22 shows the double-stranded nucleotide sequence of the HCV cDNA insert in clone 81, and the putative amino acid sequence of the polypeptide encoded therein.

5 Fig. 23 shows the HCV cDNA sequence in clone 36, the segment which overlaps the NANBV cDNA of clone 81, and the polypeptide sequence encoded within clone 36.

 Fig. 24 shows the HCV cDNA sequence in clone 37b, the segment which overlaps clone 35, and the
10 polypeptide encoded therein.

 Fig. 25 shows autoradiographs of the HCV cPCR assay on RNA derived from liver samples of chimpanzees with NANBH (Fig. 25A) and on Italian patients with NANBH (Fig. 25B).

15 Fig. 26A and 26B are graphs showing the temporal relationship between the display of liver damage, the presence of HCV RNA, and the presence of anti-HCV antibodies for two chimpanzees with NANBH.

 Fig. 27 shows the nucleotide sequence of HCV
20 cDNA in clone CA84a, the amino acids encoded therein, and the sequences which overlap with clone CA59a.

 Fig. 28 shows the HCV cDNA sequence in clone 40b, the segment which overlaps clone 37b, and the polypeptide encoded therein.

25 Fig. 29 is an autoradiograph showing the labeled amplified products of approximately 300, 30, and 3 CID of HCV genomes.

 Fig. 32 shows the nucleotide sequence of HCV cDNA in clone 40a.

30 Fig. 33 is an autoradiograph showing amplified products extended from primers derived from conserved regions of the HCV genome.

 Fig. 34 shows the HCV cDNA sequence in clone 35, the segment which overlaps clone 36, and the polypeptide
35 encoded therein.

Fig. 37 is a diagram showing the relationship of probes and primers derived from the 5'-region of HCV RNA, from which the HCV cDNAs in clones ag30a and k9-1 are derived.

5 Fig. 38 is an autoradiograph of amplified products extended from sets of primers derived from ag30a and k9-1.

Fig. 39 shows the aligned nucleotide sequences of human isolates 23 and 27 and of HCV1. Homologous
10 sequences are indicated by the symbol (*). Non homologous sequences are in small letters.

Fig. 40 shows the aligned amino acid sequences of human isolates 23 and 27 and of HCV1. Homologous sequences are indicated by the symbol (*). Non homologous
15 sequences are in small letters.

Fig. 41 shows a half-tone reproduction of an autoradiograph of a Northern blot of RNA isolated from the liver of a BB-NANBV infected chimpanzee, probed with BB-NANBV cDNA of clone 81.

20 Fig. 43 shows a half-tone reproduction of an autoradiograph of nucleic acids extracted from NANBV particles captured from infected plasma with anti-NANB₅₋₁₋₁, and probed with ³²P-labeled NANBV cDNA from clone 81.

Fig. 44 shows reproductions of autoradiographs
25 of filters containing isolated NANBV nucleic acids, probed with ³²P-labeled plus and minus strand DNA probes derived from NANBV cDNA in clone 81.

Fig. 46 shows the nucleotide consensus sequence of human isolate 23, variant sequences are shown below the
30 sequence line. The amino acids encoded in the consensus sequence are also shown.

Fig. 47 shows the nucleotide consensus sequence of human isolate 27, variant sequences are shown below the
35 sequence line. The amino acids encoded in the consensus sequence are also shown.

Fig. 48 is a graph showing the relationship of the EnvL and EnvR primers to the model flavivirus polyprotein and putative HCV polyprotein.

Fig. 49 shows a comparison of the composite
5 aligned nucleotide sequences of isolates Thorn, EC1, HCT #18, and HCV1.

Fig. 50 shows a comparison of the nucleotide sequences of EC10 and a composite of the HCV1 sequence; the EC10 sequence is on the line above the dots, and the
10 HCV1 sequence is on the line below the dots.

Fig. 51 shows a comparison of the amino acid sequences 117-308 (relative to HCV1) encoded in the "EnvL" regions of the consensus sequences of human isolates HCT #18, JH23, JH 27, Thorne, EC1, and of HCV1.

Fig. 52 shows a comparison of the amino acid
15 sequences 330-360 (relative to HCV1) encoded in the "EnvR" regions of the consensus sequences of human isolates HCT #18, JH23, JH 27, Thorne, EC1, and of HCV1.

Fig. 53 shows the nucleotide sequences of
20 individual primers in primer mixture 5'-3.

Modes for Carrying Out the Invention

The term "hepatitis C virus" (HCV) has been reserved by workers in the field for an heretofore unknown
25 etiologic agent of NANBH. The prototype isolate of HCV has been identified in U.S.S.N. 122,714 (See also E.P.O. Publication No. 318,216). The term HCV also includes new isolates of the same viral species. As an extension of this terminology, the disease caused by HCV, formerly
30 called blood-borne NANB hepatitis (BB-NANBH), is called hepatitis C. The terms NANBH and hepatitis C may be used interchangeably herein.

HCV is a viral species of which pathogenic strains cause BB-NANBH. There may also be attenuated
35 strains or defective interfering particles derived therefrom. As shown infra, the HCV genome is comprised of

RNA. It is known that RNA containing viruses have relatively high rates of spontaneous mutation, i.e., reportedly on the order of 10^{-3} to 10^{-4} per incorporated nucleotide (Fields & Knipe (1986)). Therefore, since
5 heterogeneity and fluidity of genotype are inherent in RNA viruses, there are multiple strains/isolates, which may be virulent or avirulent, within the HCV species. The compositions and methods described herein, enable the propagation, identification, detection, and isolation of
10 the various HCV strains or isolates.

Several different strains/isolates of HCV have been identified. (See *infra*). One such strain or isolate, which is a prototype, is named CDC/HCV1 (also called HCV1). Information from one strain or isolate,
15 such as a partial genomic sequence, is sufficient to allow those skilled in the art using standard techniques to isolate new strains/isolates and to identify whether such new strains/isolates are HCV. For example, several different strains/isolates are described *infra*. These
20 strains, which were obtained from a number of human sera (and from different geographical areas), were isolated utilizing the information from the genomic sequence of HCV1.

Using the techniques described in E.P.O.
25 Publication No. 318,216 and *infra*, the genomic structure and the nucleotide sequence of HCV1 genomic RNA has been deduced. The genome appears to be single-stranded RNA containing ~10,000 nucleotides. The genome is positive-stranded, and possesses a continuous, translational open
30 reading frame (ORF) that encodes a polyprotein of about 3,000 amino acids. In the ORF, the structural protein(s) appear to be encoded in approximately the first quarter of the N-terminus region, with the majority of the polyprotein responsible for non-structural proteins. When
35 compared with all known viral sequences, small but significant co-linear homologies are observed with the

non-structural proteins of the flavivirus family, and with the pestiviruses (which are now also considered to be part of the Flavivirus family).

A schematic alignment of possible regions of a
5 flaviviral polyprotein (using Yellow Fever Virus as an example), and of a putative polyprotein encoded in the major ORF of the HCV genome, is shown in Fig. 20. In the figure the possible domains of the HCV polyprotein are indicated. The flavivirus polyprotein contains, from the
10 amino terminus to the carboxy terminus, the nucleocapsid protein (C), the matrix protein (M), the envelope protein (E), and the non-structural proteins (NS) 1, 2 (a+b), 3, 4 (a+b), and 5. Based upon the putative amino acids encoded in the nucleotide sequence of HCV1, a small domain at the
15 extreme N-terminus of the HCV polyprotein appears similar both in size and high content of basic residues to the nucleocapsid protein (C) found at the N-terminus of flaviviral polyproteins. The non-structural proteins 2,3,4, and 5 (NS2-5) of HCV and of yellow fever virus
20 (YFV) appear to have counter parts of similar size and hydropathicity, although there is divergence of the amino acid sequences. However, the region of HCV which would correspond to the regions of YFV polyprotein which contains the M, E, and NS1 protein not only differs in
25 sequence, but also appears to be quite different both in size and hydropathicity. Thus, while certain domains of the HCV genome may be referred to herein as, for example, NS1, or NS2, it should be borne in mind that these designations are speculative; there may be considerable
30 differences between the HCV family and flaviviruses that have yet to be appreciated.

Different strains, isolates or subtypes of HCV are expected to contain variations at the amino acid and nucleic acids compared with HCV1. Many isolates are
35 expected to show much (i.e., more than about 40%) homology in the total amino acid sequence compared with HCV1.

However, it may also be found that there are other less homologous HCV isolates. These would be defined as HCV according to various criteria such as, for example, an ORF of approximately 9,000 nucleotides to approximately 12,000 nucleotides, encoding a polyprotein similar in size to that of HCV1, an encoded polyprotein of similar hydrophobic and/or antigenic character to that of HCV1, and the presence of co-linear peptide sequences that are conserved with HCV1. In addition, it is believed that the genome would be a positive-stranded RNA.

All HCV isolates encode at least one epitope which is immunologically identifiable (i.e., immunologically cross-reactive) with an epitope encoded in the HCV cDNAs described herein. Preferably the epitope is contained in an amino acid sequence described herein and is unique to HCV when compared to previously known pathogens. The uniqueness of the epitope may be determined by its immunological reactivity with anti-HCV antibodies and lack of immunological reactivity with antibodies to known pathogens.

HCV strains and isolates are evolutionarily related. Therefore, it is expected that the overall homology of the genomes at the nucleotide level may be about 40% or greater, probably will be about 50% or greater, probably about 60% or greater, and even more probably about 80% or greater; and in addition that there will be corresponding contiguous sequences of at least about 13 nucleotides. It should be noted, as shown infra, that there are variable and hypervariable regions within the HCV genome; therefore, the homology in these regions is expected to be significantly less than that in the overall genome. The correspondence between the putative HCV strain genomic sequence and, for example, the CDC/HCV1 cDNA sequence can be determined by techniques known in the art. For example, they can be determined by a direct comparison of the sequence information of the

polynucleotide from the putative HCV, and the HCV cDNA sequence(s) described herein. They also can be determined by hybridization of the polynucleotides under conditions which form stable duplexes between homologous regions (for example, those which would be used prior to S_1 digestion), followed by digestion with single stranded specific nuclease(s), followed by size determination of the digested fragments.

Because of the evolutionary relationship of the strains or isolates of HCV, putative HCV strains or isolates are identifiable by their homology at the polypeptide level. Generally, HCV strains or isolates are expected to be at least 40% homologous, more than about 50% homologous, probably more than about 70% homologous, and even more probably more than about 80% homologous, and some may even be more than about 90% homologous at the polypeptide level. The techniques for determining amino acid sequence homology are known in the art. For example, the amino acid sequence may be determined directly and compared to the sequences provided herein. Alternatively the nucleotide sequence of the genomic material of the putative HCV may be determined (usually via a cDNA intermediate), the putative amino acid sequence encoded therein can be determined, and the corresponding regions compared.

As used herein, a polynucleotide "derived from" a designated sequence refers to a polynucleotide sequence which is comprised of a sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10-12 nucleotides, and even more preferably at least about 15-20 nucleotides corresponding to a region of the designated nucleotide sequence. "Corresponding" means homologous to or complementary to the designated sequence. Preferably, the sequence of the region from which the polynucleotide is derived is homologous to or complementary to a sequence which is unique to an HCV genome. More preferably, the

derived sequence is homologous or complementary to a sequence that is unique to all or to a majority of HCV isolates. Whether or not a sequence is unique to the HCV genome can be determined by techniques known to those of skill in the art. For example, the sequence can be compared to sequences in databanks, e.g., Genbank, to determine whether it is present in the uninfected host or other organisms. The sequence can also be compared to the known sequences of other viral agents, including those which are known to induce hepatitis, e.g., HAV, HBV, and HDV, and to members of the Flaviviridae. The correspondence or non-correspondence of the derived sequence to other sequences can also be determined by hybridization under the appropriate stringency conditions. Hybridization techniques for determining the complementarity of nucleic acid sequences are known in the art, and are discussed infra. See also, for example, Maniatis et al. (1982). In addition, mismatches of duplex polynucleotides formed by hybridization can be determined by known techniques, including for example, digestion with a nuclease such as S1 that specifically digests single-stranded areas in duplex polynucleotides. Regions from which typical DNA sequences may be "derived" include but are not limited to, for example, regions encoding specific epitopes, as well as non-transcribed and/or non-translated regions.

The derived polynucleotide is not necessarily physically derived from the nucleotide sequence shown, but may be generated in any manner, including for example, chemical synthesis or DNA replication or reverse transcription or transcription. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with an intended use.

The term "recombinant polynucleotide" as used herein intends a polynucleotide of genomic, cDNA,

semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than
5 that to which it is linked in nature, or (3) does not occur in nature.

The term "polynucleotide" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers
10 only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA and RNA. It also includes known types of modifications, for example, labels which are known in the art, methylation, "caps", substitution of one or more of the naturally occurring
15 nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those
20 containing pendant moieties, such as, for example proteins (including for e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative
25 metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide.

As used herein, the "sense strand" of a nucleic
30 acid contains the sequence that has sequence homology to that of mRNA. The "anti-sense strand" contains a sequence which is complementary to that of the "sense strand".

As used herein, a "positive stranded genome" of a virus is one in which the genome, whether RNA or DNA, is
35 single-stranded and which encodes a viral polypeptide(s). Examples of positive stranded RNA viruses include

Togaviridae, Coronaviridae, Retroviridae, Picornaviridae, and Caliciviridae. Included also, are the Flaviviridae, which were formerly classified as Togaviridae. See Fields & Knipe (1986).

5 The term "primer" as used herein refers to an oligomer which is capable of acting as a point of initiation of synthesis of a polynucleotide strand when placed under appropriate conditions. The primer will be completely or substantially complementary to a region of
10 the polynucleotide strand to be copied. Thus, under conditions conducive to hybridization, the primer will anneal to the complementary region of the analyte strand. Upon addition of suitable reactants, (e.g., a polymerase, nucleotide triphosphates, and the like), the primer is
15 extended by the polymerizing agent to form a copy of the analyte strand. The primer may be single-stranded, or alternatively may be partially or fully double-stranded.

 The terms "analyte polynucleotide" and "analyte strand" refer to a single- or double-stranded nucleic acid
20 molecule which is suspected of containing a target sequence, and which may be present in a biological sample.

 As used herein, the term "oligomer" refers to primers and to probes. The term oligomer does not connote the size of the molecule. However, typically oligomers
25 are no greater than 1000 nucleotides, more typically are no greater than 500 nucleotides, even more typically are no greater than 250 nucleotides; they may be no greater than 100 nucleotides, and may be no greater than 75
30 nucleotides, and also may be no greater than 50 nucleotides in length.

 As used herein, the term "probe" refers to a structure comprised of a polynucleotide which forms a hybrid structure with a target sequence, due to complementarity of at least one sequence in the probe with
35 a sequence in the target region. The polynucleotide regions of probes may be composed of DNA, and/or RNA, and/

or synthetic nucleotide analogs. Included within probes are "capture probes" and "label probes". Preferably the probe does not contain a sequence complementary to sequence(s) used to prime the polymerase chain reaction
5 (PCR).

As used herein, the term "target region" refers to a region of the nucleic acid which is to be amplified and/or detected. The term "target sequence" refers to a sequence with which a probe or primer will form a stable
10 hybrid under desired conditions.

The term "capture probe" as used herein refers to a polynucleotide comprised of a single-stranded polynucleotide coupled to a binding partner. The single-stranded polynucleotide is comprised of a targeting
15 polynucleotide sequence, which is complementary to a target sequence in a target region to be detected in the analyte polynucleotide. This complementary region is of sufficient length and complementarity to the target sequence to afford a duplex of stability which is suf-
20 ficient to immobilize the analyte polynucleotide to a solid surface (via the binding partners). The binding partner is specific for a second binding partner; the second binding partner can be bound to the surface of a solid support, or may be linked indirectly via other
25 structures or binding partners to a solid support.

The term "targeting polynucleotide sequence" as used herein, refers to a polynucleotide sequence which is comprised of nucleotides which are complementary to a target nucleotide sequence; the sequence is of sufficient
30 length and complementarity with the target sequence to form a duplex which has sufficient stability for the purpose intended.

The term "binding partner" as used herein refers to a molecule capable of binding a ligand molecule with
35 high specificity, as for example an antigen and an antibody specific therefor. In general, the specific binding

partners must bind with sufficient affinity to immobilize the analyte copy/complementary strand duplex (in the case of capture probes) under the isolation conditions.

Specific binding partners are known in the art, and include, for example, biotin and avidin or streptavidin, IgG and protein A, the numerous known receptor-ligand couples, and complementary polynucleotide strands. In the case of complementary polynucleotide binding partners, the partners are normally at least about 15 bases in length, and may be at least 40 bases in length; in addition, they have a content of Gs and Cs of at least about 40% and as much as about 60%. The polynucleotides may be composed of DNA, RNA, or synthetic nucleotide analogs.

The term "coupled" as used herein refers to attachment by covalent bonds or by strong non-covalent interactions (e.g., hydrophobic interactions, hydrogen bonds, etc.). Covalent bonds may be, for example, ester, ether, phosphoester, amide, peptide, imide, carbon-sulfur bonds, carbon-phosphorus bonds, and the like.

The term "support" refers to any solid or semi-solid surface to which a desired binding partner may be anchored. Suitable supports include glass, plastic, metal, polymer gels, and the like, and may take the form of beads, wells, dipsticks, membranes, and the like.

The term "label" as used herein refers to any atom or moiety which can be used to provide a detectable (preferably quantifiable) signal, and which can be attached to a polynucleotide or polypeptide.

As used herein, the term "label probe" refers to an oligomer which is comprised of targeting polynucleotide sequence, which is complementary to a target sequence to be detected in the analyte polynucleotide. This complementary region is of sufficient length and complementarity to the target sequence to afford a duplex comprised of the "label probe" and the "target sequence" to be detected by the label. The oligomer is coupled to a

label either directly, or indirectly via a set of ligand molecules with high specificity for each other. Sets of ligand molecules with high specificity are described supra., and also includes multimers.

5 The term "multimer", as used herein, refers to linear or branched polymers of the same repeating single-stranded polynucleotide unit or different single-stranded polynucleotide units. At least one of the units has a sequence, length, and composition that permits it to
10 hybridize specifically to a first single-stranded nucleotide sequence of interest, typically an analyte or an oligomer (e.g., a label probe) bound to an analyte. In order to achieve such specificity and stability, this unit will normally be at least about 15 nucleotides in length,
15 typically no more than about 50 nucleotides in length, and preferably about 30 nucleotides in length; moreover, the content of Gs and Cs will normally be at least about 40%, and at most about 60%. In addition to such unit(s), the multimer includes a multiplicity of units that are capable
20 of hybridizing specifically and stably to a second single-stranded nucleotide of interest, typically a labeled polynucleotide or another multimer. These units are generally about the same size and composition as the multimers discussed above. When a multimer is designed to
25 be hybridized to another multimer, the first and second oligonucleotide units are heterogeneous (different), and do not hybridize with each other under the conditions of the selected assay. Thus, multimers may be label probes, or may be ligands which couple the label to the probe.

30 As used herein, the term "viral RNA", which includes HCV RNA, refers to RNA from the viral genome, fragments thereof, transcripts thereof, and mutant sequences derived therefrom.

 As used herein, a "biological sample" refers to
35 a sample of tissue or fluid isolated from an individual, including but not limited to, for example, plasma, serum,

spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs, and also samples of in vitro cell culture constituents (including but not limited to conditioned medium resulting from the growth of cells in cell culture medium, putatively virally infected cells, recombinant cells, and cell components).

Description of the Invention

10 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in
15 the literature. See e.g., Maniatis, Fitch & Sambrook, MOLECULAR CLONING; A LABORATORY MANUAL (1982); DNA CLONING, VOLUMES I AND II (D.N Glover ed. 1985); OLIGONUCLEOTIDE SYNTHESIS (M.J. Gait ed, 1984); NUCLEIC ACID HYBRIDIZATION (B.D. Hames & S.J. Higgins eds. 1984);
20 the series, METHODS IN ENZYMOLOGY (Academic Press, Inc.), particularly Vol. 154 and Vol. 155 (Wu and Grossman, and Wu, eds., respectively). All patents, patent applications, and publications mentioned herein, both supra and infra, are hereby incorporated herein by reference.

25 The useful materials and processes of the present invention are made possible by the identification of HCV as the etiologic agent of BB-NANBV, and by the provision of a family of nucleotide sequences isolated from cDNA libraries which contain HCV cDNA sequences.
30 These cDNA libraries were derived from nucleic acid sequences present in the plasma of an HCV-infected chimpanzee. The construction of one of these libraries, the "c" library (ATCC No. 40394), is described in E.P.O. Publication No. 318,216.

35 Utilizing the above-described HCV cDNA sequences, as well as that described herein, oligomers can

be constructed which are useful as reagents for detecting viral polynucleotides in biological samples. For example, from the sequences it is possible to synthesize DNA oligomers of about 8-10 nucleotides, or larger, which are useful as hybridization probes to detect the presence of HCV RNA in, for example, donated blood, blood fractions, sera of subjects suspected of harboring the virus, or cell culture systems in which the virus is replicating. In addition, the novel oligomers described herein enable further characterization of the HCV genome. Polynucleotide probes and primers derived from these sequences may be used to amplify sequences present in cDNA libraries, and/or to screen cDNA libraries for additional overlapping cDNA sequences, which, in turn, may be used to obtain more overlapping sequences. As indicated infra. and in E.P.O. Publication No. 318,216, the genome of HCV appears to be RNA comprised primarily of a large open reading frame (ORF) which encodes a large polyprotein.

In addition to the above, the information provided infra allows the identification of additional HCV strains or isolates. The isolation and characterization of the additional HCV strains or isolates may be accomplished utilizing techniques known to those of skill in the art, for example, by isolating the nucleic acids from body components which contain viral particles and/or viral RNA, creating cDNA libraries using the oligomers described infra., for screening the libraries for clones containing HCV cDNA sequences described infra., and comparing the HCV cDNAs from the new isolates with the cDNAs described in E.P.O. Publication No. 318,216 and infra. Strains or isolates which fit within the parameters of HCV, as described in the Definitions section, supra., are readily identifiable. Other methods for identifying HCV strains will be obvious to those of skill in the art, based upon the information provided herein.

Isolation of the HCV cDNA Sequences

The oligomers of the invention contain regions which form hybrid duplex structures with targeted sequences in HCV polynucleotides. The HCV polynucleotide hybridizing regions of the oligomers may be ascertained from the HCV cDNA sequence(s) provided herein, and described in E.P.O. Publication No. 318,216. A composite of HCV cDNA from HCV1, a prototypic HCV, is shown in Fig. 18. The composite sequence is based upon sequence information derived from a number of HCV cDNA clones, which were isolated from a number of HCV cDNA libraries, including the "c" library present in lambda gt11 (ATCC No. 40394), and from human serum. The HCV cDNA clones were isolated by methods described in E.P.O. Publication No. 318,216. Briefly, the majority of clones which were isolated contained sequences from the HCV cDNA "c" library which was constructed using pooled serum from a chimpanzee with chronic HCV infection and containing a high titer of the virus, i.e., at least 10^6 chimp infectious doses/ml (CID/ml). The pooled serum was used to isolate viral particles; nucleic acids isolated from these particles was used as the template in the construction of cDNA libraries to the viral genome. The initial clone, 5-1-1, was obtained by screening the "c" library with serum from infected individuals. After the isolation of the initial clone, the remainder of the sequence was obtained by screening with synthetic polynucleotide probes, the sequences of which were derived from the 5'-region and the 3'-region of the known HCV cDNA sequence(s).

The description of the methods to retrieve the cDNA sequences is mostly of historical interest. The resultant sequences (and their complements) are provided herein, and the sequences, or any portion thereof, could be prepared using synthetic methods, or by a combination of synthetic methods with retrieval of partial sequences

using methods similar to those described in E.P.O.
Publication No. 318,216.

Oligomer Probes and Primers

- 5 Using as a basis the HCV genome (as illustrated
in Fig. 18), and/or preferably conserved regions of the
HCV genome, oligomers of approximately 8 nucleotides or
more can be prepared which hybridize with the positive
strand(s) of HCV RNA or its complement, as well as to HCV
10 cDNAs. These oligomers can serve as probes for the detec-
tion (including isolation and/or labeling) of
polynucleotides which contain HCV nucleotide sequences,
and/or as primers for the transcription and/or replication
of targeted HCV sequences. The oligomers contain a
15 targeting polynucleotide sequence, which is comprised of
nucleotides which are complementary to a target HCV
nucleotide sequence; the sequence is of sufficient length
and complementarity with the HCV sequence to form a duplex
which has sufficient stability for the purpose intended.
20 For example, if the purpose is the isolation, via im-
mobilization, of an analyte containing a target HCV
sequence, the oligomers would contain a polynucleotide
region which is of sufficient length and complementarity
to the targeted HCV sequence to afford sufficient duplex
25 stability to immobilize the analyte on a solid surface,
via its binding to the oligomers, under the isolation
conditions. For example, also, if the oligomers are to
serve as primers for the transcription and/or replication
of target HCV sequences in an analyte polynucleotide, the
30 oligomers would contain a polynucleotide region of suf-
ficient length and complementarity to the targeted HCV
sequence to allow the polymerizing agent to continue
replication from the primers which are in stable duplex
form with the target sequence, under the polymerizing
35 conditions. For example, also, if the oligomers are to be
used as label probes, or are to bind to multimers, the

targeting polynucleotide region would be of sufficient length and complementarity to form stable hybrid duplex structures with the label probes and/or multimers to allow detection of the duplex. The oligomers may contain a
 5 minimum of about 4 contiguous nucleotides which are complementary to targeted HCV sequence; usually the oligomers will contain a minimum of about 8 contiguous nucleotides which are complementary to the targeted HCV sequence, and preferably will contain a minimum of about 14 contiguous
 10 nucleotides which are complementary to the targeted HCV sequence.

Suitable HCV nucleotide targeting sequences may be comprised of nucleotides which are complementary nucleotides selected from the following HCV cDNA
 15 nucleotides, which are shown in Fig. 18, ($nn_x - nn_y$ denotes from about nucleotide number x to about nucleotide number y):

20 $nn_{-340} - nn_{-330}$; $nn_{-330} - nn_{-320}$; $nn_{-320} - nn_{-310}$;
 $nn_{-310} - nn_{-300}$; $nn_{-300} - nn_{-290}$; $nn_{-290} - nn_{-280}$;
 $nn_{-280} - nn_{-270}$; $nn_{-270} - nn_{-260}$; $nn_{-260} - nn_{-250}$;
 $nn_{-250} - nn_{-240}$; $nn_{-240} - nn_{-230}$; $nn_{-230} - nn_{-220}$;
 $nn_{-220} - nn_{-210}$; $nn_{-210} - nn_{-200}$; $nn_{-200} - nn_{-190}$;
 $nn_{-190} - nn_{-180}$; $nn_{-180} - nn_{-170}$; $nn_{-170} - nn_{-160}$;
 25 $nn_{-160} - nn_{-150}$; $nn_{-150} - nn_{-140}$; $nn_{-140} - nn_{-130}$;
 $nn_{-130} - nn_{-120}$; $nn_{-120} - nn_{-110}$; $nn_{-110} - nn_{-100}$;
 $nn_{-100} - nn_{-90}$; $nn_{-90} - nn_{-80}$; $nn_{-80} - nn_{-70}$;
 $nn_{-70} - nn_{-60}$; $nn_{-60} - nn_{-50}$; $nn_{-50} - nn_{-40}$;
 $nn_{-40} - nn_{-30}$; $nn_{-30} - nn_{-20}$; $nn_{-20} - nn_{-10}$;
 30 $nn_{-10} - nn_1$; $nn_1 - nn_{10}$; $nn_{10} - nn_{20}$; $nn_{20} - nn_{30}$;
 $nn_{30} - nn_{40}$; $nn_{40} - nn_{50}$; $nn_{50} - nn_{60}$; $nn_{60} - nn_{70}$;
 $nn_{70} - nn_{80}$; $nn_{80} - nn_{90}$; $nn_{90} - nn_{100}$; $nn_{100} - nn_{110}$;
 $nn_{110} - nn_{120}$; $nn_{120} - nn_{130}$; $nn_{130} - nn_{140}$;
 $nn_{140} - nn_{150}$; $nn_{150} - nn_{160}$; $nn_{160} - nn_{170}$;
 35 $nn_{170} - nn_{180}$; $nn_{180} - nn_{190}$; $nn_{190} - nn_{200}$;
 $nn_{200} - nn_{210}$; $nn_{210} - nn_{220}$; $nn_{220} - nn_{230}$;

nn230 - nn240; nn240 - nn250; nn250 - nn260;
nn260 - nn270; nn270 - nn280; nn280 - nn290;
nn290 - nn300; nn300 - nn310; nn310 - nn320;
nn320 - nn330; nn330 - nn340; nn340 - nn350;
5 nn350 - nn360; nn360 - nn370; nn370 - nn380;
nn380 - nn390; nn390 - nn400; nn400 - nn410;
nn410 - nn420; nn420 - nn430; nn430 - nn440;
nn440 - nn450; nn450 - nn460; nn460 - nn470;
nn470 - nn480; nn480 - nn490; nn490 - nn500;
10 nn500 - nn510; nn510 - nn520; nn520 - nn530;
nn530 - nn540; nn540 - nn550; nn550 - nn560;
nn560 - nn570; nn570 - nn580; nn580 - nn590;
nn590 - nn600; nn600 - nn610; nn610 - nn620;
nn620 - nn630; nn630 - nn640; nn640 - nn650;
15 nn650 - nn660; nn660 - nn670; nn670 - nn680;
nn680 - nn690; nn690 - nn700; nn700 - nn710;
nn710 - nn720; nn720 - nn730; nn730 - nn740;
nn740 - nn750; nn750 - nn760; nn760 - nn770;
nn770 - nn780; nn780 - nn790; nn790 - nn800;
20 nn800 - nn810; nn810 - nn820; nn820 - nn830;
nn830 - nn840; nn840 - nn850; nn850 - nn860;
nn860 - nn870; nn870 - nn880; nn880 - nn890;
nn890 - nn900; nn900 - nn910; nn910 - nn920;
nn920 - nn930; nn930 - nn940; nn940 - nn950;
25 nn950 - nn960; nn960 - nn970; nn970 - nn980;
nn980 - nn990; nn990 - nn1000; nn1000 - nn1010;
nn1010 - nn1020; nn1020 - nn1030; nn1030 - nn1040;
nn1040 - nn1050; nn1050 - nn1060; nn1060 - nn1070;
nn1070 - nn1080; nn1080 - nn1090; nn1090 - nn1100;
30 nn1100 - nn1110; nn1110 - nn1120; nn1120 - nn1130;
nn1130 - nn1140; nn1140 - nn1150; nn1150 - nn1160;
nn1160 - nn1170; nn1170 - nn1180; nn1180 - nn1190;
nn1190 - nn1200; nn1200 - nn1210; nn1210 - nn1220;
nn1220 - nn1230; nn1230 - nn1240; nn1240 - nn1250;
35 nn1250 - nn1260; nn1260 - nn1270; nn1270 - nn1280;
nn1280 - nn1290; nn1290 - nn1300; nn1300 - nn1310;

nn1310 - nn1320; nn1320 - nn1330; nn1330 - nn1340;
nn1340 - nn1350; nn1350 - nn1360; nn1360 - nn1370;
nn1370 - nn1380; nn1380 - nn1390; nn1390 - nn1400;
nn1400 - nn1410; nn1410 - nn1420; nn1420 - nn1430;
5 nn1430 - nn1440; nn1440 - nn1450; nn1450 - nn1460;
nn1460 - nn1470; nn1470 - nn1480; nn1480 - nn1490;
nn1490 - nn1500; nn1500 - nn1510; nn1510 - nn1520;
nn1520 - nn1530; nn1530 - nn1540; nn1540 - nn1550;
nn1550 - nn1560; nn1560 - nn1570; nn1570 - nn1580;
10 nn1580 - nn1590; nn1590 - nn1600; nn1600 - nn1610;
nn1610 - nn1620; nn1620 - nn1630; nn1630 - nn1640;
nn1640 - nn1650; nn1650 - nn1660; nn1660 - nn1670;
nn1670 - nn1680; nn1680 - nn1690; nn1690 - nn1700;
nn1700 - nn1710; nn1710 - nn1720; nn1720 - nn1730;
15 nn1730 - nn1740; nn1740 - nn1750; nn1750 - nn1760;
nn1760 - nn1770; nn1770 - nn1780; nn1780 - nn1790;
nn1790 - nn1800; nn1800 - nn1810; nn1810 - nn1820;
nn1820 - nn1830; nn1830 - nn1840; nn1840 - nn1850;
nn1850 - nn1860; nn1860 - nn1870; nn1870 - nn1880;
20 nn1880 - nn1890; nn1890 - nn1900; nn1900 - nn1910;
nn1910 - nn1920; nn1920 - nn1930; nn1930 - nn1940;
nn1940 - nn1950; nn1950 - nn1960; nn1960 - nn1970;
nn1970 - nn1980; nn1980 - nn1990; nn1990 - nn2000;
nn2000 - nn2010; nn2010 - nn2020; nn2020 - nn2030;
25 nn2030 - nn2040; nn2040 - nn2050; nn2050 - nn2060;
nn2060 - nn2070; nn2070 - nn2080; nn2080 - nn2090;
nn2090 - nn2100; nn2100 - nn2110; nn2110 - nn2120;
nn2120 - nn2130; nn2130 - nn2140; nn2140 - nn2150;
nn2150 - nn2160; nn2160 - nn2170; nn2170 - nn2180;
30 nn2180 - nn2190; nn2190 - nn2200; nn2200 - nn2210;
nn2210 - nn2220; nn2220 - nn2230; nn2230 - nn2240;
nn2240 - nn2250; nn2250 - nn2260; nn2260 - nn2270;
nn2270 - nn2280; nn2280 - nn2290; nn2290 - nn2300;
nn2300 - nn2310; nn2310 - nn2320; nn2320 - nn2330;
35 nn2330 - nn2340; nn2340 - nn2350; nn2350 - nn2360;
nn2360 - nn2370; nn2370 - nn2380; nn2380 - nn2390;

nn2390 - nn2400; nn2400 - nn2410; nn2410 - nn2420;
nn2420 - nn2430; nn2430 - nn2440; nn2440 - nn2450;
nn2450 - nn2460; nn2460 - nn2470; nn2470 - nn2480;
nn2480 - nn2490; nn2490 - nn2500; nn2500 - nn2510;
5 nn2510 - nn2520; nn2520 - nn2530; nn2530 - nn2540;
nn2540 - nn2550; nn2550 - nn2560; nn2560 - nn2570;
nn2570 - nn2580; nn2580 - nn2590; nn2590 - nn2600;
nn2600 - nn2610; nn2610 - nn2620; nn2620 - nn2630;
nn2630 - nn2640; nn2640 - nn2650; nn2650 - nn2660;
10 nn2660 - nn2670; nn2670 - nn2680; nn2680 - nn2690;
nn2690 - nn2700; nn2700 - nn2710; nn2710 - nn2720;
nn2720 - nn2730; nn2730 - nn2740; nn2740 - nn2750;
nn2750 - nn2760; nn2760 - nn2770; nn2770 - nn2780;
nn2780 - nn2790; nn2790 - nn2800; nn2800 - nn2810;
15 nn2810 - nn2820; nn2820 - nn2830; nn2830 - nn2840;
nn2840 - nn2850; nn2850 - nn2860; nn2860 - nn2870;
nn2870 - nn2880; nn2880 - nn2890; nn2890 - nn2900;
nn2900 - nn2910; nn2910 - nn2920; nn2920 - nn2930;
nn2930 - nn2940; nn2940 - nn2950; nn2950 - nn2960;
20 nn2960 - nn2970; nn2970 - nn2980; nn2980 - nn2990;
nn2990 - nn3000; nn3000 - nn3010; nn3010 - nn3020;
nn3020 - nn3030; nn3030 - nn3040; nn3040 - nn3050;
nn3050 - nn3060; nn3060 - nn3070; nn3070 - nn3080;
nn3080 - nn3090; nn3090 - nn3100; nn3100 - nn3110;
25 nn3110 - nn3120; nn3120 - nn3130; nn3130 - nn3140;
nn3140 - nn3150; nn3150 - nn3160; nn3160 - nn3170;
nn3170 - nn3180; nn3180 - nn3190; nn3190 - nn3200;
nn3200 - nn3210; nn3210 - nn3220; nn3220 - nn3230;
nn3230 - nn3240; nn3240 - nn3250; nn3250 - nn3260;
30 nn3260 - nn3270; nn3270 - nn3280; nn3280 - nn3290;
nn3290 - nn3300; nn3300 - nn3310; nn3310 - nn3320;
nn3320 - nn3330; nn3330 - nn3340; nn3340 - nn3350;
nn3350 - nn3360; nn3360 - nn3370; nn3370 - nn3380;
nn3380 - nn3390; nn3390 - nn3400; nn3400 - nn3410;
35 nn3410 - nn3420; nn3420 - nn3430; nn3430 - nn3440;
nn3440 - nn3450; nn3450 - nn3460; nn3460 - nn3470;

nn3470 - nn3480; nn3480 - nn3490; nn3490 - nn3500;
nn3500 - nn3510; nn3510 - nn3520; nn3520 - nn3530;
nn3530 - nn3540; nn3540 - nn3550; nn3550 - nn3560;
nn3560 - nn3570; nn3570 - nn3580; nn3580 - nn3590;
5 nn3590 - nn3600; nn3600 - nn3610; nn3610 - nn3620;
nn3620 - nn3630; nn3630 - nn3640; nn3640 - nn3650;
nn3650 - nn3660; nn3660 - nn3670; nn3670 - nn3680;
nn3680 - nn3690; nn3690 - nn3700; nn3700 - nn3710;
nn3710 - nn3720; nn3720 - nn3730; nn3730 - nn3740;
10 nn3740 - nn3750; nn3750 - nn3760; nn3760 - nn3770;
nn3770 - nn3780; nn3780 - nn3790; nn3790 - nn3800;
nn3800 - nn3810; nn3810 - nn3820; nn3820 - nn3830;
nn3830 - nn3840; nn3840 - nn3850; nn3850 - nn3860;
nn3860 - nn3870; nn3870 - nn3880; nn3880 - nn3890;
15 nn3890 - nn3900; nn3900 - nn3910; nn3910 - nn3920;
nn3920 - nn3930; nn3930 - nn3940; nn3940 - nn3950;
nn3950 - nn3960; nn3960 - nn3970; nn3970 - nn3980;
nn3980 - nn3990; nn3990 - nn4000; nn4000 - nn4010;
nn4010 - nn4020; nn4020 - nn4030; nn4030 - nn4040;
20 nn4040 - nn4050; nn4050 - nn4060; nn4060 - nn4070;
nn4070 - nn4080; nn4080 - nn4090; nn4090 - nn4100;
nn4100 - nn4110; nn4110 - nn4120; nn4120 - nn4130;
nn4130 - nn4140; nn4140 - nn4150; nn4150 - nn4160;
nn4160 - nn4170; nn4170 - nn4180; nn4180 - nn4190;
25 nn4190 - nn4200; nn4200 - nn4210; nn4210 - nn4220;
nn4220 - nn4230; nn4230 - nn4240; nn4240 - nn4250;
nn4250 - nn4260; nn4260 - nn4270; nn4270 - nn4280;
nn4280 - nn4290; nn4290 - nn4300; nn4300 - nn4310;
nn4310 - nn4320; nn4320 - nn4330; nn4330 - nn4340;
30 nn4340 - nn4350; nn4350 - nn4360; nn4360 - nn4370;
nn4370 - nn4380; nn4380 - nn4390; nn4390 - nn4400;
nn4400 - nn4410; nn4410 - nn4420; nn4420 - nn4430;
nn4430 - nn4440; nn4440 - nn4450; nn4450 - nn4460;
nn4460 - nn4470; nn4470 - nn4480; nn4480 - nn4490;
35 nn4490 - nn4500; nn4500 - nn4510; nn4510 - nn4520;
nn4520 - nn4530; nn4530 - nn4540; nn4540 - nn4550;

nn4550 - nn4560; nn4560 - nn4570; nn4570 - nn4580;
nn4580 - nn4590; nn4590 - nn4600; nn4600 - nn4610;
nn4610 - nn4620; nn4620 - nn4630; nn4630 - nn4640;
nn4640 - nn4650; nn4650 - nn4660; nn4660 - nn4670;
5 nn4670 - nn4680; nn4680 - nn4690; nn4690 - nn4700;
nn4700 - nn4710; nn4710 - nn4720; nn4720 - nn4730;
nn4730 - nn4740; nn4740 - nn4750; nn4750 - nn4760;
nn4760 - nn4770; nn4770 - nn4780; nn4780 - nn4790;
nn4790 - nn4800; nn4800 - nn4810; nn4810 - nn4820;
10 nn4820 - nn4830; nn4830 - nn4840; nn4840 - nn4850;
nn4850 - nn4860; nn4860 - nn4870; nn4870 - nn4880;
nn4880 - nn4890; nn4890 - nn4900; nn4900 - nn4910;
nn4910 - nn4920; nn4920 - nn4930; nn4930 - nn4940;
nn4940 - nn4950; nn4950 - nn4960; nn4960 - nn4970;
15 nn4970 - nn4980; nn4980 - nn4990; nn4990 - nn5000;
nn5000 - nn5010; nn5010 - nn5020; nn5020 - nn5030;
nn5030 - nn5040; nn5040 - nn5050; nn5050 - nn5060;
nn5060 - nn5070; nn5070 - nn5080; nn5080 - nn5090;
nn5090 - nn5100; nn5100 - nn5110; nn5110 - nn5120;
20 nn5120 - nn5130; nn5130 - nn5140; nn5140 - nn5150;
nn5150 - nn5160; nn5160 - nn5170; nn5170 - nn5180;
nn5180 - nn5190; nn5190 - nn5200; nn5200 - nn5210;
nn5210 - nn5220; nn5220 - nn5230; nn5230 - nn5240;
nn5240 - nn5250; nn5250 - nn5260; nn5260 - nn5270;
25 nn5270 - nn5280; nn5280 - nn5290; nn5290 - nn5300;
nn5300 - nn5310; nn5310 - nn5320; nn5320 - nn5330;
nn5330 - nn5340; nn5340 - nn5350; nn5350 - nn5360;
nn5360 - nn5370; nn5370 - nn5380; nn5380 - nn5390;
nn5390 - nn5400; nn5400 - nn5410; nn5410 - nn5420;
30 nn5420 - nn5430; nn5430 - nn5440; nn5440 - nn5450;
nn5450 - nn5460; nn5460 - nn5470; nn5470 - nn5480;
nn5480 - nn5490; nn5490 - nn5500; nn5500 - nn5510;
nn5510 - nn5520; nn5520 - nn5530; nn5530 - nn5540;
nn5540 - nn5550; nn5550 - nn5560; nn5560 - nn5570;
35 nn5570 - nn5580; nn5580 - nn5590; nn5590 - nn5600;
nn5600 - nn5610; nn5610 - nn5620; nn5620 - nn5630;

nn5630 - nn5640; nn5640 - nn5650; nn5650 - nn5660;
nn5660 - nn5670; nn5670 - nn5680; nn5680 - nn5690;
nn5690 - nn5700; nn5700 - nn5710; nn5710 - nn5720;
nn5720 - nn5730; nn5730 - nn5740; nn5740 - nn5750;
5 nn5750 - nn5760; nn5760 - nn5770; nn5770 - nn5780;
nn5780 - nn5790; nn5790 - nn5800; nn5800 - nn5810;
nn5810 - nn5820; nn5820 - nn5830; nn5830 - nn5840;
nn5840 - nn5850; nn5850 - nn5860; nn5860 - nn5870;
nn5870 - nn5880; nn5880 - nn5890; nn5890 - nn5900;
10 nn5900 - nn5910; nn5910 - nn5920; nn5920 - nn5930;
nn5930 - nn5940; nn5940 - nn5950; nn5950 - nn5960;
nn5960 - nn5970; nn5970 - nn5980; nn5980 - nn5990;
nn5990 - nn6000; nn6000 - nn6010; nn6010 - nn6020;
nn6020 - nn6030; nn6030 - nn6040; nn6040 - nn6050;
15 nn6050 - nn6060; nn6060 - nn6070; nn6070 - nn6080;
nn6080 - nn6090; nn6090 - nn6100; nn6100 - nn6110;
nn6110 - nn6120; nn6120 - nn6130; nn6130 - nn6140;
nn6140 - nn6150; nn6150 - nn6160; nn6160 - nn6170;
nn6170 - nn6180; nn6180 - nn6190; nn6190 - nn6200;
20 nn6200 - nn6210; nn6210 - nn6220; nn6220 - nn6230;
nn6230 - nn6240; nn6240 - nn6250; nn6250 - nn6260;
nn6260 - nn6270; nn6270 - nn6280; nn6280 - nn6290;
nn6290 - nn6300; nn6300 - nn6310; nn6310 - nn6320;
nn6320 - nn6330; nn6330 - nn6340; nn6340 - nn6350;
25 nn6350 - nn6360; nn6360 - nn6370; nn6370 - nn6380;
nn6380 - nn6390; nn6390 - nn6400; nn6400 - nn6410;
nn6410 - nn6420; nn6420 - nn6430; nn6430 - nn6440;
nn6440 - nn6450; nn6450 - nn6460; nn6460 - nn6470;
nn6470 - nn6480; nn6480 - nn6490; nn6490 - nn6500;
30 nn6500 - nn6510; nn6510 - nn6520; nn6520 - nn6530;
nn6530 - nn6540; nn6540 - nn6550; nn6550 - nn6560;
nn6560 - nn6570; nn6570 - nn6580; nn6580 - nn6590;
nn6590 - nn6600; nn6600 - nn6610; nn6610 - nn6620;
nn6620 - nn6630; nn6630 - nn6640; nn6640 - nn6650;
35 nn6650 - nn6660; nn6660 - nn6670; nn6670 - nn6680;
nn6680 - nn6690; nn6690 - nn6700; nn6700 - nn6710;

nn6710 - nn6720; nn6720 - nn6730; nn6730 - nn6740;
nn6740 - nn6750; nn6750 - nn6760; nn6760 - nn6770;
nn6770 - nn6780; nn6780 - nn6790; nn6790 - nn6800;
nn6800 - nn6810; nn6810 - nn6820; nn6820 - nn6830;
5 nn6830 - nn6840; nn6840 - nn6850; nn6850 - nn6860;
nn6860 - nn6870; nn6870 - nn6880; nn6880 - nn6890;
nn6890 - nn6900; nn6900 - nn6910; nn6910 - nn6920;
nn6920 - nn6930; nn6930 - nn6940; nn6940 - nn6950;
nn6950 - nn6960; nn6960 - nn6970; nn6970 - nn6980;
10 nn6980 - nn6990; nn6990 - nn7000; nn7000 - nn7010;
nn7010 - nn7020; nn7020 - nn7030; nn7030 - nn7040;
nn7040 - nn7050; nn7050 - nn7060; nn7060 - nn7070;
nn7070 - nn7080; nn7080 - nn7090; nn7090 - nn7100;
nn7100 - nn7110; nn7110 - nn7120; nn7120 - nn7130;
15 nn7130 - nn7140; nn7140 - nn7150; nn7150 - nn7160;
nn7160 - nn7170; nn7170 - nn7180; nn7180 - nn7190;
nn7190 - nn7200; nn7200 - nn7210; nn7210 - nn7220;
nn7220 - nn7230; nn7230 - nn7240; nn7240 - nn7250;
nn7250 - nn7260; nn7260 - nn7270; nn7270 - nn7280;
20 nn7280 - nn7290; nn7290 - nn7300; nn7300 - nn7310;
nn7310 - nn7320; nn7320 - nn7330; nn7330 - nn7340;
nn7340 - nn7350; nn7350 - nn7360; nn7360 - nn7370;
nn7370 - nn7380; nn7380 - nn7390; nn7390 - nn7400;
nn7400 - nn7410; nn7410 - nn7420; nn7420 - nn7430;
25 nn7430 - nn7440; nn7440 - nn7450; nn7450 - nn7460;
nn7460 - nn7470; nn7470 - nn7480; nn7480 - nn7490;
nn7490 - nn7500; nn7500 - nn7510; nn7510 - nn7520;
nn7520 - nn7530; nn7530 - nn7540; nn7540 - nn7550;
nn7550 - nn7560; nn7560 - nn7570; nn7570 - nn7580;
30 nn7580 - nn7590; nn7590 - nn7600; nn7600 - nn7610;
nn7610 - nn7620; nn7620 - nn7630; nn7630 - nn7640;
nn7640 - nn7650; nn7650 - nn7660; nn7660 - nn7670;
nn7670 - nn7680; nn7680 - nn7690; nn7690 - nn7700;
nn7700 - nn7710; nn7710 - nn7720; nn7720 - nn7730;
35 nn7730 - nn7740; nn7740 - nn7750; nn7750 - nn7760;
nn7760 - nn7770; nn7770 - nn7780; nn7780 - nn7790;

nn7790 - nn7800; nn7800 - nn7810; nn7810 - nn7820;
nn7820 - nn7830; nn7830 - nn7840; nn7840 - nn7850;
nn7850 - nn7860; nn7860 - nn7870; nn7870 - nn7880;
nn7880 - nn7890; nn7890 - nn7900; nn7900 - nn7910;
5 nn7910 - nn7920; nn7920 - nn7930; nn7930 - nn7940;
nn7940 - nn7950; nn7950 - nn7960; nn7960 - nn7970;
nn7970 - nn7980; nn7980 - nn7990; nn7990 - nn8000;
nn8000 - nn8010; nn8010 - nn8020; nn8020 - nn8030;
nn8030 - nn8040; nn8040 - nn8050; nn8050 - nn8060;
10 nn8060 - nn8070; nn8070 - nn8080; nn8080 - nn8090;
nn8090 - nn8100; nn8100 - nn8110; nn8110 - nn8120;
nn8120 - nn8130; nn8130 - nn8140; nn8140 - nn8150;
nn8150 - nn8160; nn8160 - nn8170; nn8170 - nn8180;
nn8180 - nn8190; nn8190 - nn8200; nn8200 - nn8210;
15 nn8210 - nn8220; nn8220 - nn8230; nn8230 - nn8240;
nn8240 - nn8250; nn8250 - nn8260; nn8260 - nn8270;
nn8270 - nn8280; nn8280 - nn8290; nn8290 - nn8300;
nn8300 - nn8310; nn8310 - nn8320; nn8320 - nn8330;
nn8330 - nn8340; nn8340 - nn8350; nn8350 - nn8360;
20 nn8360 - nn8370; nn8370 - nn8380; nn8380 - nn8390;
nn8390 - nn8400; nn8400 - nn8410; nn8410 - nn8420;
nn8420 - nn8430; nn8430 - nn8440; nn8440 - nn8450;
nn8450 - nn8460; nn8460 - nn8470; nn8470 - nn8480;
nn8480 - nn8490; nn8490 - nn8500; nn8500 - nn8510;
25 nn8510 - nn8520; nn8520 - nn8530; nn8530 - nn8540;
nn8540 - nn8550; nn8550 - nn8560; nn8560 - nn8570;
nn8570 - nn8580; nn8580 - nn8590; nn8590 - nn8600;
nn8600 - nn8610; nn8610 - nn8620; nn8620 - nn8630;
nn8630 - nn8640; nn8640 - nn8650; nn8650 - nn8660;
30 nn8660 - nn8670; nn8670 - nn8680; nn8680 - nn8690;
nn8690 - nn8700; nn8700 - nn8710; nn8710 - nn8720;
nn8720 - nn8730; nn8730 - nn8740; nn8740 - nn8750;
nn8750 - nn8760; nn8760 - nn8770; nn8770 - nn8780;
nn8780 - nn8790; nn8790 - nn8800; nn8800 - nn8810;
35 nn8810 - nn8820; nn8820 - nn8830; nn8830 - nn8840;
nn8840 - nn8850; nn8850 - nn8860; nn8860 - nn8870;

nn₈₈₇₀ - nn₈₈₈₀; nn₈₈₈₀ - nn₈₈₉₀; nn₈₈₉₀ - nn₈₉₀₀;
nn₈₉₀₀ - nn₈₉₁₀; nn₈₉₁₀ - nn₈₉₂₀; nn₈₉₂₀ - nn₈₉₃₀;
nn₈₉₃₀ - nn₈₉₄₀; nn₈₉₄₀ - nn₈₉₅₀; nn₈₉₅₀ - nn₈₉₆₀;
nn₈₉₆₀ - nn₈₉₇₀; nn₈₉₇₀ - nn₈₉₈₀; nn₈₉₈₀ - nn₈₉₉₀;
5 nn₈₉₉₀ - nn₉₀₀₀; nn₉₀₀₀ - nn₉₀₁₀; nn₉₀₁₀ - nn₉₀₂₀;
nn₉₀₂₀ - nn₉₀₃₀; nn₉₀₃₀ - nn₉₀₄₀; nn₉₀₄₀ - nn₉₀₅₀;
nn₉₀₅₀ - nn₉₀₆₀.

The oligomer, however, need not consist only of
10 the sequence which is complementary to the targeted HCV
sequence. It may contain in addition, nucleotide
sequences or other moieties which are suitable for the
purposes for which the oligomers are used. For example,
if the oligomers are used as primers for the amplification
15 of HCV sequences via PCR, they may contain sequences
which, when in duplex, form restriction enzyme sites which
facilitate the cloning of the amplified sequences. For
example, also, if the oligomers are to be used as "capture
probes" in hybridization assays (described infra), they
20 would contain in addition a binding partner which is
coupled to the oligomer containing the nucleotide sequence
which is complementary to the targeted HCV sequence.
Other types of moieties or sequences which are useful of
which the oligomers may be comprised or coupled to, are
25 those which are known in the art to be suitable for a
variety of purposes, including the labeling of nucleotide
probes.

The preparation of the oligomers is by means
known in the art, including, for example, by methods which
30 include excision, transcription, or chemical synthesis.
The target sequences and/or regions of the genome which
are selected to which the targeting polynucleotides of the
oligomers are complementary depend upon the purpose. For
example, if the goal is to screen for the presence of HCV
35 in biological samples (e.g. blood), the preferred
oligomers would be used as probes and/or primers, and

would hybridize to conserved regions of the HCV genome. Some of the conserved regions of the HCV genome to which the oligomers may bind are described herein, for example, the regions which include nucleotide numbers from about
5 the 5-terminus to about 200, or from about 4000 to about 5000, or from about 8000 to about 9040 as shown in Fig. 18, or preferably nucleotides -318 to 174, 4056 to 4448, and 4378 to 4902. Other regions of the genome which are conserved are readily ascertainable by comparison of the
10 nucleotide sequences of various isolates of HCV, including the prototype HCV, HCV1. Methods for conducting comparisons between genotypes to determine conserved and nonconserved regions are known in the art, and examples of these methods are disclosed herein.

15 In the basic nucleic acid hybridization assay, single-stranded analyte nucleic acid (either DNA or RNA) is hybridized to a nucleic acid probe, and resulting duplexes are detected. The probes for HCV polynucleotides (natural or derived) are a length which allows the detec-
20 tion of unique viral sequences by hybridization. While 6-8 nucleotides may be a workable length, sequences of 10-12 nucleotides are preferred, and about 20 nucleotides or more appears optimal. Preferably, these sequences will derive from regions which lack heterogeneity. These
25 probes can be prepared using routine methods, including automated oligonucleotide synthetic methods. Among useful probes, for example, are those derived from the newly isolated clones disclosed herein, as well as the various oligomers useful in probing cDNA libraries, set forth
30 below. A complement to any unique portion of the HCV genome will be satisfactory. For use as probes, complete complementarity is desirable, though it may be unnecessary as the length of the fragment is increased.

For use of such probes as agents to detect the
35 presence of HCV polynucleotides (for example in screening for contaminated blood), the biological sample to be

analyzed, such as blood or serum, may be treated, if desired, to extract the nucleic acids contained therein. The resulting nucleic acid from the sample may be subjected to gel electrophoresis or other size separation techniques; alternatively, the nucleic acid sample may be dot blotted without size separation. In order to form hybrid duplexes with the targeting sequence of the probe, the targeted region of the analyte nucleic acid must be in single stranded form. Where the sequence is naturally present in single stranded form, denaturation will not be required. However, where the sequence is present in double stranded form, the sequence will be denatured. Denaturation can be carried out by various techniques known in the art. Subsequent to denaturation, the analyte nucleic acid and probe are incubated under conditions which promote stable hybrid formation of the target sequence in the probe with the putative targeted sequence in the analyte, and the resulting duplexes containing the probe(s) are detected.

Detection of the resulting duplex, if any, is usually accomplished by the use of labeled probes; alternatively, the probe may be unlabeled, but may be detectable by specific binding with a ligand which is labeled, either directly or indirectly. Suitable labels, and methods for labeling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation or kinasing), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies, and the like.

The region of the probes which are used to bind to the analyte can be made completely complementary to the HCV genome. Therefore, usually high stringency conditions are desirable in order to prevent false positives. However, conditions of high stringency should only be used if the probes are complementary to regions of the viral

genome which lack heterogeneity. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, length of time, and

- 5 concentration of formamide. These factors are outlined in, for example, Maniatis, T. (1982).

- Variations of this basic scheme which are known in the art, including those which facilitate separation of the duplexes to be detected from extraneous materials and/
- 10 or which amplify the signal from the labeled moiety, may also be used. A number of these variations are reviewed in, for example: Matthews and Kricka (1988), Analytical Biochemistry 169:1; Landegren et al. (1988), Science 242:229; and Mittlin (1989), Clinical chem. 35:1819.
- 15 These and the following publications describing assay formats are hereby incorporated by reference herein. Probes suitable for detecting HCV in these assays are comprised of sequences which hybridize with target HCV polynucleotide sequences to form duplexes with the analyte
- 20 strand, wherein the duplexes are of sufficient stability for detection in the specified assay system.

- A suitable variation is, for example, one which is described in U.S. Patent No. 4,868,105, issued Sept. 9, 1989, and in E.P.O. Publication No. 225,807 (published
- 25 June 16, 1987). These publications describe a solution phase nucleic acid hybridization assay in which the analyte nucleic acid is hybridized to a labeling probe set and to a capturing probe set. The probe-analyte complex is coupled by hybridization with a solid-supported capture
- 30 probe that is complementary to the capture probe set. This permits the analyte nucleic acid to be removed from solution as a solid phase complex. Having the analyte in the form of a solid phase complex facilitates subsequent separation steps in the assay. The labeling probe set is
- 35 complementary to a labeled probe that is bound through hybridization to the solid phase/analyte complex.

Generally, it is expected that the HCV genome sequences will be present in serum of infected individuals at relatively low levels, i.e., at approximately 10^2 - 10^3 chimp infectious doses (CID) per ml. This level may

5 require that amplification techniques be used in hybridization assays. Such techniques are known in the art. For example, the Enzo Biochemical Corporation "Bio-Bridge" system uses terminal deoxynucleotide transferase to add unmodified 3'-poly-dT-tails to a DNA probe. The

10 poly dT-tailed probe is hybridized to the target nucleotide sequence, and then to a biotin-modified poly-A. PCT Publication 84/03520 and EP Publication No. 124221 describe a DNA hybridization assay in which: (1) analyte is annealed to a single-stranded DNA probe that is com-

15plementary to an enzyme-labeled oligonucleotide; and (2) the resulting tailed duplex is hybridized to an enzyme-labeled oligonucleotide. EPA 204510 describes a DNA hybridization assay in which analyte DNA is contacted with a probe that has a tail, such as a poly-dT tail, an ampli-

20fier strand that has a sequence that hybridizes to the tail of the probe, such as a poly-A sequence, and which is capable of binding a plurality of labeled strands. A type of hybridization assay which is described in E.P.O. Publication No. 317,077 (published May 24, 1989), which

25should detect sequences at the level of approximately 10^6 /ml, utilizes nucleic acid multimers which bind to single-stranded analyte nucleic acid, and which also bind to a multiplicity of single-stranded labeled oligonucleotides. A particularly desirable technique may involve amplifica-

30tion of the target HCV sequences in sera approximately 10,000 fold (i.e., to approximately 10^6 sequences/ml), as part of the hybridization system. The amplification may be accomplished, for example, by the polymerase chain reactions (PCR) technique described by Saiki et al. (1986),

35by Mullis, U.S. Patent No. 4,683,195, and by Mullis et al. U.S. Patent No. 4,683,202. Amplification may be prior to,

or preferably subsequent to purification of the HCV target sequence. For example, amplification may be utilized in conjunction with the assay methods described in U.S. Patent No. 4,868,105, or if even further amplification is
5 desired, in conjunction with the hybridization system described in E.P.O. Publication No. 317,077.

Preferred methods for detecting HCV sequences in an analyte polynucleotide strand are based upon the hybridization detection methods described in U.S. Patent
10 No. 4,868,105 and in E.P.O. Publication No. 317,077.

These methods are solution-phase sandwich hybridization assays which utilize both capture and label probes which hybridize to target sequences in an analyte nucleic acid. In the use of these assays to screen biological samples
15 for HCV, the probes used would bind to conserved regions of the HCV genome. The capture and label probes may be interspersed in their binding to the target sequence. Alternatively, in a preferred mode the capture and label probes are in sets, and the probes of one set do not
20 intersperse with the probes of another set. In the latter mode, preferably the set(s) of multiple capture probes hybridize to the most conserved regions of the genome, while the set(s) of multiple label probes may hybridize to regions which exhibit small amounts of divergence. For
25 example, using the prototype HCV1 cDNA sequence shown in Fig. 18, probes could be used which hybridize to sequences in the region of nucleotides from about -318 to about 174, and/or nucleotides in the region of about 4378 to about 4902, and/or nucleotides in the region of from about 4056
30 to about 4448. The preferred probes would hybridize to sequences in the 5'-region of the HCV genome, since, as shown infra., this region appears to be highly conserved. Thus, preferred probes may hybridize to, for example, nucleotides from about -318 to about 174 as shown in Fig.
35 18. Probes could be used which hybridize to either the positive strand in conserved regions, and/or its comple-

ment, depending upon the purpose, for example, to detect viral genomic sequences, or to detect HCV cDNA sequences resulting from PCR amplification, or to detect replicative intermediates to the positive HCV RNA strand.

5

Detection of HCV RNA and Polynucleotides Derived Therefrom
Using an HCV/cPCR Method

A particularly useful method for detecting HCV RNA or polynucleotides derived from HCV RNA is the HCV/
10 cPCR method, which is a subject of the herein application, and which utilizes the polymerase chain reaction technique (PCR) which is described by Saiki et al. (1986), by Mullis in U.S. Pat. No. 4,683,195, and by Mullis et al. in U.S. Patent No. 4,683,202. The HCV/cPCR method utilizes prim-
15 ers and probes derived from the information provided herein concerning the nature of the HCV genome.

Generally, in the PCR technique, short oligonucleotide primers are prepared which match opposite ends of a desired sequence. The sequence between the
20 primers need not be known. A sample of polynucleotide is extracted and denatured, preferably by heat, and hybridized with oligonucleotide primers which are present in molar excess. Polymerization is catalyzed by a template- and primer-dependent polymerase in the presence of
25 deoxynucleotide triphosphates or nucleotide analogs (dNTPs). This results in two "long products" which contain the respective primers at their 5'-termini, covalently linked to the newly synthesized complements of the original strands. The replicated DNA is again de-
30 natured, hybridized with oligonucleotide primers, returned to polymerizing conditions, and a second cycle of replication is initiated. The second cycle provides the two original strands, the two long products from cycle 1, and two "short products" replicated from the long products.
35 The short products contain sequences (sense or antisense) derived from the target sequence, flanked at the 5'- and

3'-termini with primer sequences. On each additional cycle, the number of short products is replicated exponentially. Thus, this process causes the amplification of a specific target sequence.

- 5 In the method, a sample is provided which is suspected of containing HCV RNA, or a fragment thereof. The sample is usually taken from an individual suspected of having NANBH; however, other sources of the sample are included, e.g., conditioned medium or cells from in vitro
- 10 systems in which the virus has been replicated. The sample, however, must contain the target nucleic acid sequence(s).

- The sample is then subjected to conditions which allow reverse transcription of HCV RNA into HCV cDNA.
- 15 Conditions for reverse transcribing RNA are known to those of skill in the art, and are described in, for example, Maniatis et al. (1982), and in Methods in Enzymology. A preferred method of reverse transcription utilizes reverse transcriptase from a variety of sources, including re-
- 20 combinant molecules, and isolated from, for example, a retrovirus, preferably from avian myeloblastosis virus (AMV), and suitable conditions for the transcription. The HCV cDNA product of reverse transcription is in a RNA:DNA hybrid, which results from the first round of reverse
- 25 transcription; subsequently, DNA:DNA hybrids result from two or more rounds of transcription.

- The HCV cDNA resulting from reverse transcription is then subjected to PCR to amplify the target sequence. In order to accomplish this, the HCV cDNA is
- 30 denatured, and the separated strands are hybridized with primers which flank the target sequence.

- Strand separation may be accomplished by any suitable denaturing method, including physical, chemical, or enzymatic means, which are known to those of skill in
- 35 the art. A preferred method, which is physical, involves heating the nucleic acid until it is completely (>99%)

denatured. Typical heat denaturation involves temperatures ranging from about 80°C to about 105°C, for times ranging from about 1 to 10 minutes.

After hybridization of the HCV cDNA with the
5 primers, the target HCV sequences are replicated by a polymerizing means which utilizes a primer oligonucleotide to initiate the synthesis of the replicate chain. The primers are selected so that they are complementary to sequences of the HCV genome. Oligomeric primers which are
10 complementary to regions of the sense and antisense strands of HCV cDNA can be designed from the HCV cDNA sequences from the composite cDNA sequence provided in Fig. 18.

The primers are selected so that their relative
15 positions along a duplex sequence are such that an extension product synthesized from one primer, when it is separated from its template (complement), serves as a template for the extension of the other primer to yield a replicate chain of defined length.

20 The primer is preferably single stranded for maximum efficiency in amplification, but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is an
25 oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact lengths of the primers will depend on many factors, including temperature and source of the primer and use of
30 the method. For example, depending on the complexity of the target sequence, the oligonucleotide primer typically contains about 15-45 nucleotides, although it may contain more or fewer nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently
35 stable hybrid complexes with the template.

The primers used herein are selected to be "substantially" complementary to the different strands of each specific sequence to be amplified. Therefore, the primers need not reflect the exact sequence of the template, but must be sufficiently complementary to selectively hybridize with their respective strands. For example, a non-complementary nucleotide fragment may be attached to the 5'-end of the primer, with the remainder of the primer sequence being complementary to the strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the primer, provided that the primer has sufficient complementarity with the sequence of one of the strands to be amplified to hybridize therewith, and to thereby form a duplex structure which can be extended by the polymerizing means. The non-complementary nucleotide sequences of the primers may include restriction enzyme sites. Appending a restriction enzyme site to the end(s) of the target sequence would be particularly helpful for cloning of the target sequence.

It will be understood that "primer", as used herein, may refer to more than one primer, particularly in the case where there is some ambiguity in the information regarding the terminal sequence(s) of the target region to be amplified. Hence, a "primer" includes a collection of primer oligonucleotides containing sequences representing the possible variations in the sequence or includes nucleotides which allow a typical basepairing. One of the primer oligonucleotides in this collection will be homologous with the end of the target sequence. A specific case is shown in the Examples, where oligomer sets of 44-mers and 45-mers were utilized to prime the amplification of a potentially variant region of the HCV genome.

It is anticipated that there will be a variety of strains or isolates of HCV with sequences which deviate from HCV1, the prototype strain. Therefore, in order to

detect variant strains it is preferable to construct primers which hybridize to conserved regions of the HCV genome. The conserved regions may be determined by comparing the nucleotide or amino acid sequences of several HCV strains/isolates. There appear to be at least three regions of conserved amino acid in the HCV genome, described supra., from which primers may be derived. These regions are believed to be. The primers described infra., in the Examples, are derived from what are believed to be conserved regions of HCV, based upon sequence homology to that of the Flaviviruses.

The oligonucleotide primers may be prepared by any suitable method. Methods for preparing oligonucleotides of specific sequence are known in the art, and include, for example, cloning and restriction of appropriate sequences, and direct chemical synthesis. Chemical synthesis methods may include, for example, the phosphotriester method described by Narang et al. (1979), the phosphodiester method disclosed by Brown et al. (1979), the diethylphosphoramidate method disclosed in Beaucage et al. (1981), and the solid support method in U.S. Patent No. 4,458,066.

The primers may be labeled, if desired, by incorporating means detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means.

Template-dependent extension of the oligonucleotide primer(s) is catalyzed by a polymerizing agent in the presence of adequate amounts of the four deoxyribonucleotide triphosphates (dATP, dGTP, dCTP and dTTP) or analogs, in a reaction medium which is comprised of the appropriate salts, metal cations, and pH buffering system. Suitable polymerizing agents are enzymes known to catalyze primer- and template-dependent DNA synthesis. Known DNA polymerases include, for example, E. coli DNA polymerase I or its Klenow fragment, T₄ DNA polymerase, and Taq DNA polymerase. The reaction conditions for

catalyzing DNA synthesis with these DNA polymerases are known in the art.

The products of the synthesis are duplex molecules consisting of the template strands and the primer extension strands, which include the target sequence. These products, in turn, serve as template for another round of replication. In the second round of replication, the primer extension strand of the first cycle is annealed with its complementary primer; synthesis yields a "short" product which is bounded on both the 5'- and the 3'-ends by primer sequences or their complements. Repeated cycles of denaturation, primer annealing, and extension result in the exponential accumulation of the target region defined by the primers. Sufficient cycles are run to achieve the desired amount of polynucleotide containing the target region of nucleic acid. The desired amount may vary, and is determined by the function which the product polynucleotide is to serve.

The PCR method can be performed in a number of temporal sequences. For example, it can be performed step-wise, where after each step new reagents are added, or in a fashion where all of the reagents are added simultaneously, or in a partial step-wise fashion, where fresh reagents are added after a given number of steps.

In a preferred method, the PCR reaction is carried out as an automated process which utilizes a thermostable enzyme. In this process the reaction mixture is cycled through a denaturing region, a primer annealing region, and a reaction region. A machine may be employed which is specifically adapted for use with a thermostable enzyme, which utilizes temperature cycling without a liquid handling system, since the enzyme need not be added at every cycle. This type of machine is commercially available from Perkin Elmer Cetus Corp.

After amplification by PCR, the target polynucleotides are detected by hybridization with a probe

polynucleotide which forms a stable hybrid with that of the target sequence under stringent to moderately stringent hybridization and wash conditions. If it is expected that the probes will be completely complementary (i.e., about 99% or greater) to the target sequence, stringent conditions will be used. If some mismatching is expected, for example if variant strains are expected with the result that the probe will not be completely complementary, the stringency of hybridization may be lessened. However, conditions are chosen which rule out nonspecific/adventitious binding. Conditions which affect hybridization, and which select against nonspecific binding are known in the art, and are described in, for example, Maniatis et al. (1982). Generally, lower salt concentration and higher temperature increase the stringency of binding. For example, it is usually considered that stringent conditions are incubation in solutions which contain approximately 0.1 X SSC, 0.1% SDS, at about 65°C incubation/wash temperature, and moderately stringent conditions are incubation in solutions which contain approximately 1-2 X SSC, 0.1% SDS and about 50°-65°C incubation/wash temperature. Low stringency conditions are 2 X SSC and about 30°-50°C.

Probes for HCV target sequences may be derived from the HCV cDNA sequence shown in Fig. 18, or from new HCV isolates. The HCV probes may be of any suitable length which span the target region, but which exclude the primers, and which allow specific hybridization to the target region. If there is to be complete complementarity, i.e., if the strain contains a sequence identical to that of the probe, since the duplex will be relatively stable under even stringent conditions, the probes may be short, i.e., in the range of about 10-30 base pairs. If some degree of mismatch is expected with the probe, i.e., if it is suspected that the probe will hybridize to a variant region, the probe may be of greater

length, since length seems to counterbalance some of the effect of the mismatch(es). An example of this is found in the Examples, where the probe was designed to bind to potential variants of HCV1. In this case, the primers
5 were designed to bind to HCV cDNA derived from a hypothetical conserved region of the HCV genome, and the target region was one which potentially contained variations (based upon the Flavivirus model). The probe used to detect the HCV target sequences contained approximately
10 268 base pairs.

The probe nucleic acid having a sequence complementary to the target sequence may be synthesized using similar techniques described supra. for the synthesis of primer sequences. If desired, the probe may be labeled.
15 Appropriate labels are described supra.

In some cases, it may be desirable to determine the length of the PCR product detected by the probe. This may be particularly true if it is suspected that variant HCV strains may contain deletions within the target
20 region, or if one wishes to confirm the length of the PCR product. In such cases it is preferable to subject the products to size analysis as well as hybridization with the probe. Methods for determining the size of nucleic acids are known in the art, and include, for example, gel
25 electrophoresis, sedimentation in gradients, and gel exclusion chromatography.

The presence of the target sequence in a biological sample is detected by determining whether a hybrid has been formed between the HCV polynucleotide probe and
30 the nucleic acid subjected to the PCR amplification technique. Methods to detect hybrids formed between a probe and a nucleic acid sequence are known in the art. For example, for convenience, an unlabeled sample may be transferred to a solid matrix to which it binds, and the
35 bound sample subjected to conditions which allow specific hybridization with a labeled probe; the solid matrix is

than examined for the presence of the labeled probe. Alternatively, if the sample is labeled, the unlabeled probe is bound to the matrix, and after the exposure to the appropriate hybridization conditions, the matrix is
5 examined for the presence of label. Other suitable hybridization assays are described supra.

Determination of Variant HCV Sequences Using PCR

In order to identify variant HCV strains, and
10 thereby to design probes for those variants, the above described HCV/cPCR method is utilized to amplify variant regions of the HCV genome, so that the nucleotide sequences of these variant target regions can be determined. Generally, variant types of HCV might be
15 expected to occur in different geographic locations than that in which the HCV1 strain is predominant, for example, Japan, Africa, etc.; or in different vertebrate species which are also infected with the virus. Variant HCV may also arise during passage in tissue culture systems, or be
20 the result of spontaneous or induced mutations.

In order to amplify the variant target region, primers are designed to flank the suspect region, and preferably are complementary to conserved regions. Primers to two regions of HCV which are probably conserved,
25 based upon the Flavivirus model, are described in the Examples. These primers and probes may be designed utilizing the sequence information for the HCV1 strain provided in Fig. 18.

Analysis of the nucleotide sequence of the
30 target region(s) may be by direct analysis of the PCR amplified products. A process for direct sequence analysis of PCR amplified products is described in Saiki et al. (1988).

Alternatively, the amplified target sequence(s)
35 may be cloned prior to sequence analysis. A method for the direct cloning and sequence analysis of enzymatically

amplified genomic segments has been described by Scharf (1986). In the method, the primers used in the PCR technique are modified near their 5'-ends to produce convenient restriction sites for cloning directly into, for example, an M13 sequencing vector. After amplification, the PCR products are cleaved with the appropriate restriction enzymes. The restriction fragments are ligated into the M13 vector, and transformed into, for example, a JM 103 host, plated out, and the resulting plaques are screened by hybridization with a labeled oligonucleotide probe. Other methods for cloning and sequence analysis are known in the art.

Universal Primers for Flaviviruses and for HCV

Studies of the nature of the genome of the HCV, utilizing probes derived from the HCV cDNA, as well as sequence information contained within the HCV cDNA, are suggestive that HCV is a Flavi-like virus. These studies are described in E.P.O. publication No. 318,216 owned by the herein assignee, and which is incorporated herein in its entirety. A comparison of the HCV cDNA sequence derived from the HCV cDNA clones with known sequences of a number of Flaviviruses show that HCV contains sequences which are homologous to conserved sequences in the Flaviviruses. These conserved sequences may allow the creation of primers which may be universal in their application for amplification of target regions of Flaviviruses, and for HCV. These sequences are the 16-mer or smaller sequences from the 3'-termini of the primers described in the Examples. Identification of the species is then accomplished utilizing a probe specific for the species. The genomes of a number of Flaviviruses are known in the art, and include, for example, Japanese Encephalitis Virus (Sumiyoshi et al. (1987)), Yellow Fever Virus (Rice et al. (1985)), Dengue Type 2 Virus (Hahn et al. (1988)), Dengue Type 4 Virus (Mackow (1987)), and West

Nile Virus (Castle et al. (1986)). Identification of HCV RNA is accomplished utilizing a probe specific for HCV, the sequence of which can be determined the HCV cDNA sequences provided herein.

- 5 Alternatively, utilization of sets of probe(s) designed to account for codon degeneracy and therefore contain common sequences to the Flaviviruses and to HCV, as determined by a comparison of HCV amino acid sequences with the known sequences of the Flaviviruses, allows a
10 general detection system for these viruses.

Construction of Desired DNA Sequences

- Synthetic oligonucleotides may be prepared using an automated oligonucleotide synthesizer as described by
15 Warner (1984). If desired the synthetic strands may be labeled with ^{32}P by treatment with polynucleotide kinase in the presence of ^{32}P -ATP, using standard conditions for the reaction.

- DNA sequences, including those isolated from
20 cDNA libraries, may be modified by known techniques, including, for example site directed mutagenesis, as described by Zoller (1982). Briefly, the DNA to be modified is packaged into phage as a single stranded sequence, and converted to a double stranded DNA with DNA
25 polymerase using, as a primer, a synthetic oligonucleotide complementary to the portion of the DNA to be modified, and having the desired modification included in its own sequence. The resulting double stranded DNA is transformed into a phage supporting host bacterium.
30 Cultures of the transformed bacteria, which contain replications of each strand of the phage, are plated in agar to obtain plaques. Theoretically, 50% of the new plaques contain phage having the mutated sequence, and the remaining 50% have the original sequence. Replicates of
35 the plaques are hybridized to labeled synthetic probe at temperatures and conditions which permit hybridization

with the correct strand, but not with the unmodified sequence. The sequences which have been identified by hybridization are recovered and cloned.

5 Kits for Screening for HCV Derived Polynucleotides

Oligomers which are probes and/or primers for amplification and/or screening of samples for HCV can be packaged into kits. Kits for screening for HCV sequences include the oligomeric probe DNAs. Kits for amplification of HCV sequences may include the oligomeric primers used in the amplification. The kits usually contain the probes or primers in a premeasured or predetermined amount, as well as other suitably packaged reagents and materials, in separate suitable containers, needed for the particular hybridization and/or amplification protocol(s). For example, the kit may contain standards, buffers, supports, enzymes, substrates, label probes, binding partners, and/or instructions for conducting the test.

20

Examples

Described below are examples of the present invention which are provided only for illustrative purposes, and not to limit the scope of the present invention.

25

Isolation and Sequence of Overlapping

HCV cDNA Clones 13i, 26j, CA59a, CA84a, CA156e and CA167b

The clones 13i, 26j, CA59a, CA84a, CA156e and CA167b were isolated from the lambda-gt11 library which contains HCV cDNA (ATCC No. 40394), the preparation of which is described in E.P.O. Publication No. 318,216 (published 31 May 1989), and WO 89/04669 (published 1 June 1989). Screening of the library was with the probes described infra., using the method described in Huynh (1985). The frequencies with which positive clones appeared with the respective probes was about 1 in 50,000.

The isolation of clone 13i was accomplished using a synthetic probe derived from the sequence of clone 12f. The sequence of the probe was:

5' GAA CGT TGC GAT CTG GAA GAC AGG GAC AGG 3'.

The isolation of clone 26j was accomplished using a probe derived from the 5'-region of clone K9-1. The sequence of the probe was:

10

5' TAT CAG TTA TGC CAA CGG AAG CGG CCC CGA 3'.

The isolation procedures for clone 12f and for clone k9-1 (also called K9-1) are described in E.P.O. Publication No. 318,216, and their sequences are shown in Figs. 1 and 2, respectively. The HCV cDNA sequences of clones 13i and 26j, are shown in Figs. 4 and 5, respectively. Also shown are the amino acids encoded therein, as well as the overlap of clone 13i with clone 12f, and the overlap of clone 26j with clone 13i. The sequences for these clones confirmed the sequence of clone K9-1. Clone K9-1 had been isolated from a different HCV cDNA library (See E.P.O. Publication No. 218,316).

Clone CA59a was isolated utilizing a probe based upon the sequence of the 5'-region of clone 26j. The sequence of this probe was:

5' CTG GTT AGC AGG GCT TTT CTA TCA CCA CAA 3'.

30 A probe derived from the sequence of clone CA59a was used to isolate clone CA84a. The sequence of the probe used for this isolation was:

5' AAG GTC CTG GTA GTG CTG CTG CTA TTT GCC 3'.

35

Clone CA156e was isolated using a probe derived from the sequence of clone CA84a. The sequence of the probe was:

5 5' ACT GGA CGA CGC AAG GTT GCA ATT GCT CTA 3'.

Clone CA167b was isolated using a probe derived from the sequence of clone CA 156e. The sequence of the probe was:

10

5' TTC GAC GTC ACA TCG ATC TGC TTG TCG GGA 3'.

The nucleotide sequences of the HCV cDNAs in clones CA59a, CA84a, CA156e, and CA167b, are shown Figs. 15 6, 7, 8, and 9, respectively. The amino acids encoded therein, as well as the overlap with the sequences of relevant clones, are also shown in the figures.

Creation of "pi" HCV cDNA Library

20 A library of HCV cDNA, the "pi" library, was constructed from the same batch of infectious chimpanzee plasma used to construct the lambda-gt11 HCV cDNA library (ATCC No. 40394) described in E.P.O. Publication No. 318,216, and utilizing essentially the same techniques. 25 However, construction of the pi library utilized a primer-extension method, in which the primer for reverse transcriptase was based on the sequence of clone CA59a. The sequence of the primer was:

30 5' GGT GAC GTG GGT TTC 3'.

Isolation and Sequence of Clone pil4a

Screening of the "pi" HCV cDNA library described supra., with the probe used to isolate clone CA167b (See 35 supra.) yielded clone pil4a. The clone contains about 800 base pairs of cDNA which overlaps clones CA167b, CA156e,

CA84a and CA59a, which were isolated from the lambda gt-11 HCV cDNA library (ATCC No. 40394). In addition, pil4a also contains about 250 base pairs of DNA which are upstream of the HCV cDNA in clone CA167b.

5

Isolation and Sequence of Clones CA216a, CA290a and aq30a

Based on the sequence of clone CA167b a synthetic probe was made having the following sequence:

10

5' GGC TTT ACC ACG TCA CCA ATG ATT GCC CTA 3'

The above probe was used to screen the , which yielded clone CA216a, whose HCV sequences are shown in Fig. 10.

Another probe was made based on the sequence of
15 clone CA216a having the following sequence:

5' TTT GGG TAA GGT CAT CGA TAC CCT TAC GTG 3'

Screening the lambda-gt11 library (ATCC No. 40394) with
20 this probe yielded clone CA290a, the HCV sequences therein being shown in Fig. 11.

In a parallel approach, a primer-extension cDNA library was made using nucleic acid extracted from the same infectious plasma used in the original lambda-gt11
25 cDNA library described above. The primer used was based on the sequence of clones CA216a and CA290a:

5' GAA GCC GCA CGT AAG 3'

30 The cDNA library was made using methods similar to those described previously for libraries used in the isolation of clones pil4a and k9-1. The probe used to screen this library was based on the sequence of clone CA290a:

35

5' CCG GCG TAG GTC GCG CAA TTT GGG TAA 3'

Clone ag30a was isolated from the new library with the above probe, and contained about 670 basepairs of HCV sequence. See Fig. 12. Part of this sequence overlaps the HCV sequence of clones CA216a and CA290a. About 300
5 base-pairs of the ag30a sequence, however, is upstream of the sequence from clone CA290a. The non-overlapping sequence shows a start codon (*) and stop codons that may indicate the start of the HCV ORF. Also indicated in Fig. 12 are putative small encoded peptides (#) which may play
10 a role in regulating translation, as well as the putative first amino acid of the putative polypeptide (/), and downstream amino acids encoded therein.

Isolation and Sequence of Clone CA205a

15 Clone CA205a was isolated from the original lambda gt-11 library (ATCC No. 40394), using a synthetic probe derived from the HCV sequence in clone CA290a (Fig. 11). The sequence of the probe was:

20 5' TCA GAT CGT TGG TGG AGT TTA CTT GTT GCC 3'.

The sequence of the HCV cDNA in CA205a, shown in Fig. 13, overlaps with the cDNA sequences in both clones ag30a and CA290a. The overlap of the sequence with that of CA290a
25 is shown by the dotted line above the sequence (the figure also shows the putative amino acids encoded in this fragment).

As observed from the HCV cDNA sequences in clones CA205a and ag30a, the putative HCV polyprotein appears to begin at the ATG start codon; the HCV sequences
30 in both clones contain an in-frame, contiguous double stop codon (TGATAG) forty two nucleotides upstream from this ATG. The HCV ORF appears to begin after these stop codons, and to extend for at least 8907 nucleotides (See
35 the composite HCV cDNA shown in Fig. 18).

Isolation and Sequence of Clone 18g

Based on the sequence of clone ag30a (See Fig. 12) and of an overlapping clone from the original lambda gt-11 library (ATCC No. 40394), CA230a, a synthetic probe was made having the following sequence:

5' CCA TAG TGG TCT GCG GAA CCG GTG AGT ACA 3'.

- 10 Screening of the original lambda-gt11 HCV cDNA library with the probe yielded clone 18g, the HCV cDNA sequence of which is shown in Fig. 14. Also shown in the figure are the overlap with clone ag30a, and putative polypeptides encoded within the HCV cDNA.
- 15 The cDNA in clone 18g (C18g or 18g) overlaps that in clones ag30a and CA205a, described supra. The sequence of C18g also contains the double stop codon region observed in clone ag30a. The polynucleotide region upstream of these stop codons presumably represents part
- 20 of the 5'-region of the HCV genome, which may contain short ORFs, and which can be confirmed by direct sequencing of the purified HCV genome. These putative small encoded peptides may play a regulatory role in translation. The region of the HCV genome upstream of that
- 25 represented by C18g can be isolated for sequence analysis using essentially the technique described in E.P.O. Publication No. 318,216 for isolating cDNA sequences upstream of the HCV cDNA sequence in clone 12f. Essentially, small synthetic oligonucleotide primers of
- 30 reverse transcriptase, which are based upon the sequence of C18g, are synthesized and used to bind to the corresponding sequence in HCV genomic RNA. The primer sequences are proximal to the known 5'-terminal of C18g, but sufficiently downstream to allow the design of probe
- 35 sequences upstream of the primer sequences. Known standard methods of priming and cloning are used. The

resulting cDNA libraries are screened with sequences upstream of the priming sites (as deduced from the elucidated sequence of C18g). The HCV genomic RNA is obtained from either plasma or liver samples from

5 individuals with NANBH. Since HCV appears to be a Flavi-like virus, the 5'-terminus of the genome may be modified with a "cap" structure. It is known that Flavivirus genomes contain 5'-terminal "cap" structures. (Yellow

10 Fever virus, Rice et al. (1988); Dengue virus, Hahn et al (1988); Japanese Encephalitis Virus (1987)).

Isolation and Sequence of Clones from
the beta-HCV cDNA library

Clones containing cDNA representative of the 3'-

15 terminal region of the HCV genome were isolated from a cDNA library constructed from the original infectious chimpanzee plasma pool which was used for the creation of the HCV cDNA lambda-gt11 library (ATCC No. 40394), described in E.P.O. Publication No. 318,216. In order to

20 create the DNA library, RNA extracted from the plasma was "tailed" with poly rA using poly (rA) polymerase, and cDNA was synthesized using oligo(dT)₁₂₋₁₈ as a primer for reverse transcriptase. The resulting RNA:cDNA hybrid was digested with RNAase H, and converted to double stranded

25 HCV cDNA. The resulting HCV cDNA was cloned into lambda-gt10, using essentially the technique described in Huynh (1985), yielding the beta (or b) HCV cDNA library. The procedures used were as follows.

An aliquot (12ml) of the plasma was treated with

30 proteinase K, and extracted with an equal volume of phenol saturated with 0.05M Tris-Cl, pH 7.5, 0.05% (v/v) beta-mercaptoethanol, 0.1% (w/v) hydroxyquinolone, 1 mM EDTA. The resulting aqueous phase was re-extracted with the phenol mixture, followed by 3 extractions with a 1:1

35 mixture containing phenol and chloroform:isoamyl alcohol (24:1), followed by 2 extractions with a mixture of

chloroform and isoamyl alcohol (1:1). Subsequent to adjustment of the aqueous phase to 200 mM with respect to NaCl, nucleic acids in the aqueous phase were precipitated overnight at -20°C , with 2.5 volumes of cold absolute ethanol. The precipitates were collected by centrifugation at 10,000 RPM for 40 min., washed with 70% ethanol containing 20 mM NaCl, and with 100% cold ethanol, dried for 5 min. in a dessicator, and dissolved in water.

The isolated nucleic acids from the infectious chimpanzee plasma pool were tailed with poly rA utilizing poly-A polymerase in the presence of human placenta ribonuclease inhibitor (HPRI) (purchased from Amersham Corp.), utilizing MS2 RNA as carrier. Isolated nucleic acids equivalent to that in 2 ml of plasma were incubated in a solution containing TMN (50 mM Tris HCl, pH 7.9, 10 mM MgCl_2 , 250 mM NaCl, 2.5 mM MnCl_2 , 2 mM dithiothreitol (DTT)), 40 micromolar alpha- ^{32}P ATP, 20 units HPRI (Amersham Corp.), and about 9 to 10 units of RNase free poly-A polymerase (BRL). Incubation was for 10 min. at 37°C , and the reactions were stopped with EDTA (final concentration about 250 mM). The solution was extracted with an equal volume of phenol-chloroform, and with an equal volume of chloroform, and nucleic acids were precipitated overnight at -20°C with 2.5 volumes of ethanol in the presence of 200 mM NaCl.

Isolation of Clone b5a

The beta HCV cDNA library was screened by hybridization using a synthetic probe, which had a sequence based upon the HCV cDNA sequence in clone 15e. The isolation of clone 15e is described in E.P.O. Publication No. 318,216, and its sequence is shown in Fig. 3. The sequence of the synthetic probe was:

5' ATT GCG AGA TCT ACG GGG CCT GCT ACT CCA 3'.

Screening of the library yielded clone beta-5a (b5a), which contains an HCV cDNA region of approximately 1000 base pairs. The 5'-region of this cDNA overlaps clones 35f, 19g, 26g, and 15e (these clones are described supra).

5 The region between the 3'-terminal poly-A sequence and the 3'-sequence which overlaps clone 15e, contains approximately 200 base pairs. This clone allows the identification of a region of the 3'-terminal sequence the HCV genome.

10 The sequence of b5a is contained within the sequence of the HCV cDNA in clone 16jh (described infra). Moreover, the sequence is also present in CC34a, isolated from the original lambda-gt11 library (ATCC No. 40394). (The original lambda-gt11 library is referred to herein as
15 the "C" library).

Isolation and Sequence of Clones Generated by PCR
Amplification of the 3'-Region of the HCV Genome

Multiple cDNA clones have been generated which
20 contain nucleotide sequences derived from the 3'-region of the HCV genome. This was accomplished by amplifying a targeted region of the genome by a polymerase chain reaction technique described in Saiki et al. (1986), and in Saiki et al. (1988), which was modified as described
25 below. The HCV RNA which was amplified was obtained from the original infectious chimpanzee plasma pool which was used for the creation of the HCV cDNA lambda-gt11 library (ATCC No. 40394) described in E.P.O. Publication No. 318,216. Isolation of the HCV RNA was as described supra.
30 The isolated RNA was tailed at the 3'-end with ATP by E. coli poly-A polymerase as described in Sippel (1973), except that the nucleic acids isolated from chimp serum were substituted for the nucleic acid substrate. The tailed RNA was then reverse transcribed into cDNA by
35 reverse transcriptase, using an oligo dT-primer adapter,

essentially as described by Han (1987), except that the components and sequence of the primer-adapter were:

	<u>Stuffer</u>	<u>NotI</u>	<u>SP6 Promoter</u>	<u>Primer</u>
5	AATTC	GCGGCCGC	CATACGATTTAGGTGACACTATAGAA	T ₁₅

The resultant cDNA was subjected to amplification by PCR using two primers:

10	<u>Primer</u>	<u>Sequence</u>
	JH32 (30mer)	ATAGCGGCCGCCCTCGATTGCGAGATCTAC
	JH11 (20mer)	AATTCGGGCGGCCGCCATACGA

The JH32 primer contained 20 nucleotide sequences
 15 hybridizable to the 5'-end of the target region in the cDNA, with an estimated T_m of 66°C. The JH11 was derived from a portion of the oligo dT-primer adapter; thus, it is specific to the 3'-end of the cDNA with a T_m of 64°C. Both primers were designed to have a recognition site for
 20 the restriction enzyme, NotI, at the 5'-end, for use in subsequent cloning of the amplified HCV cDNA.

The PCR reaction was carried out by suspending the cDNA and the primers in 100 microliters of reaction mixture containing the four deoxynucleoside triphosphates,
 25 buffer salts and metal ions, and a thermostable DNA polymerase isolated from Thermus aquaticus (Taq polymerase), which are in a Perkin Elmer Cetus PCR kit (N801-0043 or N801-0055). The PCR reaction was performed for 35 cycles in a Perkin Elmer Cetus DNA thermal cycler.
 30 Each cycle consisted of a 1.5 min denaturation step at 94°C, an annealing step at 60°C for 2 min, and a primer extension step at 72°C for 3 min. The PCR products were subjected to Southern blot analysis using a 30 nucleotide probe, JH34, the sequence of which was based upon that of
 35 the 3'-terminal region of clone 15e. The sequence of JH34 is:

5' CTT GAT CTA CCT CCA ATC ATT CAA AGA CTC 3'.

The PCR products detected by the HCV cDNA probe ranged in size from about 50 to about 400 base pairs.

In order to clone the amplified HCV cDNA, the PCR products were cleaved with NotI and size selected by polyacrylamide gel electrophoresis. DNA larger than 300 base pairs was cloned into the NotI site of pUC18S. The vector pUC18S is constructed by including a NotI polylinker cloned between the EcoRI and Sall sites of pUC18. The clones were screened for HCV cDNA using the JH34 probe. A number of positive clones were obtained and sequenced. The nucleotide sequence of the HCV cDNA insert in one of these clones, 16jh, and the amino acids encoded therein, are shown in Fig. 15. A nucleotide heterogeneity, detected in the sequence of the HCV cDNA in clone 16jh as compared to another clone of this region, is indicated in the figure.

Isolation and Sequence of Clone 6k

Based on the sequence of clone 16jh and clone b5a (see supra), a synthetic probe was made having the following sequence:

5' TCT TCA ACT GGG CAG TAA GAA CAA AGC TCA 3'.

Screening of the original lambda-gt11 HCV cDNA library (described in E.P.O. Publication No. 318,216) with the probe yielded clones with a frequency of approximately 1 in 10^6 ; one of these was called clone 6k (also called C6k), the HCV cDNA sequence of which is shown in Fig. 16. Also shown in the figure are the overlap with clone 16jh, and putative polypeptides encoded within the HCV cDNA. Sequence information on the HCV cDNA in clone 6k was obtained from only one strand. Information on the deposit

of this clone is provided infra, wherein the clone is listed as Lambda gt11 C6k. Confirmation of the C6K sequence as part of an ORF encoding HCV1 polypeptide has been obtained by sequencing other overlapping clones.

5

Isolation and Sequence of Clone p131jh

A clone containing sequence from the 3'-region of the HCV genome, and which contains an in-frame stop codon, was isolated essentially as described supra., for the isolation of clones generated by PCR amplification of the 3' region of the genome, except that HCV1 RNA was converted to cDNA using the oligonucleotide

15 5' AAT TCG CGG CCG CCA TAC GAT TTA GGT GAC
 ACT ATA GAA T₁₅ 3'.

The cDNA was then amplified by the PCR reaction using the primers:

20 5' TTC GCG GCC GCT ACA GCG GGG GAG ACA T 3'

and

25 5' AAT TCG CGG CCG CCA TAC GA 3'.

After amplification, the PCR products were precipitated with spermine, digested with NotI, and extracted with phenol. The purified products were cloned into the NotI site of pUC18S, and HCV positive clones were selected using the oligonucleotide:

5' CGA TGA AGG TTG GGG TAA ACA CTC CGG CCT 3'.

35 The HCV cDNA in one clone, designated p131jh, is shown in Fig. 17. This clone contains an in-frame stop codon for the large ORF contained in the HCV genome.

Isolation and Sequence of Clone 5'-clone32

A clone containing sequence from the 5'-region of the HCV genome, upstream of the sequence in clone b114a, was isolated and the nucleotide sequence determined by a modification of the method for the isolation and sequence of clones generated by PCR amplification of the 3'-region of the genome, described in U.S.S.N. 456,637, which is incorporated by reference. Generally, a target region of the genome was amplified by the PCR technique described in Saiki et al. (1986), and in Saiki et al (1988). The HCV RNA which was amplified was obtained by extracting human serum (U.S. clinical isolate, HCV27) using a cold guanidinium thiocyanate method described by Han et al. (1987). The extracted RNA was converted into single stranded cDNA with reverse transcriptase, using a primer, JH94, which is complementary to nucleotides -250 to -223 of the HCV genome (see Fig. 18). The sequence of JH94 is:

20

5' CCT GCG GCC GCA CGA CAC TCA TAC TAA 3'.

Conversion of single- to double-stranded HCV cDNA was accomplished by tailing the DNA with approximately 20 to 50 dA residues using terminal deoxynucleotidyl transferase (Sambrook et al. (1989), MOLECULAR CLONING), and replicating the tailed molecule using the following oligo-dT primer-adaptor, which contains a NotI site, and an sp6 promoter:

30

<u>Stuffer</u>	<u>NotI</u>	<u>SP6 Promoter</u>	<u>Primer</u>
AATTC	GCGGCCGC	CATACGATTTAGGTGACACTATAGAA	T ₁₅

The resultant cDNA was subjected to amplification by PCR using two primers, JH94 (described supra.) and JH11, which has the following sequence.

35

<u>Primer</u>	<u>Sequence</u>
JH11 (20mer)	AATTCGGGCGGCCCATACGA


5 The PCR reaction was carried out by suspending
the cDNA and the primers in 100 microliters of reaction
mixture containing the four deoxynucleoside triphosphates,
buffer salts and metal ions, and a thermostable DNA
polymerase isolated from Thermus aquaticus (Taq
10 polymerase), which are in a Perkin Elmer Cetus PCR kit
(N801-0043 or N801-0055). The PCR reaction was performed
for 35 cycles in a Perkin Elmer Cetus DNA thermal cycler.
Each cycle consisted of a 1.5 min denaturation step at
94°C, an annealing step at 60°C for 2 min, and a primer
15 extension step at 72°C for 3 min.

The PCR products were digested with NotI, and
cloned into pUC18S. Clones containing HCV nucleotide
sequences were obtained by screening with a probe, Alex90,
which is derived from nucleotides -312 to -283 of the HCV1
20 genome, and which has the sequence:

5' ACC ATG AAT CAC TCC CCT GTG AGG AAC TAC 3'.

The HCV cDNAs in the isolated clones were sequenced by the
25 dideoxy chain termination method (Sanger et al. (1977)).
The sequence of HCV cDNA in one of the isolated clones,
5'-clone32, spans the region of nucleotides -224 to -341
in Fig. 18.

An analysis of the nucleotide sequence of the
30 HCV cDNA showed that the replicate of the HCV RNA strand
contains a GC-rich stretch which may be capable of forming
a stable hairpin structure:



In the structure, the dashed lines indicate possible hydrogen bonds between complementary nucleotides.

10 A search in the computer database, Genbank, revealed that homologous sequences were absent from known viral sequences. Thus, this sequence may be unique to the 5'-terminus of the HCV genome.

A hairpin structure may serve as a recognition
15 signal for a transcriptase and/or it may contribute to the
stability of the RNA at the 5'-terminus.

Compiled HCV cDNA Sequences

An HCV cDNA sequence has been compiled from a series of overlapping clones derived from various HCV cDNA libraries described herein, and in E.P.O. Publication No. 318,216. The clones from which Fig. 18 has been derived are clone 5'-32, b114a, 18g, ag30a, CA205a, CA290a, CA216a, p114a, CA167b, CA156e, CA84a, CA59a, K9-1 (also called k9-1), 26j, 131, 12f, 14i, 11b, 7f, 7e, 8h, 33c, 40b, 37b, 35, 36, 81, 32, 33b, 25c, 14c, 8f, 33f, 33g, 39c, 35f, 19g, 26g, 15e, b5a, 16jh, C6k and p131jh. The methods for isolation of these clones, as well as their sequences, are discussed herein, and in E.P.O. Publication No. 318,216, which is incorporated herein by reference. In Fig. 18, the three dashes above the sequence indicate the position of the putative initiator methionine codon.

Clone b114a overlaps with clones 18g, ag30a, and CA205a, except that clone b114a contains an extra two
35 nucleotides upstream of the sequence in clone 18g (i.e.,

5'-CA). These extra two nucleotides have been included in the HCV genomic sequence shown in Fig. 18.

It should be noted that although several of the clones described supra. have been obtained from libraries other than the original HCV cDNA lambda-gt11 C library (ATCC No. 40394), these clones contain HCV cDNA sequences which overlap HCV cDNA sequences in the original library. Thus, essentially all of the HCV sequence is derivable from the original lambda-gt11 C library (ATCC No. 40394) which was used to isolate the first HCV cDNA clone (5-1-1). The isolation of clone 5-1-1 is described in E.P.O. Publication No. 318,216, which is incorporated herein by reference.

The putative sequence of the major HCV polyprotein encoded in the composite of HCV1 cDNA is also shown. The first amino acid in the sequence is the putative initiator methionine of the large ORF. The variant amino acids, due to the clonal heterogeneities, are indicated above the sequence. Since the lambda gt11 library was created from serum obtained from one individual (see E.P.O. Publication No. 318,216), the results suggest that variant viral sequences (both nucleotide and amino acid) are present in that individual.

An examination of the composite HCV cDNA sequence shows that besides the large ORF, there are a number of ORFs upstream of that encoding the polyprotein, and within the sequence encoding the polyprotein there are a large number of smaller ORFs in the other two translational frames. The ORFs upstream of the HCV polyprotein are shown in the Table immediately below.

Table
ORFs Upstream of that Encoding the Large
HCV Polyprotein

5			
	<u>Nucl. #</u>	<u>Translation Frame</u>	<u>Amino Acid Sequence</u>
	-310	1	MNHSPVRNYCLHAESV
	-329	3	MGATLHHESLPCEELL
			SSRRKRLAMALV
10	-246	2	MSVVQPPGPPLPGEP
	-127	1	MPGDLGVPPQDC

The reading frame, position, and size of the ORFs downstream of the sequence encoding the putative initiator MET of the polyprotein are shown in the Table below. The major polyprotein is that translated from reading frame 2.

Table
ORFs Downstream of the Putative Initiator MET
Encoding Sequence

20		<u>Reading Frame</u>	<u>Size(aa)</u>	<u>Position(bp)</u>
		1	168	696
		1	105	2343
25		1	119	5616
		2	3025	-42
		3	160	5
		3	111	1667
		3	148	6893

30 In addition to the above, an examination of the sequence which is complementary to the genomic strand of HCV RNA also contains several small ORFs. One of these ORFs, which is complementary to nucleotides -341 to +837 in the HCV RNA sequence, encodes a polypeptide of 385 amino acids.

Comparison of the Sequences of 5'-Regions
Obtained from HCV Isolates from Different
Geographical Locations

5 Nucleotide sequences from the 5'- regions of HCV isolates from the U.S.A. (HCV18, HCV27), from Italy (HCV11, HCV124), and from Korea (HCVK1) were compared.

Isolation of the HCV cDNA sequences was essentially as described supra., for the isolation of 5'-
 10 clone32, except for the following. The extracted RNA was reverse-transcribed into cDNA using as primers either JH51 or r16, which are complementary to HCV nucleotides -90 to -73 and 366 to 383, respectively. The sequences of these primers are as follows.

15

<u>Primer</u>	<u>Sequence</u>
JH51	5' CCC AAC ACT ACT CGG CTA 3'
r16	5' CAC GTA AGG GTA TCG ATG 3'

20 Amplification of the HCV dsDNA was by the PCR method using JH93 and JH52 as 5'- and 3'- primers, respectively. The HCV sequence in JH93 is derived from HCV nucleotides -317 to -296, that in JH52 is from HCV nucleotides -93 to -117; the nucleotide numbers are indicated in parentheses below
 25 the sequences. In JH52 the underlined dinucleotide has been mutated to create the NotI site. The sequences of these primers are the following.

(Primer)	<u>Stuffer</u>	<u>NotI</u>	<u>HCV sequence</u>
30 (JH93)	5' TTC	GCGGCCGC	ACTCCATGAATCACTCCCC 3'
		(-317)	(-296)
(JH52)	5' AGTCTT	GCGG <u>CCGC</u>	ACGCCCAAATC 3'
	(-93)		(-117)

35

After amplification, the PCR products were cleaved by NotI, and cloned into pUC18S. The HCV cDNAs were sequenced either by direct sequencing after amplification by PCR, or alternatively, the cloned HCV cDNAs were
5 sequenced by the primer extension and the dideoxy method. Primer extension and the dideoxy method of sequencing were performed as described supra., for the sequence of 5'-clone32.

The PCR method for direct sequencing used Alex90
10 (see supra. for the sequence) as the 5'-primer, and r25 as the 3'-primer. Alex90 is derived from HCV nucleotides -312 to -283, and r25 is derived from nucleotides 365 to 342 (See Fig. 18). The sequence of r25 is:

15 5' ACC TTA CCC AAA TTG CGC GAC CTA 3'.

A comparison of the sequences of the 5'-region of HCV27, HCVK1, HCVI1, HCVI24, and HCV18 with the sequence of the prototype HCV, HCV1, showed the following.
20 The examined 5'- region is highly conserved amongst the 5 HCV isolates. The sequences appeared to be identical except for one nucleotide which was deleted at position -171 in HCVI24, and for the ambiguity in four nucleotides at positions -222 to -219 in isolate HCVK1.

25 The high levels of sequence conservation in this region may reflect the role of this region in viral replication, and/or transcription, and/or translation.

30 Sequence Variations in HCV Isolates
from Different Individuals

Isolates of HCV which contain sequences which deviate from CDC/HCV1 were identified in human individuals, some of whom were serologically positive for anti-C100-3 antibodies (EC10 was antibody negative).
35 Identification of these new isolates was accomplished by cloning and sequencing segments of the HCV genome which

had been amplified by the PCR technique using CDC/HCl sequences. Amplification was accomplished essentially based on an HCV/cPCR method. The method utilizes primers and probes based upon the HCV cDNA sequences described
5 herein. The first step in the method is the synthesis of a cDNA to either the HCV genome, or its replicative intermediate, using reverse transcriptase. After synthesis of the HCV cDNA, and prior to amplification, the RNA in the sample is degraded by techniques known in the art. A
10 designated segment of the HCV cDNA is then amplified by the use of the appropriate primers. The amplified sequences are cloned, and clones containing the amplified sequences are detected by a probe which is complementary to a sequence lying between the primers, but which does
15 not overlap the primers.

HCV Isolates Isolated from Humans in the U.S.

Blood samples which were used as a source of HCV virions were obtained from the American Red Cross in
20 Charlotte, North Carolina, and from the Community Blood Center of Kansas, Kansas City, Missouri. The samples were screened for antibodies to the HCV C100-3 antigen using an ELISA assay as described in E.P.O. Publication No. 318,216, and subjected to supplemental Western blot
25 analysis using a polyclonal goat anti-human HRP to measure anti-HCV antibodies. Two samples, #23 and #27, from the American Red Cross and from the Community Blood Center of Kansas, respectively, were determined to be HCV positive by these assays.
30 Viral particles present in the serum of these samples were isolated by ultracentrifugation under the conditions described by Bradley et al. (1985). RNA was extracted from the particles by digestion with proteinase K and SDS at final concentrations of 10 micrograms/ml
35 proteinase K, and 0.1% SDS; digestion was for 1 hour at 37°C. Viral RNA was further purified by extraction with

chloroform-phenol, as described in E.P.O. Publication No. 318,216.

HCV RNA in the preparation of RNA was reverse transcribed into cDNA essentially as described in E.P.O. Publication No. 318,216, except that the oligonucleotide JHC 7, which corresponds to the cDNA sequence 1958-1939, and which has the following sequence, was used as primer for the reverse transcriptase reaction.

10 JHC 7: CCA GCG GTG GCC TGG TAT TG.

After both strands of the cDNA were synthesized, the resulting cDNA was then amplified by the PCR method essentially as described supra. for the isolation of clones generated by PCR amplification, except that the oligonucleotide primers used, i.e., JHC 6 and ALX 80, were designed to amplify a 1080 nucleotide segment of the HCV genome from CDC/HCV1 nucleotides 673 to 1751. The primers, in addition, are designed to incorporate a NOT I restriction site at the 3'-end of the PCR product, and a blunt end at the 5'-terminus. The sequences of the primers is:

25 ALX 80: TTT GGG TAA GGT CAT CGA TAC CCT TAC GTG;

and

JHC 6: ATA TGC GGC CGC CTT CCG TTG GCA TAA.

30 ALX 80 corresponds to nucleotides 673-702 of the CDC/HCV1 sequence; JHC 6 corresponds to nucleotides 1752-1738 of the HCV1 (in addition there are 12 extra nucleotides which encode a NotI site). The designation of nucleotides in JHC 6, i.e., a declining number, indicates the placement
35 in the anti-sense strand.

After PCR amplification with the above described primers, the blunt end terminus was converted into a NOT I site as follows. A homopolymer tail of 15 dGs was attached to the PCR product using terminal deoxynucleotide transferase, and the products were again subjected to amplification by PCR using as primers JHC 6 and JHC 13. The latter primer, JHC 13, the sequence of which follows, is designed to contain a NOT I site in addition to an SP6 phage promoter. (The SP6 promoter is described in GENETIC ENGINEERING, J. Setlow Ed. (1988)).

JHC 13: AAT TCG CGG CCG CCA TAC GAT TTA GGT GAC
ACT ATA GAA CCC CCC CCC CCC CCC.

In order to clone the amplified HCV cDNA, the PCR products were cleaved with NotI, precipitated with spermine to remove free oligonucleotides (Hoopes et al. (1981)), and cloned into the NotI site of pUC18S (see Section IV.A.34.). The HCV cDNAs in three clones derived from each HCV isolate, were subjected to sequence analysis. Analysis was essentially by the method described in Chen and Seeburg (1985).

Consensus sequences of the clones derived from HCV in samples 23 and 27 are shown in Fig. 46 and Fig. 47, respectively. The variable sequences are also shown in these figures, as are the amino acids encoded in the consensus sequences.

Fig. 39 and Fig. 40 show comparisons of the aligned positive strand nucleotide sequences (Fig. 39) and putative amino acid sequences (Fig. 40) of samples 23, 27, and HCV1. The amino acid sequence of HCV1 in Fig. 39 represents amino acid numbers 129-467 of the HCV polyprotein encoded by the large ORF in the HCV genomic RNA. An examination of Fig. 46 and Fig. 47 show that there are variations in the sequences of the three isolated clones. The sequence variations at the

nucleotide level and the amino acid level are summarized in the table immediately below. In the table, the polypeptides designated S and NS1 represent amino acid numbers 130 to ~380, and 380 to ~470, respectively. The numbering is from the putative initiator methionine. The terminology S and NS1 is based upon the positioning of the sequences encoding the polypeptides using the Flavivirus model. As discussed above, however, recent evidence suggests that there is not total correlation between HCV and the Flaviviruses with regard to viral polypeptide domains, particularly in the putative E/NS1 domains. Indeed, HCV polypeptides and their coding domains may exhibit substantial deviation from the Flavivirus model.

15

Table
Sequence Homology

	Nucleotide Encoding			Amino Acid Encoded		
	overall	S	NS1	overall	S	NS1
	%	%	%	%	%	%
HCV1/HCV23	93	95	91	92	95	87
HCV1/HCV27	89	93	84	89	95	82
HCV23/HCV27	89	93	85	90	93	84

25

Although there are variations in the newly isolated HCV sequences, the cloned sequences from samples 23 and 27 (called HCV23 and HCV27) each contain 1019 nucleotides, indicating a lack of deletion and addition mutants in this region in the selected clones. The sequences in Figs. 39 and 40 also show that the isolated sequences are not rearranged in this region.

30

A comparison of the consensus sequences for HCV1 and for the other isolates of HCV is summarized in the Table, supra. The sequence variations between the chimpanzee isolate HCV1, and the HCVs isolated from humans

35

are about the same as that seen between the HCVs of human origin.

It is of interest that the sequence variations in two of the putative domains is not uniform. The
5 sequence in a putative S region appears to be relatively constant, and randomly scattered throughout the region. In contrast, a putative NS1 region has a higher degree of variability than the overall sequence, and the variation appears to be in a hypervariable pocket of about 28 amino
10 acids which is located about 70 amino acids downstream from the putative N-terminus of the putative polyprotein.

Although it may be argued that the detected variations were introduced during the amplification process, it is unlikely that all of the variations are from
15 this result. It has been estimated that Taq polymerase introduces errors into a sequence at approximately one base per 10 kilobases of DNA template per cycle (Saiki et al. (1988)). Based upon this estimate, up to 7 errors may have been introduced during the PCR amplification of the
20 1019 bp DNA fragment. However, the three subclones of HCV-23 and HCV-27 yielded 29 and 14 base variations, respectively. The following suggest that these variations are naturally occurring. About 60% of the base changes are silent mutations which do not change the amino acid
25 sequence. Variations introduced by the Taq polymerase during PCR amplification would be expected to occur randomly; however, the results show that the variant sequences are clustered in at least one specific region. Moreover, a consensus sequence was derived by sequencing
30 multiple different clones derived from the PCR amplified products.

HCV Isolates from Humans in
Italy and in the U.S.

35 Segments of HCV RNA present in different isolates were amplified by the HCV/cPCR method. These

segments span a region of ~0.6Kb to ~1.6Kb downstream from the methionine encoding start codon of the putative HCV polyprotein. The isolates are from biological specimens obtained from HCV infected individuals. More

- 5 specifically, isolate HCT #18 is from human plasma from an individual in the U.S.A., EC1 and EC10 are from a liver biopsy of an Italian patient, and Th is from a peripheral blood mononucleocyte fraction of an American patient. Comparable segments of HCV RNA have been isolated from a
10 chimpanzee.

- RNA was extracted from the human plasma specimens using phenol:CHCl₃:isoamyl alcohol extraction. Either 0.1 ml or 0.01 ml of plasma was diluted to a final volume of 1.0 ml, with a TENB/proteinase K/SDS solution
15 (0.05 M Tris-HCL, pH 8.0, 0.001 M EDTA, 0.1 M NaCl, 1 mg/ml Proteinase K, and 0.5% SDS) containing 10 to 40 micrograms/ml polyadenylic acid, and incubated at 37°C for 60 minutes. After this proteinase K digestion, the resultant plasma fractions were deproteinized by extrac-
20 tion with TE (50 mM Tris-HCl, pH 8.0, 1 mM EDTA) saturated phenol, pH 6.5. The phenol phase was separated by centrifugation, and was reextracted with TENB containing 0.1% SDS. The resulting aqueous phases from each extrac-
25 tion were pooled, and extracted twice with an equal volume of phenol/chloroform/isoamyl alcohol [1:1(99:1)], and then twice with an equal volume of a 99:1 mixture of chloroform/isoamyl alcohol. Following phase separation by centrifugation, the aqueous phase was brought to a final concentration of 0.2 M Na Acetate, and the nucleic acids
30 were precipitated by the addition of two volumes of ethanol. The precipitated nucleic acids were recovered by ultracentrifugation in a SW 41 rotor at 38 K, for 60 minutes at 4°C, or in a microfuge for 10 minutes at 10K, 4°C.

RNA extracted from the liver biopsy was provided by Dr. F. Bonino, Ospedale Maggiore di S. Giovanni Battista, Torino, Italy.

The mononucleocyte fraction was obtained by sedimentation of the individual's aliquot of blood through Ficoll-Paque® (Pharmacia Corp), using the manufacturer's directions. Total RNA was extracted from the fraction using the guanidinium thiocyanate procedure described in E.P.O. Publication No. 318,216 (See also Choo et al (1989)).

Synthesis of HCV cDNA from the samples was accomplished using reverse transcriptase, and primers derived from clone 156e and from clone K91. These primers, which are anti-sense relative to the genomic RNA, have the following sequences.

156e16B: 5' CGA CAA GAA AGA CAG A 3',
 and
K91/16B 5' CGT TGG CAT AAC TGA T 3'.

20 Following ethanol precipitation, the precipitated RNA or nucleic acid fraction was dried, and resuspended in DEPC treated distilled water. Secondary structures in the nucleic acids were disrupted by heating
25 at 65°C for 10 minutes, and the samples were immediately cooled on ice. cDNA was synthesized using 1 to 3 micrograms of total RNA from liver, or from nucleic acids (or RNA) extracted from 10 to 100 microliters of plasma. The synthesis utilized reverse transcriptase, and was in a 25
30 microliter reaction, using the protocol specified by the manufacturer, BRL. All reaction mixtures for cDNA synthesis contained 23 units of the RNAase inhibitor, RNASIN™ (Fisher/Promega). Following cDNA synthesis, the reaction mixtures were diluted with water, boiled for 10
35 minutes, and quickly chilled on ice.

Each set of samples was subjected to two rounds of PCR amplification. The primers for the reactions were selected to amplify regions designated "EnvL" and EnvR". The "EnvL" region encompasses nucleotides 669-1243, and putative amino acids 117 to 308; the "EnvR" region encompasses nucleotides 1215-1629, and encodes putative amino acids 300-408 (the putative amino acids are numbered starting from the putative methionine initiation codon). The relationship of these regions relative to the putative polyprotein encoded in the HCV cDNA, and to the polypeptides encoded in the Flavivirus model is shown in Fig. 48.

The primers for the first round of PCR reactions were derived from the HCV cDNA sequences in either clone ag30a, clone 156e, or clone k9-1. The primers used for the amplification of the EnvL region were 156e16B (shown supra), and ag30a16A for the sense strand; the amplification of the EnvR region utilized the primer K91/16B (shown supra), and 156e16a for the sense strand. The sequences of the sense strand primers are the following.

For EnvL, ag30a16A: 5' CTC TAT GGC AAT GAG G 3',

and

For EnvR, 156e16A: 5' AGC TTC GAC GTC ACA T 3' .

The PCR reactions were performed essentially according to the manufacturer's directions (Cetus-Perkin-Elmer), except for the addition of 1 microgram of RNase A. The reactions were carried out in a final volume of 100 microliters. The PCR was performed for 30 cycles, utilizing a regimen of 94°C (1 min), 37°C (2 min), and 72°C (3 min), with a 7 minute extension at 72°C for the last cycle. The samples were then extracted with phenol:CHCl₃, ethanol precipitated two times, resuspended in 10 mM Tris

HCl, pH 8.0, and concentrated using Centricon-30 (Amicon) filtration. This procedure efficiently removes oligonucleotides less than 30 nucleotides in size; thus, the primers from the first round of PCR amplification are removed.

The Centricon-30 concentrated samples were then subjected to a second round of PCR amplification using probes designed from clones 202a and 156e for the EnvL region, and from 156e and 59a for the EnvR region. The primers for amplification of the EnvL region have the following sequences.

202aEnv41a: 5' CTT GAA TTC GCA ATT TGG GTA
AGG TCA TCG ATA CCC TTA CG 3'

15

and

156e38B': 5' CTT GAA TTC GAT AGA GCA ATT
GCA ACC TTG CGT CGT CC 3'.

20

The primers for amplification of the EnvR region in RNAs derived from humans have the following sequences.

156e38A': 5' CTT GAA TTC GGA CGA CGC AAG
GTT GCA ATT GCT CTA TC 3'

25

and

59aEnv39C: 5' CTT GAA TTC CAG CCG GTG TTG
AGG CTA TCA TTG CAG TTC 3'.

30

Amplification by PCR was for 35 cycles utilizing a regimen of 94°C (1 min), 60°C (1 min), and 72°C (2 min), with a 7 minute extension at 72°C for the last cycle. The samples were then extracted with phenol:CHCl₃, precipitated two times, and digested with EcoRI. The PCR reaction products

35

were analyzed by separation of the products by electrophoresis on 6% polyacrylamide gels. DNA of approximately the estimated size of the expected PCR product was electroeluted from the gels, and subcloned into either
5 a pGEM-4 plasmid vector or into lambda gt11. The expected product sizes for the EnvL and EnvR after the first round of amplification are 615 bp and 683 bp, respectively; after the second round of amplification the expected product sizes for EnvL and EnvR are 414 bp and 575 bp,
10 respectively. The plasmids containing the amplified products were used to transform host cells; the pGEM-4 plasmid was used to transform DH5-alpha, and lambda gt11 was used to transform C600 delta-HFL. Clones of the transformed cells which either hybridized to the appropri-
15 ate HCV probes (described below), or those which had inserts of the correct size were selected. The inserts were then cloned in M13 and sequenced.

The probes for all of the HCV/cPCR products consisted of ^{32}P labeled sections of HCV cDNA which had
20 been prepared by PCR amplification of a region of clone 216 (using CA216a16A and 216a16B as primers), and of clone 84 (using CA84a16A and CA84a16B or CA84a16C as primers); ^{32}P was introduced into the PCR products by nick translation. The probes for the first and second round of EnvL
25 amplification were from clone 216. Those for the first round of EnvR amplification were from 84 (i.e., CA84a16A and CA84a16B), for the second round of EnvL amplification were CA84a16A and CA84a16C. These probes did not overlap the primers used in the HCV/cPCR reactions. The sequence
30 of the primers for the PCR amplification of the probes is in the following table.

Table

	Primer	Clone	Sequence
5	CA216a16A	216	5' TGA ACT ATG CAA CAG G 3'
	CA216a16B	216	5' GGA GTG TGC AGG ATG G 3'
	CA84a16A	84	5' AAG GTT GCA ATT GCT C 3'
	CA84a16B	84	5' ACT AAC AGG ACC TTC G 3'
	CA84a16C	84	5' TAA CGG GTC ACC GCA T 3'

10

Sequence information on variants in the EnvL region was obtained from 3 clones from HCT #18, 2 clones from TH, 3 clones from EC1, and from the HCV1 clones described in E.P.O. Publication No. 318,216, and supra. A comparison of the composite nucleotide sequence of each isolate derived from these clones is shown in Fig. 49. In the figure, each sequence is shown 5' to 3' for the sense strand for the EnvL region, and the sequences have been aligned. The vertical lines and capital letters indicate sequence homology, the absence of a line and an uncapitalized letter indicates a lack of homology. The sequences shown in the lines are as follows: line 1, Thorn; line 2, EC1; line 3, HCT #18; line 4, HCV1.

Sequence information on variants in the EnvR region was obtained from two clones of EC10, and from the HCV1 clones described in E.P.O. Publication No. 318,216 and supra.. The two EC10 clones differed by only one nucleotide. A comparison of the nucleotide sequences of EC10(clone 2) and a composite of the HCV1 sequences is shown in Fig. 50; each sequence is shown 5' to 3' for the sense strand of the EnvR region, and the sequences have been aligned. The double dots between the sequences indicate sequence homology.

A comparison of the amino acid sequences encoded in the EnvL (amino acids #117-308) and EnvR region (amino acids #300-438) for each of the isolates is shown in Fig.

51 and Fig. 52, respectively. Included in the Figures are sequences for the isolates JH23 and JH27, described supra. Also indicated are sequences from a Japanese isolate; these sequences were provided by Dr. T. Miyamura, Japan.

- 5 In the figures, the amino acid sequence for the region is given in its entirety for HCV1, and the non-homologous amino acids in the various isolates are indicated.

As seen in Fig. 51, In the EnvL region there is overall about a 93% homology between HCV1 and the other
 10 isolates. HCT18, Th, and EC1 have about a 97% homology with HCV1; JH23 and JH27 have about 96% and about 95% homology, respectively, with HCV1. Fig. 52 shows that the homologies in the EnvR region are significantly less than in the EnvL region; moreover, one subregion appears to be
 15 hypervariable (i.e., from amino acid 383-405). This data is summarized in the Table immediately below.

Table
Homology of EnvR Region

20

Isolate	Percent Homology with HCV1	
	AA330-AA438	AA383-AA405
JH23(U.S.)	83	57
JH27(U.S.)	80	39
25 Japanese	73	48
EC10 (Italy)	84	48

Detection of Positive and Negative Strand
5'-HCV RNA in Serum

30

The RNA in HCV27, isolated from serum, was analyzed for the presence of positive and negative strands using the PCR method. The PCR method was performed essentially as described above, except for the following. The extracted HCV27 RNA was reverse transcribed into
 35 single-stranded cDNA using as a primer either Alex90 or JH52 (see supra. for the sequences). The sequence of

Alex90 matches that in nucleotides -312 to -283 of the positive strand of HCV RNA, whereas JH52 matches that of nucleotides -117 to -93 of the negative strand. the resulting single-stranded HCV cDNAs were each separately amplified by PCR using Alex90 and JH52. Detection of the amplified products was accomplished by Southern blotting, using Alex89 as the probe. Alex89 matches nucleotide numbers -203 to -175 of HCV RNA. The sequence of Alex89 is:

10

5' CCA TAG TGG TCT GCG GAA CCG GTG AGT ACA 3'.

The analysis indicated that, by this method, the signals of the amplified products of both RNA strands were of equal intensity. These results are suggestive that HCV RNA in the 5'-region may exist as double-stranded RNA.

Probes for Sandwich Hybridization for HCV

This example exemplifies the sets of label and capture probes useful to detect HCV RNA in biological samples, using essentially the assay described in U.S. Patent No. 4,868,105. The method is a solution-phase sandwich hybridization assay which utilizes both capture and label probes which hybridize to target sequences in an analyte nucleic acid. In the screening of biological samples for HCV, the probes used bind to conserved regions of the HCV genome, and the HCV binding regions are selected for their uniqueness to the HCV genome. The regions which bind to the binding partner of the capture probe, or the portion of the label probe which binds to the labeling moiety (or to an amplifying multimer if the method described in E.P.O. Publication No. 317,077 is used), are selected such that they do not bind to any of the known sequences in the databank or in HCV, and which have the appropriate content of Gs and Cs to allow stable duplex formation with their complements under the selec-

tion conditions. The capture and label probes are in sets, and the probes of one set do not intersperse with the probes of another set. These probes are comprised of sequences which are complementary to the following 5 nucleotide sequences in the coding strand of the prototype HCV cDNA sequence shown in Fig. 18.

Set 1

10	<u>Probe type</u>	<u>Probe Number</u>	<u>Complement of</u> <u>Nucleotide Numbers</u>
	Capture	42.XT1.1	-318 to -289
	Capture	42.XT1.2	-285 to -256
	Capture	42.XT1.3	-252 to -223
15	Capture	42.XT1.4	-219 to -190
	Label	42.LLA2C.5	-186 to -157
	Label	42.LLA2C.6	-153 to -124
	Label	42.LLA2C.7	-120 to -91
	Label	42.LLA2C.8	-87 to -58
20	Label	42.LLA2C.9	-54 to -25
	Label	42.LLA2C.10	-21 to 9
	Label	42.LLA2C.11	13 to 42
	Label	42.LLA2C.12	46 to 75
	Label	42.LLA2C.13	79 to 108
25	Label	42.LLA2C.14	112 to 141
	Label	42.LLA2C.15	145 to 174

30

35

Set 2

	<u>Probe type</u>	<u>Probe Number</u>	<u>Complement of Nucleotide Numbers</u>
5	Capture	42.16.XT1	4378 to 4407
	Capture	42.17.XT1	4411 to 4440
	Capture	42.18.XT1	4444 to 4473
	Capture	42.19.XT1	4477 to 4506
10	Capture	42.20.XT1	4510 to 4539
	Label	42.21.LLA2C	4543 to 4572
	Label	42.22.LLA2C	4576 to 4605
	Label	42.23.LLA2C	4609 to 4638
	Label	42.24.LLA2C	4642 to 4671
15	Label	42.25.LLA2C	4675 to 4704
	Label	42.26.LLA2C	4708 to 4737
	Label	42.27.LLA2C	4771 to 4770
	Label	42.28.LLA2C	4774 to 4803
	Label	42.29.LLA2C	4807 to 4836
20	Label	42.30.LLA2C	4840 to 4869
	Label	42.31.LLA2C	4873 to 4902

25

30

35

Set 3

5	<u>Probe type</u>	<u>Probe Number</u>	<u>Complement of</u> <u>Nucleotide Numbers</u>
	Capture	42.32.XT1	4056 to 4085
	Capture	42.33.XT1	4089 to 4085
	Capture	42.34.XT1	4122 to 4151
10	Capture	42.35.XT1	4155 to 4184
	Label	42.36.LLA2C	4188 to 4217
	Label	42.37.LLA2C	4221 to 4250
	Label	42.38.LLA2C	4254 to 4283
	Label	42.39.LLA2C	4287 to 4316
15	Label	42.40.LLA2C	4230 to 4349
	Label	42.41.LLA2C	4353 to 4382
	Label	42.42.LLA2C	4386 to 4415
	Label	42.43.LLA2C	4419 to 4448

- 20 In the above sets, each capture probe contains, in addition to the sequences complementary to the HCV sequences, the following sequence downstream of the HCV sequence (i.e., at the 3'-end):

25 5' CTT CTT TGG AGA AAG TGG TG 3'.

- The sequence common to each capture probe is complementary to a sequence in the binding partner(s), so that after hybridization, the duplex can be captured via affixation to the solid phase.

30 Also, in each set, each label probe contains, in addition to the sequences complementary to the HCV sequences, the following sequence downstream of the HCV sequence:

35 5' TTA GGC ATA GGA CCC GTG TC 3'.

If the method described in E.P.O. Publication No. 317,077 is used, the sequence common to each label probe is complementary to a sequence in a multimer, to allow hybrid duplex formation with that multimer.

The sequences of the probes in the above sets are shown in Fig. 19.

Detection of HCV Polynucleotide Sequences

Using PCR Amplification

In the generalized method for amplification of HCV RNA by cPCR it is contemplated that the RNA strand is a virion or mRNA strand, which is a "sense" strand. However, it is also possible that replicative intermediate forms may also be detected which would be "anti-sense"; in this case the primer would be "sense". An RNA sense strand containing the target region is hybridized with an anti-sense primer which primes the synthesis of the replicate strand containing the target. cDNA to the RNA template is synthesized with a primer- and template-dependent reverse transcriptase. The cDNA in the resulting RNA:cDNA hybrid is released by denaturation and treatment with RNase. Primers are annealed to the cDNA, and extended with a primer- and template-dependent DNA polymerase. The products are denatured, re-annealed to primers, and a second round of synthesis is conducted. A number of cycles are run until the amplified product containing the target region is in a desired amount, which is at least a detectable level.

Detection of Amplified HCV Nucleic Acid Sequences derived from HCV Nucleic Acid Sequences in Liver and Plasma Specimens from Chimpanzees with NANBH

HCV nucleic acids present in liver and plasma of chimpanzees with NANBH, and not in control chimpanzees, were amplified using essentially the polymerase chain re-

action (PCR) technique described by Saiki et al. (1986). The primer oligonucleotides were derived from the HCV cDNA sequences in clone 81 (Fig. 22), or clones 36 (Fig. 23) and 37b (Fig. 24). The amplified sequences were detected
5 by gel electrophoresis and a modified Southern blotting method, using as probes the appropriate cDNA oligomer or nick-translated cDNA sequence with a sequence from the region between, but not including, the two primers.

Samples of RNA containing HCV sequences to be
10 examined by the amplification system were isolated from liver biopsies of three chimpanzees with NANBH, and from two control chimpanzees. The isolation of the poly A⁺ RNA fraction was by the guanidinium thiocyanate procedure described in Maniatis et al. (1982).

15 Samples of RNA which were to be examined by the amplification system were also isolated from the plasmas of two chimpanzees with NANBH, and from one control chimpanzee, as well as from a pool of plasmas from control chimpanzees. One infected chimpanzee had a titer equal to
20 or greater than 10⁶ CID/ml, and the other infected chimpanzee had a titer equal to or greater than 10⁵ CID/ml.

The nucleic acids were extracted from the plasma as follows. Either 0.1 ml or 0.01 ml of plasma was
25 diluted to a final volume of 1.0 ml, with a TENB/proteinase K/SDS solution (0.05 M Tris-HCL, pH 8.0, 0.001 M EDTA, 0.1 M NaCl, 1 mg/ml Proteinase K, and 0.5% SDS) containing 10 micrograms/ml polyadenylic acid, and incubated at 37°C for 60 minutes. After this proteinase K
30 digestion, the resultant plasma fractions were deproteinized by extraction with TE (10.0 mM Tris-HCL, pH 8.0, 1 mM EDTA) saturated phenol. The phenol phase was separated by centrifugation, and was reextracted with TENB containing 0.1% SDS. The resulting aqueous phases from
35 each extraction were pooled, and extracted twice with an equal volume of phenol/chloroform/isoamyl alcohol

[1:1(99:1)], and then twice with an equal volume of a 99:1 mixture of chloroform/isoamyl alcohol. Following phase separation by centrifugation, the aqueous phase was brought to a final concentration of 0.2 M Na Acetate, and
5 the nucleic acids were precipitated by the addition of two volumes of ethanol. The precipitated nucleic acids were recovered by ultracentrifugation in a SW 41 rotor at 38 K, for 60 minutes at 4°C.

In addition to the above, the high titer
10 chimpanzee plasma and the pooled control plasma alternatively were extracted with 50 micrograms of poly A carrier by the procedure of Chomczynski and Sacchi (1987). This procedure uses an acid guanidinium thiocyanate extraction. RNA was recovered by centrifugation at 10,000
15 RPM for 10 minutes at 4°C in an Eppendorf microfuge.

On two occasions, prior to the synthesis of cDNA in the PCR reaction, the nucleic acids extracted from plasma by the proteinase K/SDS/phenol method were further purified by binding to and elution from S and S Elutip-R
20 Columns. The procedure followed was according to the manufacturer's directions.

The cDNA used as a template for the PCR reaction was derived from the nucleic acids (either total nucleic acids or RNA) prepared as described above. Following
25 ethanol precipitation, the precipitated nucleic acids were dried, and resuspended in DEPC treated distilled water. Secondary structures in the nucleic acids were disrupted by heating at 65°C for 10 minutes, and the samples were immediately cooled on ice. cDNA was synthesized using 1
30 to 3 micrograms of total chimpanzee RNA from liver, or from nucleic acids (or RNA) extracted from 10 to 100 microliters of plasma. The synthesis utilized reverse transcriptase, and was in a 25 microliter reaction, using the protocol specified by the manufacturer, BRL. The
35 primers for cDNA synthesis were those also utilized in the PCR reaction, described below. All reaction mixtures for

The primers for cDNA synthesis and for the PCR reactions were derived from the HCV cDNA sequences in either clone 81, clone 36, or clone 37b. (The HCV cDNA sequences of clones 81, 36, and 37b are shown in Figs. 22, 23, and 24, respectively.) The sequences of the two 16-mer primers derived from clone 81 were:

The sequence of the primer from clone 36 was:

The sequence of the primer from clone 37b was:

In the PCR reactions, the primer pairs consisted of either the two 16-mers derived from clone 81, or the 16-mer from clone 36 and the 16-mer from clone 37b.

The PCR reaction products were analyzed by separation of the products by alkaline gel electrophoresis, followed by Southern blotting, and detec-

tion of the amplified HCV-cDNA sequences with a ^{32}P -labeled internal oligonucleotide probe derived from a region of the HCV cDNA which does not overlap the primers. The PCR reaction mixtures were extracted with phenol/
5 chloroform, and the nucleic acids precipitated from the aqueous phase with salt and ethanol. The precipitated nucleic acids were collected by centrifugation, and dissolved in distilled water. Aliquots of the samples were subjected to electrophoresis on 1.8% alkaline agarose
10 gels. Single stranded DNA of 60, 108, and 161 nucleotide lengths were co-electrophoresed on the gels as molecular weight markers. After electrophoresis, the DNAs in the gel were transferred onto Biorad Zeta Probe[™] paper. Prehybridization and hybridization, and wash conditions
15 were those specified by the manufacturer (Biorad).

The probes used for the hybridization-detection of amplified HCV cDNA sequences were the following. When the pair of PCR primers were derived from clone 81, the probe was an 108-mer with a sequence corresponding to that
20 which is located in the region between the sequences of the two primers. When the pair of PCR primers were derived from clones 36 and 37b, the probe was the nick-translated HCV cDNA insert derived from clone 35, the nucleotide sequence of which is shown in Fig. 34. The
25 primers are derived from nucleotides 155-170 of the clone 37b insert, and 206-268 of the clone 36 insert. The 3'-end of the HCV cDNA insert in clone 35 overlaps nucleotides 1-186 of the insert in clone 36; and the 5'-end of clone 35 insert overlaps nucleotides 207-269 of the
30 insert in clone 37b. (Compare Figs. 23, 34 and 24.) Thus, the cDNA insert in clone 35 spans part of the region between the sequences of the clone 36 and 37b derived primers, and is useful as a probe for the amplified sequences which include these primers.

35 Analysis of the RNA from the liver specimens was according to the above procedure utilizing both sets of

primers and probes. The RNA from the liver of the three chimpanzees with NANBH yielded positive hybridization results for amplification sequences of the expected size (161 and 586 nucleotides for 81 and 36 and 37b, 5 respectively), while the control chimpanzees yielded negative hybridization results. The same results were achieved when the experiment was repeated three times.

Analysis of the nucleic acids and RNA from plasma was also according to the above procedure utilizing 10 the primers and probe from clone 81. The plasmas were from two chimpanzees with NANBH, from a control chimpanzee, and pooled plasmas from control chimpanzees. Both of the NANBH plasmas contained nucleic acids/RNA which yielded positive results in the PCR amplified assay, 15 while both of the control plasmas yielded negative results. These results have been repeatedly obtained several times.

Defective viruses have been known to occur in RNA viruses. By using PCR technology it is possible to 20 design primers to amplify sequences of the HCV genome. By analysis of the amplified products, it is expected to be able to identify both defective versions of the viral genome as well as wild-type viral species. Accordingly, using two primers based on known HCV sequence, one can 25 predict accurately the expected size of the PCR product. Any larger species observed by gel electrophoresis and hybridization analysis could represent potential variant genomes. Alternatively, any smaller species observed in this fashion might represent defective agents. Analyses 30 of these types would be useful in confirming the exact origin of the known HCV sequence, whether it is indeed a wild-type viral sequence or a defective genome. Techniques and methods for these analyses are well known in the art and have been previously described. This 35 methodology will enable one skilled in the art to obtain

related (wild-type or defective) forms of the viral genome.

Detection of Sequences in Captured Particles

5 Which When Amplified by PCR

Hybridize to HCV cDNA Derived from Clone 81

The RNA in captured particles was obtained as described below. The analysis for sequences which hybridize to the HCV cDNA derived from clone 81 was carried out
10 utilizing the PCR amplification procedure, as described supra., except that the hybridization probe was a kinased oligonucleotide derived from the clone 81 cDNA sequence. The results showed that the amplified sequences hybridized with the HCV cDNA probe.

15 Particles were captured from HCV infected chimpanzee plasma using polystyrene beads coated with an immunopurified antibody directed against the polypeptide encoded in clone 5-1-1. The procedure for producing the immunopurified antibody preparation is described in E.P.O.
20 Publication No. 318,216, which is commonly owned by the herein assignee, and which is incorporated herein by reference. Briefly, the HCV polypeptide encoded within clone 5-1-1 was expressed as a fusion polypeptide with superoxide dismutase (SOD). This was accomplished by
25 subcloning the clone 5-1-1 cDNA insert into the expression vector pSODcfl (Steimer et al. (1986)). DNA isolated from pSODcfl was treated with BamHI and EcoRI, and the following linker was ligated into the linear DNA created by the restriction enzymes:

30

5' GAT CCT GGA ATT CTG ATA AGA
CCT TAA GAC TAT TTT AA 3'

After cloning, the plasmid containing the insert was
35 isolated. Plasmid containing the insert was restricted with EcoRI. The HCV cDNA insert in clone 5-1-1 was

excised with EcoRI, and ligated into this EcoRI linearized plasmid DNA. The DNA mixture was used to transform E. coli strain D1210 (Sadler et al. (1980)). Recombinants with the 5-1-1 cDNA in the correct orientation for expression of the ORF were identified by restriction mapping and nucleotide sequencing. Recombinant bacteria from one clone were induced to express the SOD-NANB₅₋₁₋₁ polypeptide by growing the bacteria in the presence of IPTG. The fusion polypeptide was purified from the recombinant E. coli by differential extraction of the cell extracts with urea, followed by chromatography on anion and cation exchange columns. The purified SOD-NANB₅₋₁₋₁ polypeptide was attached to a nitrocellulose membrane. Antibody in samples of HCV infected serum was absorbed to the matrix-bound polypeptide. After washing to remove non-specifically bound materials and unbound materials, the bound antibody was released from the bound polypeptide.

20 cPCR Method to Detect HCV RNA in Liver
 and in Serum from Individuals with NANBH.

 The reliability and utility of a modified form of the PCR assay, i.e., a cPCR assay, for detecting HCV infection was determined by performing the assay on total liver RNA and on serum from infected individuals. In the cPCR assay, putative viral RNA in the sample is reverse transcribed into cDNA with reverse transcriptase; a segment of the resulting cDNA is then amplified utilizing a modified version of the PCR technique described by Saiki et al. (1986). The primers for the cPCR technique are derived from HCV RNA, which can be identified by the family of HCV cDNAs provided herein. Amplified product corresponding to the HCV-RNA is detected utilizing a probe derived from the family of HCV cDNAs provided herein.

35 The cPCR/HCV assay used in these studies were performed utilizing the following methods for the prepara-

tion of RNA, the reverse transcription of the RNA into cDNA, the amplification of specific segments of the cDNA by PCR, and the analysis of the PCR products.

RNA was extracted from liver utilizing the
5 guanidium isothiocyanate method for preparing total RNA described in Maniatis et al. (1982).

In order to isolate total RNA from plasma, the plasma was diluted five- to ten-fold with TENB (0.1 M NaCl, 50 mM Tris-HCl, pH 8.0, 1 mM EDTA) and incubated in
10 a Proteinase K/SDS solution (0.5% SDS, 1 mg/ml Proteinase K, 20 micrograms/ml Poly A carrier) for 60 to 90 minutes at 37°C. The samples were extracted once with phenol (pH 6.5), the resulting organic phase was re-extracted once
15 both extractions were pooled and extracted twice with an equal volume of phenol/CHCl₃/isoamyl alcohol [1:1(99:1)]. The resulting aqueous phases were extracted with an equal volume of CHCl₃/isoamyl alcohol (99:1) twice, and ethanol precipitated using 0.2 M sodium acetate, pH 6.5, and 2.5
20 volumes of 100% ethanol; precipitation was overnight at -20°C.

The cDNA used as a template for the PCR reaction was prepared utilizing the designated samples for preparation of the corresponding cDNAs. Each RNA sample
25 (containing either 2 micrograms of heat denatured total chimpanzee liver RNA, RNA from 2 microliters of plasma, or 10% of the RNA extracted from 10mm X 4 mm cylindrical human liver biopsies) was incubated in a 25 microliter reaction containing 1 micromolar of each primer, 1
30 millimolar of each deoxyribonucleotide triphosphate (dNTP), 50 millimolar Tris-HCL, pH 8.3, 5 millimolar MgCl₂, 5 millimolar dithiothreitol (DTT), 73 millimolar KCl, 40 units of RNase inhibitor (RNASIN), and 5 units of AMV reverse transcriptase. The incubation was for 60
35 minutes at 37°C. Following cDNA synthesis, the reactions

were diluted with 50 microliters of deionized water (DIW), boiled for 10 minutes, and cooled on ice.

Amplification of a segment of the HCV cDNA was performed utilizing two synthetic oligomer 16-mer primers whose sequences were derived from HCV cDNA clones 36 (anti-sense) and 37b (sense). The sequence of the primer from clone 36 was:

5' GCA TGT CAT GAT GTA T 3'.

10

The sequence of the primer from clone 37b was:

5' ACA ATA CGT GTG TCA C 3'.

15 The primers were used at a final concentration of 1 micromolar each. In order to amplify the segment of HCV cDNA which is flanked by the primers, the cDNA samples were incubated with 0.1 microgram of RNase A and the PCR reactants of the Perkin Elmer Cetus PCR kit (N801-0043 or
20 N801-0055) according to the manufacturer's instructions. The PCR reaction was performed for either 30 cycles or 60 cycles in a Perkin Elmer Cetus DNA thermal cycler. Each cycle consisted of a 1 minute denaturation step at 94°C, an annealing step of 2 minutes at 37°C, and an extension
25 step of 3 minutes at 72°C. However, the extension step in the final cycle (30 or 60) was 7 minutes rather than 3 minutes. After amplification the samples were extracted with an equal volume of phenol: chloroform (1:1), followed by extraction with an equal volume of chloroform, and then
30 the samples were precipitated with ethanol containing 0.2 M sodium acetate.

The cPCR products were analyzed as follows. The products were subjected to electrophoresis on 1.8% alkaline agarose gels according to Murakawa et al. (1988),
35 and transferred onto Zeta™ Probe paper (BioRad Corp.) by blotting gels overnight in 0.4 M NaOH. The blots were

neutralized in 2 X SSC (1 X SSC contains 0.15 M NaCl, 0.015 M sodium citrate), prehybridized in 0.3 M NaCl, 15 mM sodium phosphate buffer, pH 6.8, 15 mM EDTA, 1.0% SDS, 0.5% nonfat milk (Carnation Co.), and 0.5 mg/ml sonicated denatured salmon sperm DNA. The blots to be analyzed for HCV cDNA fragments were hybridized to a ³²P-labeled probe generated by nick translation of the HCV cDNA insert sequence in clone 35, described in E.P.O. Publication No. 318,216. After hybridization, the blots were washed in 0.1 X SSC (1 X SSC contains 0.15M NaCl, 0.01M Na citrate) at 65°C, dried, and autoradiographed. The expected product size is 586 nucleotides in length; products which hybridized with the probe and migrated in the gels in this size range were scored as positive for viral RNA.

As a control, cPCR primers designed to amplify alpha-1 anti-trypsin mRNA was performed to verify the presence of RNA in each sample analyzed. The coding region of the alpha-1 anti-trypsin gene is described in Rosenberg et al. (1984). Synthetic oligomer 16-mer primers designed to amplify a 365 nucleotide fragment of the coding region of the alpha-1 antitrypsin gene were derived from nucleotides 22-37 (sense) and nucleotides 372-387 (antisense). The PCR products were detected using a ³²P nick-translated probe which lies between, and not including, the cDNA/PCR primer sequences.

Due to the extreme sensitivity of the PCR reaction, all samples were run a minimum of three times. All false positive signals were eliminated when the following precautions were taken: 1) eliminating aerosols by using screw capped tubes with rubber O-ring seals; 2) pipetting with Ramin Microman™ positive displacement pipettors with disposable pistons/capillaries; and 3) selecting the oligonucleotide sequences for the cDNA and PCR primers from two non-contiguous cDNA clones.

Detection of HCV RNA in Liver Samples by a cPCR Method

The cPCR assay was performed on total RNA isolated from livers of three chimpanzees experimentally infected with a NANBH agent, and from liver biopsies of Italian patients diagnosed as having chronic NANBH.

Fig. 25A shows the results of the cPCR assay using 1 microgram of each preparation of total liver RNA. The RNA was isolated from liver samples of a chimpanzee in the chronic phase of NANBH (910) (lane 1), two chimpanzees in the acute phase of infection (1028 and 508) (lanes 2 and 3, respectively). PCR was performed on the samples in lanes 1-3 for 30 cycles and the autoradiogram of the blot containing those lanes was exposed for 5 hours. cDNA from 1 microgram of total RNA from acutely infected animal 1028 (lane 4), and three uninfected chimpanzees (lanes 5-7), were amplified for 60 cycles and the autoradiograms containing those lanes were exposed for 7 days. ³²P labeled MspI-digested pBR322 DNA served as markers on all the autoradiograms. It may be seen from the results that cDNA corresponding to HCV RNA was seen only in the samples from chimpanzees with NANBH, whether acute or chronic (lanes 1, 3, and 4). The cPCR products in these lanes migrated between marker fragments of 527 and 622 nucleotides (not shown).

Fig. 25B shows the results of the cPCR assay using 10% of the RNA extracted from 10mm X 4mm liver biopsy cylinders from 15 chronic NANB patients (lanes 1-15), one patient with cryptogenic liver disease (lane 16) and one control sample from a patient with chronic Hepatitis B (lane 17). Amplification by PCR was for 30 cycles and the autoradiogram for the blots were exposed for 4 days, except that lane 1 was exposed for 15 hours. As seen from the results, 9/15 (60%) of the human samples were positive for HCV RNA (lanes 1, 2, 4, 6, 7, 10-13). One patient diagnosed with cryptogenic liver disease (lane 16)

and one patient with a chronic HBV infection (lane 17) were repeatedly negative in the cPCR assay.

Comparison of the HCV/cPCR Assay on Human Liver Biopsies
and RIA of Serum Using HCV C100-3 Polypeptide

5 SOD/HCV C100-3 polypeptide (also called C100) is a recombinant fusion polypeptide which contains 363 viral amino acids. The polypeptide is useful for detecting antibodies to HCV (See Kuo et al. (1989)). The method for
10 preparing C100 is described in E.P.O. Publication No. 318,216.

Radioimmune assay using C100 was performed on the sera collected from the same 17 human patients whose liver samples were subjected to HCV/cPCR assay as
15 described supra. The sera was collected on the same day as the liver biopsies. The assay was performed essentially as described in E.P.O. Publication No. 318,216, which is commonly owned and incorporated herein by reference. Briefly, Microtiter plates (Immulon 2, Removeawell
20 strips) were coated with 0.1 microgram of purified C100. The coated plates were incubated for 1 hour at 37°C with the serum samples (100 microliters of a 1:100 dilution) or appropriate controls. After incubation, the unbound material was removed, the plates were washed, and
25 complexes of human antibody-C100 were detected by incubation with ¹²⁵I-labeled sheep anti-human immunoglobulin. Unbound labeled antibody was removed by aspiration, and the plates were washed. The radioactivity in individual wells was determined.

30 The results of the RIA showed that sixty-seven percent of these samples were positive for anti-C100 antibodies. Sera from the patient diagnosed with cryptogenic liver disease was positive for anti-C100 antibodies, although the levels of viral RNA were undetectable in the
35 patient's liver in this sample. The level of correlation between the presence of anti-C100 antibodies and HCV RNA

was seventy percent; two patients who were negative for antibodies by RIA had significant levels of HCV RNA in their livers (data not shown).

The results indicate that virus is frequently present in the liver of patients with circulating anti-C100 antibodies, and confirms claims that the presence of anti-C100 antibodies accurately reflects exposure to HCV. Moreover, taken together, these results indicate that HCV of this type accounts for NANBH in at least 75% of the patients in this study, and that the predominant strain of HCV in Italy appears to be closely related to the strain of HCV prevalent in the United States.

HCV/cPCR Assay of Sera: Detection of Viral RNA
in Acute Phase Infection in Chimpanzees

The temporal relationship between the display of liver damage, the presence of HCV RNA, and the presence of anti-HCV antibodies was monitored in serum from two experimentally infected chimpanzees with NANBH (nos. 771 and 910). Liver damage was determined by alanine amino transferase (ALT) levels; the presence of HCV RNA was determined by the HCV cPCR assay described above; anti-HCV antibodies were detected utilizing the C100 RIA.

The HCV/cPCR analysis was performed on RNA extracted from 1 microliter of chimpanzee plasma. Serum was taken from chimpanzee 771 on days 25, 32, 70 and 88 post-infection; cPCR was performed for 30 cycles and the autoradiogram was exposed for 18 days. Serum was taken from chimpanzee 910 on days 11, 28, and 67 post-infection; cPCR was performed for 60 cycles and the autoradiogram was exposed for 5 days.

The results of the assays are shown in Fig. 26A for chimpanzee 771, and Fig. 26B for chimpanzee 910. From a comparison of Figs. 26A and 26B, it appears that an early, well defined peak of ALT values during acute

hepatitis correlates with the presence of viral RNA in the infected individual.

The data also indicate that the presence of HCV RNA, which is indicative of a state of viremia, precedes the presence of anti-HCV antibodies. Chimpanzee 771 (Fig. 26A) exhibited a clearly defined acute episode of post-transfusion NANBH at 28 days, as characterized by an initial peak of ALT levels. HCV RNA was detected in the serum collected at day 25, and at day 32. However, during this acute phase, anti-HCV antibodies were absent. In contrast, at day 70 HCV RNA was below the experimental level of detection, and anti-HCV antibodies were rising. At day 88, HCV RNA remained undetectable, while anti-HCV antibodies were significantly increased over that of day 70.

The results obtained from the sera of chimpanzee 910 were somewhat similar in pattern, although the time of HCV antibodies induced by the infection were not detected during the acute phase of the disease, which extended to at least day 67; the anti-HCV antibodies detected by RIA at day 11 were due to passive immunization of animal 910 with antibodies from the plasma used to inoculate the animal. Anti-HCV antibodies were found in chimpanzee 910 serum during the later, chronic phase of the infection (data not shown).

It should be noted that low ALT values in plasma from individuals with chronic NANBH do not necessarily correlate with weak virus production. A pool of 17 different plasma samples taken from chimpanzee 910 over a period of two to three and one-half years post inoculation was monitored for ALT levels and for HCV RNA. The ALT values of the samples did not exceed 45 mU/ml; nevertheless, titration studies indicated high titers of HCV (3×10^6 CID/ml). cPCR was carried out for 30 cycles, and the autoradiogram was exposed for 15 hours; the cPCR analysis clearly showed the presence of viral RNA (data not shown).

HCV/cPCR Assay of Sera: Detection of Viral RNA
in Acute Phase Infection in Humans

Plasma from a human surgical patient collected
5 during early acute NANBH was examined for HCV RNA and for
anti-HCV antibodies, utilizing the HCV/cPCR assay and
C100-RIA, respectively. The HCV/cPCR assay was conducted
utilizing 1 microliter of plasma from the patient, and
from four human controls with known pedigrees; cPCR was
10 performed for thirty cycles, and after hybridization and
washing the autoradiogram was exposed for eight hours.

The results showed that the serum collected from
the surgical patient during the acute phase of infection
contained a high level of viral RNA, and that anti-HCV
15 antibodies were not detectable by the C100-RIA (data now
shown). (The acute phase plasma from the surgical patient
was known to have a high titer of NANBH infectious agent
[$10^{6.5}$ CID/ml, as determined by Feinstone et al. (1981);
Feinstone et al. (1983)]). It should be noted, however,
20 that this patient did sero-convert to anti-HCV antibodies
by the C100-RIA approximately 9 months after infection.
The serum from the pedigreed human control plasmas were
negative in both the HCV/cPCR assay and C100-RIA.

25 Sensitivity of HCV/cPCR Assay

The sensitivity of the HCV/cPCR assay was
determined by analyzing ten-fold serial dilutions of a
plasma pool of known titer. The chimpanzee plasma had a
titer of $\sim 3 \times 10^5$ CID/ml, and RNA was extracted from ten-
30 fold dilutions of 1 microliter of the plasma. cPCR was
performed for 30 cycles, and after hybridization and wash-
ing, the autoradiogram was exposed for 15 hours. The cPCR
products resulting from amplification of ~ 300 , ~ 30 , and ~ 3
CID of HCV genomes are shown in lanes 1-3, respectively of
35 Fig. 29. The samples in lanes 1 and 2 were detectable on
autoradiograms exposed for 2 hours.

Since the average titer of HCV in infected individuals is believed to be between approximately 100 to 10,000 CID/ml of plasma, this data suggests that the HCV/cPCR assay may be clinically useful.

5

HCV/cPCR Assay for Variant HCV Strains

Primers, consisting of a set of oligomer 44-mers and a set of oligomer 45-mers, were designed to amplify strains of HCV which are similar or identical to the HCV isolate from which the cDNA sequence in Fig. 18 is derived. The premise underlying the design of these primers is our discovery that HCV is a Flavi-like virus. Members of the Flaviviridae family, when compared to HCV, have two major conserved sets of amino acid sequences, TATPPG and QRRGR, in the putative NS3 region of these viruses. Several other smaller sets may be seen, for example, GDD in the putative NS5 region. Other sets are determinable by comparison of the known amino acid sequences with that of HCV. This information was deduced from the sequences for several members of Flaviviridae which have been described, including Japanese Encephalitis Virus (Sumiyoshi et al. (1987)), Yellow Fever Virus (Rice et al. (1985)), Dengue Type 2 Virus (Hahn et al. (1988)), Dengue Type 4 Virus (Mackow (1987)), and West Nile Virus (Castle et al. (1986)). The conserved amino acid sequences and codon utilization are in the table immediately following.

30

35

Conserved Amino Acid (A.A.) Sequences
Among Flaviviruses and HCV

5	Virus	# of first A.A.	A.A.					
			T	A	T	P	G	
	HCV	1348	5' ACC	GCC	ACC	CCT	CCG	GCC 3'
	Yellow Fever	1805	ACA	GCC	ACA	CCG	CCT	GGG
	West Nile	1818	ACG	GCA	ACG	CCA	CCC	GGG
	Dengue-4	1788	ACC	GCA	ACC	CCT	CCC	GGA
	JEV	1957	ACA	GCG	ACC	CCG	CCT	GGA

HCV sense primer (44mer)=

10 5' ACC GCC ACC CCX CC 3'
(X = A, T, C, or G)

15	Virus	# of first A.A.	A.A.				
			Q	R	R	G	R
	HCV	1486	5' CAA	CGT	CGG	GGC	AGG 3'
	Yellow Fever	1946	CAA	AGG	AGG	GGG	CGC
	West Nile	1959	CAG	CGG	AGA	GGA	CGC
	Dengue-4	1929	CAG	AGA	AGA	GGG	CGA
	JEV	1820	CAA	CGG	AGG	GGC	AGA

HCV antisense primer (45mer)=

20 3' GTX GCA GCC CCG TCC 5'
(X = T or C)

25 Note: the primer sequence was chosen to minimize the number of nucleotide degeneracies at the 3'-end of the primer sequence and to maximize the number of nucleotides at the 3'-end of each primer which exactly match any of the possible nucleotide sequences, or the complement thereof, encoding the conserved amino acids indicated above.

30 The 44-mer and 45-mer oligomer primers were designed so that the sequences encoding these amino acids were incorporated within the primer. Moreover, they contain degeneracies at the 3'-end of each primer, and are derived from two different regions of the HCV genome which are present in clone 40b (See Fig. 28), and which are derived from the region encoding putative NS3 of HCV. The formulae for the oligonucleotide primers in the sets are:

35

5' GAC TGC GGG GGC GAG ACT GGT TGT GCT CGC
ACC GCC ACC CCX CC 3'

5 where X is A,T,G, or C; and

5' TCT GTA GAT GCC TGG CTT CCC CCT GCC AGT
CCT GCC CCG ACT YTG 3'

10 where Y is T or C.

The HCV/cPCR assay was carried out utilizing these primers to amplify HCV RNA in chimpanzee 910 plasma. The assay method was essentially as described in Section supra., except that the 44-mer and 45-mer sets of oligomer
15 primers were substituted for the primers derived from clone 36 and clone 37b. In addition, detection of amplified HCV cDNA was by hybridization with a probe derived from clone 40a, the sequence of which is shown in Fig. 32.

The probe was prepared by amplifying a segment
20 of clone 40a utilizing the PCR method described supra., and 18-mer primers containing the following sequences:

5' GAG ACA TCT CAT CTT CTG 3'

25 and

5' GAG CGT GAT TGT CTC AAT 3'.

After amplification, the probe preparation was labeled
30 with ³²P by nick translation.

Fig. 33 shows an autoradiograph of the Southern blots probed with the sequence derived from Clone 40a. ³²P labeled MspI digested pBR322 DNA fragments served as markers (lane 1). The predicted size of the PCR product
35 resulting from amplification using these primers is 490

nucleotides (nt). Duplicate reactions are shown in lanes 2 and 3.

Analysis for Variants of the 5'-Region of HCV

5 Based upon the Flavivirus model, the 5'-region
HCV cDNA which is flanked by the regions represented in
clones ag30a and k9-1 encodes a segment of putative
envelope and/or matrix protein(s) (E/M). Serum obtained
from the chimpanzee from which the HCV cDNA "c" library,
10 was constructed was analyzed by HCV/cPCR to determine
whether variants within this target region were present.

 The HCV/cPCR assay was performed essentially as
described supra., for the isolation of clone 5'-32, except
for the primers and probes used. Fig. 37 shows the
15 relationship of the primers and probes (and the clones
from which they were derived) to that of the target region
of HCV cDNA. One set of PCR primers, ag30a16A and
K91Env16B, were derived from clones ag30a and k9-1, which
are upstream and downstream, respectively, of the target
20 sequence. The expected size of the cPCR product primed by
ag30a16A and K91Env16B is 1.145 kb based upon the
confirmed sequence of HCV cDNA. Two other sets of PCR
primers covering the region amplified using ag30a16A and
K91Env16B, and overlapping each other were also used for
25 PCR amplification of HCV RNA in the serum. Thus, in this
case the PCR reactions were run using as one set of prim-
ers ag30a16A and CA156e16B, and as the second set of prim-
ers CA156e16A and k91Env16B. The expected PCR product
sizes for these pairs were 615 nucleotides (NT) and 683
30 NT, respectively. The table immediately following lists
the primer, the clone from which it was derived, and the
primer sequence.

Table

	Primer	Clone	Sequence
5	ag30a16A	ag30a	5' CTC TAT GGC AAT GAG G 3'
	K91Env16B	k9-1	5' CGT TGG CAT AAC TGA T 3'
	CA156e16B	156	5' CGA CAA GAA AGA CAG A 3'
	CA156e16A	156	5' AGC TTC GAC GTC ACA T 3'
	CA216a16A	216	5' TGA ACT ATG CAA CAG G 3'
10	CA216a16B	216	5' GGA GTG TGC AGG ATG G 3'
	CA84a16A	84	5' AAG GTT GCA ATT GCT C 3'
	CA84a16B	84	5' ACT AAC AGG ACC TTC G 3'

The probes for all of the HCV/cPCR products consisted of ³²P labeled sections of HCV cDNA which had been prepared by PCR amplification of a region of clone 216 (using CA216a16A and 216a16B as primers), and of clone 84 (using CA84a16A and CA84a16B as primers); ³²P was introduced into the PCR products by nick translation. These probes did not overlap the primers used in the HCV/cPCR reactions.

Fig. 38 shows an autoradiograph of a Southern blot in which the HCV/cPCR products were hybridized with the ³²P-labeled probes. The HCV/cPCR product extended from primers ag30a16A and K91Env16B (lane 1) was approximately 1.1Kb; no other PCR products were observed in a 15 hour exposure. The HCV products extended from the primer sets ag30a15A/CA156e16B (lane 2) and CA156e16A/K91Env16B (lane 3) were approximately 625NT and approximately 700 NT, respectively. The size of the PCR products were determined by comparison with the relative migrations of fragments resulting from the digestion of pBR322 with MspI and of PhiX 174 digested with HaeIII (lane 5).

The above study will detect insertions or deletions as small as approximately 20NT to 50NT and DNA rearrangements altering the size of the target DNA. The

results in Fig. 38 confirm that there is only 1 major species of cDNA derived from the E/M region of the HCV in the chimpanzee serum.

5 Amplification for Cloning of HCV cDNA Sequences
 Utilizing the PCR and Primers Derived from
 Conserved Regions of Flavivirus Genomic Sequences

Our discovery that HCV is a flavi-like virus, allows a strategy for cloning uncharacterized HCV cDNA
10 sequences utilizing the PCR technique, and primers derived from the regions encoding conserved amino acid sequences in flaviviruses. Generally, one of the primers is derived from a defined HCV genomic sequence, and the other primer which flanks a region of unsequenced HCV polynucleotide is
15 derived from a conserved region of the flavivirus genome. The flavivirus genomes are known to contain conserved sequences within the NS1, and E polypeptides, which are encoded in the 5'-region of the flavivirus genome. Thus, to isolate cDNA sequences derived from putatively
20 comparable regions of the HCV genome, upstream primers are designed which are derived from the conserved sequences within these flavivirus polypeptides. The downstream primers are derived from an upstream end of the known portion of the HCV cDNA.

25 Because of the degeneracy of the code, it is probable that there will be mismatches between the flavivirus probes and the corresponding HCV genomic sequence. Therefore a strategy which is similar to the one described by Lee (1988) is used. The Lee procedure
30 utilizes mixed oligonucleotide primers complementary to the reverse translation products of an amino acid sequence; the sequences in the mixed primers takes into account every codon degeneracy for the conserved amino acid sequence.

35 Three sets of primer mixes are generated, based on the amino acid homologies found in several

-111-

flaviviruses, including Dengue-2,4 (D-2,4), Japanese Encephalitis Virus (JEV), Yellow Fever (YF), and West Nile Virus (WN). The primer mixture derived from the most upstream conserved sequence (5'-1), is based upon the amino acid sequence gly-trp-gly, which is part of the conserved sequence asp-arg-gly-trp-gly-aspN found in the E protein of D-2, JEV, YF, and WN. The next primer mixture (5'-2) is based upon a downstream conserved sequence in E protein, phe-asp-gly-asp-ser-tyr-ileu-phe-gly-asp-ser-tyr-ileu, and is derived from phe-gly-asp; the conserved sequence is present in D-2, JEV, YF, and WN. The third primer mixture (5'-3), is based on the amino acid sequence arg-ser-cys, which is part of the conserved sequence cys-cys-arg-ser-cys in the NS1 protein of D-2, D-4, JEV, YF, and WN. The individual primers which form the mixture in 5'-3 are shown in Fig. 53. In addition to the varied sequences derived from conserved region, each primer in each mixture also contains a constant region at the 5'-end which contains a sequence encoding sites for restriction enzymes, HindIII, MboI, and EcoRI.

The downstream primer, ssc5h20A, is derived from a nucleotide sequence in clone 5h, which contains HCV cDNA with sequences with overlap those in clones 14i and 11b. The sequence of ssc5h20A is

5' GTA ATA TGG TGA CAG AGT CA 3'.

An alternative primer, ssc5h34A, may also be used. This primer is derived from a sequence in clone 5h, and in addition contains nucleotides at the 5'-end which create a restriction enzyme site, thus facilitating cloning. The sequence of ssc5h34A is

5' GAT CTC TAG AGA AAT CAA TAT GGT GAC AGA GTC A 3'.

The PCR reaction, which was initially described by Saiki et al. (1986), is carried out essentially as described in Lee et al. (1988), except that the template for the cDNA is RNA isolated from HCV infected chimpanzee liver, or from viral particles isolated from HCV infected chimpanzee serum. In addition, the annealing conditions are less stringent in the first round of amplification (0.6M NaCl, and 25°C), since the part of the primer which will anneal to the HCV sequence is only 9 nucleotides, and there could be mismatches. Moreover, if ssc5h34A is used, the additional sequences not derived from the HCV genome tend to destabilize the primer-template hybrid. After the first round of amplification, the annealing conditions can be more stringent (0.066M NaCl, and 32°C-37°C), since the amplified sequences now contain regions which are complementary to, or duplicates of the primers. In addition, the first 10 cycles of amplification are run with Klenow enzyme I, under appropriate PCR conditions for that enzyme. After the completion of these cycles, the samples are extracted, and run with Taq polymerase, according to kit directions, as furnished by Cetus/Perkin-Elmer.

After the amplification, the amplified HCV cDNA sequences are detected by hybridization using a probe derived from clone 5h. This probe is derived from sequences upstream of those used to derive the primer, and does not overlap the sequences of the clone 5h derived primers. The sequence of the probe is

5' CCC AGC GGC GTA CGC GCT GGA CAC GGA GGT GGC CGC GTC
GTG TGG CGG TGT TGT TCT CGT CGG GTT GAT GGC GC 3'.

Industrial Applicability

The methods described herein, as well as the oligomers, both probes and primers, derived from HCV cDNA, and kits containing them, are useful for the accurate, relatively simple, and economic determination of the presence of HCV in biological samples, more particularly in blood which may be used for transfusions, and in individuals suspected of having HCV an infection. Moreover, these methods and oligomers may be useful for detecting an earlier stage of HCV infection than are immunological assays based upon the use of a recombinant HCV polypeptides. Also, an amplified polynucleotide hybridization assay detects HCV RNA in occasional samples which are anti-HCV antibody negative. Thus, the probes and primers described herein may be used amplified hybridization assays, in conjunction with an immunoassays based on HCV polypeptides to more completely identify infections due to HCV, and HCV-infected biological specimens, including blood.

The information provided herein allows the design of primers and/or probes which are derived from conserved regions of the HCV genome. The provision of these primers and probes makes available a general method which will detect variant HCV strains, and which will be of use in the screening of blood and blood products.

If the primers used in the method are derived from conserved regions of the HCV genome, the method should aid in the detection and/or identification of variant strains of HCV. This, in turn, should lead to the development of additional immunological reagents for the detection and diagnosis of HCV, as well as the development of additional polynucleotide reagents for detection and or treatment of HCV.

In addition, sets of primers and probes designed from the conserved amino acid sequences of Flaviviruses

and HCV allow for a universal detection method for these infectious agents.

The following listed materials are on deposit under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC), 12301 Parklawn Dr., Rockville, Maryland 20852, and have been assigned the following Accession Numbers.

	<u>lambda-gt11</u>	<u>ATCC No.</u>	<u>Deposit Date</u>
10	HCV cDNA library	40394	1 Dec. 1987
	clone 81	40388	17 Nov. 1987
	clone 91	40389	17 Nov. 1987
	clone 1-2	40390	17 Nov. 1987
	clone 5-1-1	40391	18 Nov. 1987
15	clone 12f	40514	10 Nov. 1988
	clone 35f	40511	10 Nov. 1988
	clone 15e	40513	10 Nov. 1988
	clone K9-1	40512	10 Nov. 1988
	JSC 308	20879	5 May 1988
20	pS356	67683	29 April 1988

In addition, the following deposits were made on 11 May 1989.

	<u>Strain</u>	<u>Linkers</u>	<u>ATCC No.</u>
25	D1210 (Cf1/5-1-1)	EF	67967
	D1210 (Cf1/81)	EF	67968
	D1210 (Cf1/CA74a)	EF	67969
	D1210 (Cf1/35f)	AB	67970
30	D1210 (Cf1/279a)	EF	67971
	D1210 (Cf1/C36)	CD	67972
	D1210 (Cf1/13i)	AB	67973
	D1210 (Cf1/C33b)	EF	67974
	D1210 (Cf1/CA290a)	AB	67975
35	HB101 (AB24/C100 #3R)		67976

The following derivatives of strain D1210 were deposited on 3 May 1989.

	<u>Strain Derivative</u>	<u>ATCC No.</u>
5	pCF1CS/C8f	67956
	pCF1AB/C12f	67952
	pCF1EF/14c	67949
	pCF1EF/15e	67954
	pCF1AB/C25c	67958
10	pCF1EF/C33c	67953
	pCF1EF/C33f	67050
	pCF1CD/33g	67951
	pCF1CD/C39c	67955
	pCF1EF/C40b	67957
15	pCF1EF/CA167b	67959

The following strains were deposited on May 12, 1989.

	<u>Strain</u>	<u>ATCC No.</u>
20	Lambda gt11(C35)	40603
	Lambda gt10(beta-5a)	40602
	D1210 (C40b)	67980
	D1210 (M16)	67981

25 The following biological materials were deposited on March 23, 1990.

	<u>Material</u>	<u>ATCC No.</u>
30	5'-clone32 (in pUC18S)	68276

35

CLAIMS

1. An oligomer capable of hybridizing to an HCV
 5 sequence in an analyte polynucleotide strand, wherein the
 oligomer is comprised of an HCV targeting sequence com-
 plementary to at least 4 contiguous nucleotides of HCV
 cDNA shown in Fig. 18.

10 2. The oligomer of claim 1, wherein the target-
 ing sequence is comprised of nucleotides which are com-
 plementary to nucleotides selected from the following HCV
 cDNA nucleotides shown in Fig. 18, (nn_x - nn_y denotes
 nucleotide number x to nucleotide number y)):

15
 nn-340 - nn-330; nn-330 - nn-320; nn-320 - nn-310;
 nn-310 - nn-300; nn-300 - nn-290; nn-290 - nn-280;
 nn-280 - nn-270; nn-270 - nn-260; nn-260 - nn-250;
 nn-250 - nn-240; nn-240 - nn-230; nn-230 - nn-220;
 20 nn-220 - nn-210; nn-210 - nn-200; nn-200 - nn-190;
 nn-190 - nn-180; nn-180 - nn-170; nn-170 - nn-160;
 nn-160 - nn-150; nn-150 - nn-140; nn-140 - nn-130;
 nn-130 - nn-120; nn-120 - nn-110; nn-110 - nn-100;
 nn-100 - nn-90; nn-90 - nn-80; nn-80 - nn-70;
 25 nn-70 - nn-60; nn-60 - nn-50; nn-50 - nn-40;
 nn-40 - nn-30; nn-30 - nn-20; nn-20 - nn-10;
 nn-10 - nn₁; nn₁ - nn₁₀; nn₁₀ - nn₂₀; nn₂₀ - nn₃₀;
 nn₃₀ - nn₄₀; nn₄₀ - nn₅₀; nn₅₀ - nn₆₀; nn₆₀ - nn₇₀;
 nn₇₀ - nn₈₀; nn₈₀ - nn₉₀; nn₉₀ - nn₁₀₀; nn₁₀₀ - nn₁₁₀;
 30 nn₁₁₀ - nn₁₂₀; nn₁₂₀ - nn₁₃₀; nn₁₃₀ - nn₁₄₀;
 nn₁₄₀ - nn₁₅₀; nn₁₅₀ - nn₁₆₀; nn₁₆₀ - nn₁₇₀;
 nn₁₇₀ - nn₁₈₀; nn₁₈₀ - nn₁₉₀; nn₁₉₀ - nn₂₀₀;
 nn₂₀₀ - nn₂₁₀; nn₂₁₀ - nn₂₂₀; nn₂₂₀ - nn₂₃₀;
 nn₂₃₀ - nn₂₄₀; nn₂₄₀ - nn₂₅₀; nn₂₅₀ - nn₂₆₀;
 35 nn₂₆₀ - nn₂₇₀; nn₂₇₀ - nn₂₈₀; nn₂₈₀ - nn₂₉₀;
 nn₂₉₀ - nn₃₀₀; nn₃₀₀ - nn₃₁₀; nn₃₁₀ - nn₃₂₀;

nn320 - nn330; nn330 - nn340; nn340 - nn350;
nn350 - nn360; nn360 - nn370; nn370 - nn380;
nn380 - nn390; nn390 - nn400; nn400 - nn410;
nn410 - nn420; nn420 - nn430; nn430 - nn440;
5 nn440 - nn450; nn450 - nn460; nn460 - nn470;
nn470 - nn480; nn480 - nn490; nn490 - nn500;
nn500 - nn510; nn510 - nn520; nn520 - nn530;
nn530 - nn540; nn540 - nn550; nn550 - nn560;
nn560 - nn570; nn570 - nn580; nn580 - nn590;
10 nn590 - nn600; nn600 - nn610; nn610 - nn620;
nn620 - nn630; nn630 - nn640; nn640 - nn650;
nn650 - nn660; nn660 - nn670; nn670 - nn680;
nn680 - nn690; nn690 - nn700; nn700 - nn710;
nn710 - nn720; nn720 - nn730; nn730 - nn740;
15 nn740 - nn750; nn750 - nn760; nn760 - nn770;
nn770 - nn780; nn780 - nn790; nn790 - nn800;
nn800 - nn810; nn810 - nn820; nn820 - nn830;
nn830 - nn840; nn840 - nn850; nn850 - nn860;
nn860 - nn870; nn870 - nn880; nn880 - nn890;
20 nn890 - nn900; nn900 - nn910; nn910 - nn920;
nn920 - nn930; nn930 - nn940; nn940 - nn950;
nn950 - nn960; nn960 - nn970; nn970 - nn980;
nn980 - nn990; nn990 - nn1000; nn1000 - nn1010;
nn1010 - nn1020; nn1020 - nn1030; nn1030 - nn1040;
25 nn1040 - nn1050; nn1050 - nn1060; nn1060 - nn1070;
nn1070 - nn1080; nn1080 - nn1090; nn1090 - nn1100;
nn1100 - nn1110; nn1110 - nn1120; nn1120 - nn1130;
nn1130 - nn1140; nn1140 - nn1150; nn1150 - nn1160;
nn1160 - nn1170; nn1170 - nn1180; nn1180 - nn1190;
30 nn1190 - nn1200; nn1200 - nn1210; nn1210 - nn1220;
nn1220 - nn1230; nn1230 - nn1240; nn1240 - nn1250;
nn1250 - nn1260; nn1260 - nn1270; nn1270 - nn1280;
nn1280 - nn1290; nn1290 - nn1300; nn1300 - nn1310;
nn1310 - nn1320; nn1320 - nn1330; nn1330 - nn1340;
35 nn1340 - nn1350; nn1350 - nn1360; nn1360 - nn1370;
nn1370 - nn1380; nn1380 - nn1390; nn1390 - nn1400;

nn₁₄₀₀ - nn₁₄₁₀; nn₁₄₁₀ - nn₁₄₂₀; nn₁₄₂₀ - nn₁₄₃₀;
nn₁₄₃₀ - nn₁₄₄₀; nn₁₄₄₀ - nn₁₄₅₀; nn₁₄₅₀ - nn₁₄₆₀;
nn₁₄₆₀ - nn₁₄₇₀; nn₁₄₇₀ - nn₁₄₈₀; nn₁₄₈₀ - nn₁₄₉₀;
nn₁₄₉₀ - nn₁₅₀₀; nn₁₅₀₀ - nn₁₅₁₀; nn₁₅₁₀ - nn₁₅₂₀;
5 nn₁₅₂₀ - nn₁₅₃₀; nn₁₅₃₀ - nn₁₅₄₀; nn₁₅₄₀ - nn₁₅₅₀;
nn₁₅₅₀ - nn₁₅₆₀; nn₁₅₆₀ - nn₁₅₇₀; nn₁₅₇₀ - nn₁₅₈₀;
nn₁₅₈₀ - nn₁₅₉₀; nn₁₅₉₀ - nn₁₆₀₀; nn₁₆₀₀ - nn₁₆₁₀;
nn₁₆₁₀ - nn₁₆₂₀; nn₁₆₂₀ - nn₁₆₃₀; nn₁₆₃₀ - nn₁₆₄₀;
nn₁₆₄₀ - nn₁₆₅₀; nn₁₆₅₀ - nn₁₆₆₀; nn₁₆₆₀ - nn₁₆₇₀;
10 nn₁₆₇₀ - nn₁₆₈₀; nn₁₆₈₀ - nn₁₆₉₀; nn₁₆₉₀ - nn₁₇₀₀;
nn₁₇₀₀ - nn₁₇₁₀; nn₁₇₁₀ - nn₁₇₂₀; nn₁₇₂₀ - nn₁₇₃₀;
nn₁₇₃₀ - nn₁₇₄₀; nn₁₇₄₀ - nn₁₇₅₀; nn₁₇₅₀ - nn₁₇₆₀;
nn₁₇₆₀ - nn₁₇₇₀; nn₁₇₇₀ - nn₁₇₈₀; nn₁₇₈₀ - nn₁₇₉₀;
nn₁₇₉₀ - nn₁₈₀₀; nn₁₈₀₀ - nn₁₈₁₀; nn₁₈₁₀ - nn₁₈₂₀;
15 nn₁₈₂₀ - nn₁₈₃₀; nn₁₈₃₀ - nn₁₈₄₀; nn₁₈₄₀ - nn₁₈₅₀;
nn₁₈₅₀ - nn₁₈₆₀; nn₁₈₆₀ - nn₁₈₇₀; nn₁₈₇₀ - nn₁₈₈₀;
nn₁₈₈₀ - nn₁₈₉₀; nn₁₈₉₀ - nn₁₉₀₀; nn₁₉₀₀ - nn₁₉₁₀;
nn₁₉₁₀ - nn₁₉₂₀; nn₁₉₂₀ - nn₁₉₃₀; nn₁₉₃₀ - nn₁₉₄₀;
nn₁₉₄₀ - nn₁₉₅₀; nn₁₉₅₀ - nn₁₉₆₀; nn₁₉₆₀ - nn₁₉₇₀;
20 nn₁₉₇₀ - nn₁₉₈₀; nn₁₉₈₀ - nn₁₉₉₀; nn₁₉₉₀ - nn₂₀₀₀;
nn₂₀₀₀ - nn₂₀₁₀; nn₂₀₁₀ - nn₂₀₂₀; nn₂₀₂₀ - nn₂₀₃₀;
nn₂₀₃₀ - nn₂₀₄₀; nn₂₀₄₀ - nn₂₀₅₀; nn₂₀₅₀ - nn₂₀₆₀;
nn₂₀₆₀ - nn₂₀₇₀; nn₂₀₇₀ - nn₂₀₈₀; nn₂₀₈₀ - nn₂₀₉₀;
nn₂₀₉₀ - nn₂₁₀₀; nn₂₁₀₀ - nn₂₁₁₀; nn₂₁₁₀ - nn₂₁₂₀;
25 nn₂₁₂₀ - nn₂₁₃₀; nn₂₁₃₀ - nn₂₁₄₀; nn₂₁₄₀ - nn₂₁₅₀;
nn₂₁₅₀ - nn₂₁₆₀; nn₂₁₆₀ - nn₂₁₇₀; nn₂₁₇₀ - nn₂₁₈₀;
nn₂₁₈₀ - nn₂₁₉₀; nn₂₁₉₀ - nn₂₂₀₀; nn₂₂₀₀ - nn₂₂₁₀;
nn₂₂₁₀ - nn₂₂₂₀; nn₂₂₂₀ - nn₂₂₃₀; nn₂₂₃₀ - nn₂₂₄₀;
nn₂₂₄₀ - nn₂₂₅₀; nn₂₂₅₀ - nn₂₂₆₀; nn₂₂₆₀ - nn₂₂₇₀;
30 nn₂₂₇₀ - nn₂₂₈₀; nn₂₂₈₀ - nn₂₂₉₀; nn₂₂₉₀ - nn₂₃₀₀;
nn₂₃₀₀ - nn₂₃₁₀; nn₂₃₁₀ - nn₂₃₂₀; nn₂₃₂₀ - nn₂₃₃₀;
nn₂₃₃₀ - nn₂₃₄₀; nn₂₃₄₀ - nn₂₃₅₀; nn₂₃₅₀ - nn₂₃₆₀;
nn₂₃₆₀ - nn₂₃₇₀; nn₂₃₇₀ - nn₂₃₈₀; nn₂₃₈₀ - nn₂₃₉₀;
nn₂₃₉₀ - nn₂₄₀₀; nn₂₄₀₀ - nn₂₄₁₀; nn₂₄₁₀ - nn₂₄₂₀;
35 nn₂₄₂₀ - nn₂₄₃₀; nn₂₄₃₀ - nn₂₄₄₀; nn₂₄₄₀ - nn₂₄₅₀;
nn₂₄₅₀ - nn₂₄₆₀; nn₂₄₆₀ - nn₂₄₇₀; nn₂₄₇₀ - nn₂₄₈₀;

nn2480 - nn2490; nn2490 - nn2500; nn2500 - nn2510;
nn2510 - nn2520; nn2520 - nn2530; nn2530 - nn2540;
nn2540 - nn2550; nn2550 - nn2560; nn2560 - nn2570;
nn2570 - nn2580; nn2580 - nn2590; nn2590 - nn2600;
5 nn2600 - nn2610; nn2610 - nn2620; nn2620 - nn2630;
nn2630 - nn2640; nn2640 - nn2650; nn2650 - nn2660;
nn2660 - nn2670; nn2670 - nn2680; nn2680 - nn2690;
nn2690 - nn2700; nn2700 - nn2710; nn2710 - nn2720;
nn2720 - nn2730; nn2730 - nn2740; nn2740 - nn2750;
10 nn2750 - nn2760; nn2760 - nn2770; nn2770 - nn2780;
nn2780 - nn2790; nn2790 - nn2800; nn2800 - nn2810;
nn2810 - nn2820; nn2820 - nn2830; nn2830 - nn2840;
nn2840 - nn2850; nn2850 - nn2860; nn2860 - nn2870;
nn2870 - nn2880; nn2880 - nn2890; nn2890 - nn2900;
15 nn2900 - nn2910; nn2910 - nn2920; nn2920 - nn2930;
nn2930 - nn2940; nn2940 - nn2950; nn2950 - nn2960;
nn2960 - nn2970; nn2970 - nn2980; nn2980 - nn2990;
nn2990 - nn3000; nn3000 - nn3010; nn3010 - nn3020;
nn3020 - nn3030; nn3030 - nn3040; nn3040 - nn3050;
20 nn3050 - nn3060; nn3060 - nn3070; nn3070 - nn3080;
nn3080 - nn3090; nn3090 - nn3100; nn3100 - nn3110;
nn3110 - nn3120; nn3120 - nn3130; nn3130 - nn3140;
nn3140 - nn3150; nn3150 - nn3160; nn3160 - nn3170;
nn3170 - nn3180; nn3180 - nn3190; nn3190 - nn3200;
25 nn3200 - nn3210; nn3210 - nn3220; nn3220 - nn3230;
nn3230 - nn3240; nn3240 - nn3250; nn3250 - nn3260;
nn3260 - nn3270; nn3270 - nn3280; nn3280 - nn3290;
nn3290 - nn3300; nn3300 - nn3310; nn3310 - nn3320;
nn3320 - nn3330; nn3330 - nn3340; nn3340 - nn3350;
30 nn3350 - nn3360; nn3360 - nn3370; nn3370 - nn3380;
nn3380 - nn3390; nn3390 - nn3400; nn3400 - nn3410;
nn3410 - nn3420; nn3420 - nn3430; nn3430 - nn3440;
nn3440 - nn3450; nn3450 - nn3460; nn3460 - nn3470;
nn3470 - nn3480; nn3480 - nn3490; nn3490 - nn3500;
35 nn3500 - nn3510; nn3510 - nn3520; nn3520 - nn3530;
nn3530 - nn3540; nn3540 - nn3550; nn3550 - nn3560;

nn3560 - nn3570; nn3570 - nn3580; nn3580 - nn3590;
nn3590 - nn3600; nn3600 - nn3610; nn3610 - nn3620;
nn3620 - nn3630; nn3630 - nn3640; nn3640 - nn3650;
nn3650 - nn3660; nn3660 - nn3670; nn3670 - nn3680;
5 nn3680 - nn3690; nn3690 - nn3700; nn3700 - nn3710;
nn3710 - nn3720; nn3720 - nn3730; nn3730 - nn3740;
nn3740 - nn3750; nn3750 - nn3760; nn3760 - nn3770;
nn3770 - nn3780; nn3780 - nn3790; nn3790 - nn3800;
nn3800 - nn3810; nn3810 - nn3820; nn3820 - nn3830;
10 nn3830 - nn3840; nn3840 - nn3850; nn3850 - nn3860;
nn3860 - nn3870; nn3870 - nn3880; nn3880 - nn3890;
nn3890 - nn3900; nn3900 - nn3910; nn3910 - nn3920;
nn3920 - nn3930; nn3930 - nn3940; nn3940 - nn3950;
nn3950 - nn3960; nn3960 - nn3970; nn3970 - nn3980;
15 nn3980 - nn3990; nn3990 - nn4000; nn4000 - nn4010;
nn4010 - nn4020; nn4020 - nn4030; nn4030 - nn4040;
nn4040 - nn4050; nn4050 - nn4060; nn4060 - nn4070;
nn4070 - nn4080; nn4080 - nn4090; nn4090 - nn4100;
nn4100 - nn4110; nn4110 - nn4120; nn4120 - nn4130;
20 nn4130 - nn4140; nn4140 - nn4150; nn4150 - nn4160;
nn4160 - nn4170; nn4170 - nn4180; nn4180 - nn4190;
nn4190 - nn4200; nn4200 - nn4210; nn4210 - nn4220;
nn4220 - nn4230; nn4230 - nn4240; nn4240 - nn4250;
nn4250 - nn4260; nn4260 - nn4270; nn4270 - nn4280;
25 nn4280 - nn4290; nn4290 - nn4300; nn4300 - nn4310;
nn4310 - nn4320; nn4320 - nn4330; nn4330 - nn4340;
nn4340 - nn4350; nn4350 - nn4360; nn4360 - nn4370;
nn4370 - nn4380; nn4380 - nn4390; nn4390 - nn4400;
nn4400 - nn4410; nn4410 - nn4420; nn4420 - nn4430;
30 nn4430 - nn4440; nn4440 - nn4450; nn4450 - nn4460;
nn4460 - nn4470; nn4470 - nn4480; nn4480 - nn4490;
nn4490 - nn4500; nn4500 - nn4510; nn4510 - nn4520;
nn4520 - nn4530; nn4530 - nn4540; nn4540 - nn4550;
nn4550 - nn4560; nn4560 - nn4570; nn4570 - nn4580;
35 nn4580 - nn4590; nn4590 - nn4600; nn4600 - nn4610;
nn4610 - nn4620; nn4620 - nn4630; nn4630 - nn4640;

nn4640 - nn4650; nn4650 - nn4660; nn4660 - nn4670;
nn4670 - nn4680; nn4680 - nn4690; nn4690 - nn4700;
nn4700 - nn4710; nn4710 - nn4720; nn4720 - nn4730;
nn4730 - nn4740; nn4740 - nn4750; nn4750 - nn4760;
5 nn4760 - nn4770; nn4770 - nn4780; nn4780 - nn4790;
nn4790 - nn4800; nn4800 - nn4810; nn4810 - nn4820;
nn4820 - nn4830; nn4830 - nn4840; nn4840 - nn4850;
nn4850 - nn4860; nn4860 - nn4870; nn4870 - nn4880;
nn4880 - nn4890; nn4890 - nn4900; nn4900 - nn4910;
10 nn4910 - nn4920; nn4920 - nn4930; nn4930 - nn4940;
nn4940 - nn4950; nn4950 - nn4960; nn4960 - nn4970;
nn4970 - nn4980; nn4980 - nn4990; nn4990 - nn5000;
nn5000 - nn5010; nn5010 - nn5020; nn5020 - nn5030;
nn5030 - nn5040; nn5040 - nn5050; nn5050 - nn5060;
15 nn5060 - nn5070; nn5070 - nn5080; nn5080 - nn5090;
nn5090 - nn5100; nn5100 - nn5110; nn5110 - nn5120;
nn5120 - nn5130; nn5130 - nn5140; nn5140 - nn5150;
nn5150 - nn5160; nn5160 - nn5170; nn5170 - nn5180;
nn5180 - nn5190; nn5190 - nn5200; nn5200 - nn5210;
20 nn5210 - nn5220; nn5220 - nn5230; nn5230 - nn5240;
nn5240 - nn5250; nn5250 - nn5260; nn5260 - nn5270;
nn5270 - nn5280; nn5280 - nn5290; nn5290 - nn5300;
nn5300 - nn5310; nn5310 - nn5320; nn5320 - nn5330;
nn5330 - nn5340; nn5340 - nn5350; nn5350 - nn5360;
25 nn5360 - nn5370; nn5370 - nn5380; nn5380 - nn5390;
nn5390 - nn5400; nn5400 - nn5410; nn5410 - nn5420;
nn5420 - nn5430; nn5430 - nn5440; nn5440 - nn5450;
nn5450 - nn5460; nn5460 - nn5470; nn5470 - nn5480;
nn5480 - nn5490; nn5490 - nn5500; nn5500 - nn5510;
30 nn5510 - nn5520; nn5520 - nn5530; nn5530 - nn5540;
nn5540 - nn5550; nn5550 - nn5560; nn5560 - nn5570;
nn5570 - nn5580; nn5580 - nn5590; nn5590 - nn5600;
nn5600 - nn5610; nn5610 - nn5620; nn5620 - nn5630;
nn5630 - nn5640; nn5640 - nn5650; nn5650 - nn5660;
35 nn5660 - nn5670; nn5670 - nn5680; nn5680 - nn5690;
nn5690 - nn5700; nn5700 - nn5710; nn5710 - nn5720;

nn5720 - nn5730; nn5730 - nn5740; nn5740 - nn5750;
nn5750 - nn5760; nn5760 - nn5770; nn5770 - nn5780;
nn5780 - nn5790; nn5790 - nn5800; nn5800 - nn5810;
nn5810 - nn5820; nn5820 - nn5830; nn5830 - nn5840;
5 nn5840 - nn5850; nn5850 - nn5860; nn5860 - nn5870;
nn5870 - nn5880; nn5880 - nn5890; nn5890 - nn5900;
nn5900 - nn5910; nn5910 - nn5920; nn5920 - nn5930;
nn5930 - nn5940; nn5940 - nn5950; nn5950 - nn5960;
nn5960 - nn5970; nn5970 - nn5980; nn5980 - nn5990;
10 nn5990 - nn6000; nn6000 - nn6010; nn6010 - nn6020;
nn6020 - nn6030; nn6030 - nn6040; nn6040 - nn6050;
nn6050 - nn6060; nn6060 - nn6070; nn6070 - nn6080;
nn6080 - nn6090; nn6090 - nn6100; nn6100 - nn6110;
nn6110 - nn6120; nn6120 - nn6130; nn6130 - nn6140;
15 nn6140 - nn6150; nn6150 - nn6160; nn6160 - nn6170;
nn6170 - nn6180; nn6180 - nn6190; nn6190 - nn6200;
nn6200 - nn6210; nn6210 - nn6220; nn6220 - nn6230;
nn6230 - nn6240; nn6240 - nn6250; nn6250 - nn6260;
nn6260 - nn6270; nn6270 - nn6280; nn6280 - nn6290;
20 nn6290 - nn6300; nn6300 - nn6310; nn6310 - nn6320;
nn6320 - nn6330; nn6330 - nn6340; nn6340 - nn6350;
nn6350 - nn6360; nn6360 - nn6370; nn6370 - nn6380;
nn6380 - nn6390; nn6390 - nn6400; nn6400 - nn6410;
nn6410 - nn6420; nn6420 - nn6430; nn6430 - nn6440;
25 nn6440 - nn6450; nn6450 - nn6460; nn6460 - nn6470;
nn6470 - nn6480; nn6480 - nn6490; nn6490 - nn6500;
nn6500 - nn6510; nn6510 - nn6520; nn6520 - nn6530;
nn6530 - nn6540; nn6540 - nn6550; nn6550 - nn6560;
nn6560 - nn6570; nn6570 - nn6580; nn6580 - nn6590;
30 nn6590 - nn6600; nn6600 - nn6610; nn6610 - nn6620;
nn6620 - nn6630; nn6630 - nn6640; nn6640 - nn6650;
nn6650 - nn6660; nn6660 - nn6670; nn6670 - nn6680;
nn6680 - nn6690; nn6690 - nn6700; nn6700 - nn6710;
nn6710 - nn6720; nn6720 - nn6730; nn6730 - nn6740;
35 nn6740 - nn6750; nn6750 - nn6760; nn6760 - nn6770;
nn6770 - nn6780; nn6780 - nn6790; nn6790 - nn6800;

nn6800 - nn6810; nn6810 - nn6820; nn6820 - nn6830;
nn6830 - nn6840; nn6840 - nn6850; nn6850 - nn6860;
nn6860 - nn6870; nn6870 - nn6880; nn6880 - nn6890;
nn6890 - nn6900; nn6900 - nn6910; nn6910 - nn6920;
5 nn6920 - nn6930; nn6930 - nn6940; nn6940 - nn6950;
nn6950 - nn6960; nn6960 - nn6970; nn6970 - nn6980;
nn6980 - nn6990; nn6990 - nn7000; nn7000 - nn7010;
nn7010 - nn7020; nn7020 - nn7030; nn7030 - nn7040;
nn7040 - nn7050; nn7050 - nn7060; nn7060 - nn7070;
10 nn7070 - nn7080; nn7080 - nn7090; nn7090 - nn7100;
nn7100 - nn7110; nn7110 - nn7120; nn7120 - nn7130;
nn7130 - nn7140; nn7140 - nn7150; nn7150 - nn7160;
nn7160 - nn7170; nn7170 - nn7180; nn7180 - nn7190;
nn7190 - nn7200; nn7200 - nn7210; nn7210 - nn7220;
15 nn7220 - nn7230; nn7230 - nn7240; nn7240 - nn7250;
nn7250 - nn7260; nn7260 - nn7270; nn7270 - nn7280;
nn7280 - nn7290; nn7290 - nn7300; nn7300 - nn7310;
nn7310 - nn7320; nn7320 - nn7330; nn7330 - nn7340;
nn7340 - nn7350; nn7350 - nn7360; nn7360 - nn7370;
20 nn7370 - nn7380; nn7380 - nn7390; nn7390 - nn7400;
nn7400 - nn7410; nn7410 - nn7420; nn7420 - nn7430;
nn7430 - nn7440; nn7440 - nn7450; nn7450 - nn7460;
nn7460 - nn7470; nn7470 - nn7480; nn7480 - nn7490;
nn7490 - nn7500; nn7500 - nn7510; nn7510 - nn7520;
25 nn7520 - nn7530; nn7530 - nn7540; nn7540 - nn7550;
nn7550 - nn7560; nn7560 - nn7570; nn7570 - nn7580;
nn7580 - nn7590; nn7590 - nn7600; nn7600 - nn7610;
nn7610 - nn7620; nn7620 - nn7630; nn7630 - nn7640;
nn7640 - nn7650; nn7650 - nn7660; nn7660 - nn7670;
30 nn7670 - nn7680; nn7680 - nn7690; nn7690 - nn7700;
nn7700 - nn7710; nn7710 - nn7720; nn7720 - nn7730;
nn7730 - nn7740; nn7740 - nn7750; nn7750 - nn7760;
nn7760 - nn7770; nn7770 - nn7780; nn7780 - nn7790;
nn7790 - nn7800; nn7800 - nn7810; nn7810 - nn7820;
35 nn7820 - nn7830; nn7830 - nn7840; nn7840 - nn7850;
nn7850 - nn7860; nn7860 - nn7870; nn7870 - nn7880;

nn7880 - nn7890; nn7890 - nn7900; nn7900 - nn7910;
nn7910 - nn7920; nn7920 - nn7930; nn7930 - nn7940;
nn7940 - nn7950; nn7950 - nn7960; nn7960 - nn7970;
nn7970 - nn7980; nn7980 - nn7990; nn7990 - nn8000;
5 nn8000 - nn8010; nn8010 - nn8020; nn8020 - nn8030;
nn8030 - nn8040; nn8040 - nn8050; nn8050 - nn8060;
nn8060 - nn8070; nn8070 - nn8080; nn8080 - nn8090;
nn8090 - nn8100; nn8100 - nn8110; nn8110 - nn8120;
nn8120 - nn8130; nn8130 - nn8140; nn8140 - nn8150;
10 nn8150 - nn8160; nn8160 - nn8170; nn8170 - nn8180;
nn8180 - nn8190; nn8190 - nn8200; nn8200 - nn8210;
nn8210 - nn8220; nn8220 - nn8230; nn8230 - nn8240;
nn8240 - nn8250; nn8250 - nn8260; nn8260 - nn8270;
nn8270 - nn8280; nn8280 - nn8290; nn8290 - nn8300;
15 nn8300 - nn8310; nn8310 - nn8320; nn8320 - nn8330;
nn8330 - nn8340; nn8340 - nn8350; nn8350 - nn8360;
nn8360 - nn8370; nn8370 - nn8380; nn8380 - nn8390;
nn8390 - nn8400; nn8400 - nn8410; nn8410 - nn8420;
nn8420 - nn8430; nn8430 - nn8440; nn8440 - nn8450;
20 nn8450 - nn8460; nn8460 - nn8470; nn8470 - nn8480;
nn8480 - nn8490; nn8490 - nn8500; nn8500 - nn8510;
nn8510 - nn8520; nn8520 - nn8530; nn8530 - nn8540;
nn8540 - nn8550; nn8550 - nn8560; nn8560 - nn8570;
nn8570 - nn8580; nn8580 - nn8590; nn8590 - nn8600;
25 nn8600 - nn8610; nn8610 - nn8620; nn8620 - nn8630;
nn8630 - nn8640; nn8640 - nn8650; nn8650 - nn8660;
nn8660 - nn8670; nn8670 - nn8680; nn8680 - nn8690;
nn8690 - nn8700; nn8700 - nn8710; nn8710 - nn8720;
nn8720 - nn8730; nn8730 - nn8740; nn8740 - nn8750;
30 nn8750 - nn8760; nn8760 - nn8770; nn8770 - nn8780;
nn8780 - nn8790; nn8790 - nn8800; nn8800 - nn8810;
nn8810 - nn8820; nn8820 - nn8830; nn8830 - nn8840;
nn8840 - nn8850; nn8850 - nn8860; nn8860 - nn8870;
nn8870 - nn8880; nn8880 - nn8890; nn8890 - nn8900;
35 nn8900 - nn8910; nn8910 - nn8920; nn8920 - nn8930;
nn8930 - nn8940; nn8940 - nn8950; nn8950 - nn8960;

nn₈₉₆₀ - nn₈₉₇₀; nn₈₉₇₀ - nn₈₉₈₀; nn₈₉₈₀ - nn₈₉₉₀;
nn₈₉₉₀ - nn₉₀₀₀; nn₉₀₀₀ - nn₉₀₁₀; nn₉₀₁₀ - nn₉₀₂₀;
nn₉₀₂₀ - nn₉₀₃₀; nn₉₀₃₀ - nn₉₀₄₀; nn₉₀₄₀ - nn₉₀₅₀;
nn₉₀₅₀ - nn₉₀₆₀.

5

3. The oligomer of claim 1, wherein the targeting sequence is comprised of a sequence which is complementary to a sequence of at least 8 nucleotides present in a conserved HCV nucleotide sequence in HCV RNA.

10

4. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from the 5'-terminus to about 200 in Fig. 18.

15

5. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 4000 to about 5000 in Fig. 18.

20

6. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 8000 to about 9040 as shown in Fig. 18.

25

7. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about -318 to about 174 as shown in Fig. 18.

30

8. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about or from about 4056 to about 4448 as shown in Fig. 18.

35

9. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 4378 to about 4902 as shown in Fig. 18.

5

10. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 4042 to about 4059 as shown in Fig. 18.

10

11. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 4456 to about 4470, as shown in Fig. 18.

15

12. The oligomer of claim 3, wherein the conserved sequence is located in the sequence of nucleotide numbers from about 8209 to about 8217, as shown in Fig. 18.

20

13. The oligomer of claim 3, which is a capture probe.

14. The oligomer of claim 3, which is a label
25 probe.

15. The oligomer of claim 3, which is a primer.

30 16. A process for detecting an HCV sequence in an analyte strand suspected of containing an HCV polynucleotide, wherein the HCV polynucleotide comprises a selected target region, said process comprising:

35 (a) providing an oligomer capable of hybridizing to an HCV sequence in an analyte polynucleotide strand, wherein the oligomer is comprised of an HCV targeting

sequence complementary to at least 4 contiguous nucleotides of HCV cDNA shown in Fig. 18

(b) incubating the analyte strand with the oligomer of (a) which allow specific hybrid duplexes to
5 form between the targeting sequence and the target sequence; and

(d) detecting hybrids formed between target region, if any, and the oligomer.

10

17. The process of claim 16 which further comprises:

(a) providing a set oligomers which are primers for the polymerase chain reaction method and which flank
15 the target region; and

(b) amplifying the target region via a polymerase chain reaction method.

18. A kit for detecting an HCV target sequence
20 in an analyte strand, comprising the oligomer of claim 1 packaged in a suitable container.

19. A method for preparing blood free of HCV comprising:

25 (a) providing analyte nucleic acids from a sample of blood suspected of containing an HCV target sequence;

(b) providing an oligomer capable of hybridizing to the HCV sequence in an analyte polynucleotide
30 strand, if any, wherein the oligomer is comprised of an HCV targeting sequence complementary to a sequence of at least 8 nucleotides present in a conserved HCV nucleotide sequence in HCV RNA;

(c) reacting (a) with (b) under conditions
35 which allow the formation of a polynucleotide duplex

between the targeting sequence and the target sequence, if any;

(d) detecting a duplex formed in (c), if any;

and

5 (e) saving the blood from which complexes were not detected in (d).

10

15

20

25

30

35

1 / 86

FIG. 1 Translation of DNA 12f

IlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeuGluAlaCysAsn
 1 CCATATTAAATCAGGATGTACGTGGAGGGTCCGAACACAGGCTGGAGCTGCCCTGCA
 GGATAAATTTTAGTCCATACATGCACCTCCCGAGCTTGTTGTCCGACCTTCGACGGACGT
 TrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSerGluLeuSerProLeu
 61 ACTGGACGGGGCGGAACGTTGCGATCTGGAAGACAGGACAGGTCCGAGCTCAGCCCCGT
 TGACCTGGCCCCCGCTTGCAACGCTAGACCTTCTGTCCCTGTCCAGGCTCGAGTCGGGCA
 LeuLeuThrThrThrGlnTrpGlnValLeuProCysSerPheThrThrLeuProAlaLeu
 121 TACTGCTGACCACTACACAGTGGCAGTCCCTCCCGTGTTCCTTCACACCTACAGCCT
 ATGACGACTGCTGATGTGTCAACCTCCAGGAGGACAAAGGAGTGTGGATGGTCGGA
 SerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGlnTyrLeuTyrGlyVal
 181 TGTCCACCGGCTCATCCACCTCCACCAGAACATTTGGACGTGCAGTACTTGTACGGGG
 ACAGGTGGCCGGAGTAGGTGGAGGTGCTTTGTAACACCTGCACGTCATGAACATGCCCC

 GlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValValLeuLeuPheLeuLeu
 241 TGGGGTCAAGCATCGCGTCCCTGGGCCATTAAAGTGGGAGTACGTCGTTCTCCTGTTCCCTTC
 ACCCCAGTTCGTAGCGCAGGACCCCGGTAAATTCACCTCATGCAGCAAGAGACAAGGAAG

 LeuAlaAspAlaArgValCysSerCysLeuTrpMetMetLeuLeuIleSerGlnAlaGlu
 301 TGCTTGCAGACGCGCGCTGCTCCTGCTTGTTGGATGATGCTACTCATATCCCCAAGCGG
 ACGAACGTCTGCGCGCGCAGACGAGGACGAACACCTACTACGATGAGTATAGGGTTTCGCC
 -----Overlap with 14i-----
 AlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAlaGlyThrHisGlyLeu
 361 AGCGCGCTTTGGAGAACCTCGTAATACTTAATGCAGCATCCCTGGCCGGACGACGGTC
 TCCGCCGAAACCTCTTGGAGCATTAATGAATTACGTCGTAGGACCGGCCCTGCGGTGCCAG

 Val
 421 TTGTATC
 AACATAG

SUBSTITUTE SHEET

2 / 86

FIG. 2-1 Translation of DNA k9-1

GlyCysProGluArgLeuAlaSerCysArgProLeuThrAspPheAspGlnGlyTrpGly
 1 CAGGCTGTCTGAGAGGCTAGCCAGCTGCCGACCCCTTACCGATTTTGACCAAGGCTGGG
 GTCCGACAGGACTCTCCGATCCGATCGGCTCGACGGCTGGGAATGGCTAAACTGGTCCCGACCC

ProIleSerTyrAlaAsnGlySerGlyProAspGlnArgProTyrCysTrpHisTyrPro
 61 GCCCTATCAGTTATGCCAACGGAAGCGGCCCGACACGACGCCCCCTACTGCTGGCACTACC
 CGGGATAGTCAATACGGTTGCCCTTCGCCGGGCTGGTCGCGGGATGACGACCGTGATGG

ProLysProCysGlyIleValProAlaLysSerValCysGlyProValTyrCysPheThr
 121 CCCCAAACCTTGCGGTATTGTGCCCGAAGAGTGTGTGGTCCGGTATATTGCTTCA
 GGGGTTTTGGAACGCCATAACACGGGGCTTCTCACACACACACGAGGCCATATAACGAAGT

ProSerProValValGlyThrThrAspArgSerGlyAlaProThrTyrSerTrpGly
 181 CTCCCAGCCCCGTGGTGGGAACGACCGACAGGTCCGGCGCGCCACCTACAGCTGGG
 GAGGGTCGGGGCACCAACCCCTTGCTGGCTGTCCAGCCCGCGCGGTGGATGTCGACCC

GluAsnAspThrAspValPheValLeuAsnAsnThrArgProProLeuGlyAsnTrpPhe
 241 GTGAAAATGATACGGACGTCTTCGTCCTTAACAATACCAAGCCACCGCTGGGCAATTGGT
 CACTTTTACTATGCCCTGCAGAACGAGGAATTGTTATGGTCCGGTGGCGACCCGTTAACCA

GlyCysThrTrpMetAsnSerThrGlyPheThrLysValCysGlyAlaProProCysVal
 301 TCGGTGTACCTGGATGAACCTCAACTGGATTCAACAAAGTGTGCGGAGCGCCTCCTTG
 AGCCCAACATGGACCTACTTGAGTTGACCTAAGTGGTTTACACACGCTTCGCGGAGGAACAC

SUBSTITUTE SHEET

3/86

FIG. 2-2

IleGlyGlyAlaGlyAsnAsnThrLeuHisCysProThrAspCysPheArgLysHisPro
361 TCATCGAGGGCGGCAACAACACCTGCACCTGCCACTGATTGCTTCCGCAAGCATC
AGTAGCCTCCCCCGCGTGTGTGTGGACGTGACGGGTGACTAACGAAGCGTTCGTAG

AspAlaThrTyrSerArgCysGlySerGlyProTrpIleThrProArgCysLeuValAsp
421 CGGACGCCACATACTCTGGTGGCTCCGGTCCCTGGATCACACCCAGGTGCCGTGTCG
GCCTGCGGTGTATGAGAGCCACGCCGAGGCCAGGACCTAGTGTGGTCCACGGACCAGC

TyrProTyrArgLeuTrpHisTyrProCysThrIleAsnTyrThrIlePheLysIleArg
481 ACTACCCGTATAGGCTTTGGCATTAATCCTGTACCATCAACTACATATATTAAATCA
TGATGGCCATATCCGAAACCGTAATAGGAACATGGTAGTTGATGTGATATAAATTTTAGT

MetTyrValGlyGlyValGluHisArgLeuGluAlaAlaCysAsnTrpThrArgGlyGlu
541 GGATGTACGTGGGAGGGTCGAGCACAGGCTGGAAGCTGCCCTGCAACTGGACGCGGGCG
CCTACATGCACCCCTCCCCAGCTCGTGTCCGACCTTCGACGGACGTTGACCTGCGCCCCCGC

ArgCysAspLeuGluAspArgAspArgSerGluLeuSerProLeuLeuThrThrThr
601 AACGTTGCGATCTGGAAGATAGGACAGGTCCGAGCTCAGCCCGTTACTGTGACCACTA
TTGCAACGCTAGACCTTCTATCCCTGTCCAGGCTCGAGTCGGGCAATGACGACTGGTGAT

GlnTrpGlnValLeuProCysSerPheThrThrLeuProAlaLeuSerThrGlyLeuIle
661 CACAGTGGCAGGTCTCCGTTCTCCTTCAACAACCTGCCAGCCTTGTCACCGCCCTCA
GTGTACCGTCCAGGAGGCCACAAAGGAAGTGTGGGACGTCGGAAACAGGTGGCGCGGAGT

SUBSTITUTE SHEET

5 / 86

FIG. 2-4

TyrGlyMetTrpProLeuLeuLeuLeuAlaLeuProGlnArgAlaTyrAlaLeu
1021 TCTACGGGATGTGGCCTCTCCTCCTGCTCCTGTGGCGTTGCCCGAGCGGGCTACGCGC
AGATGCCCTACACCGGAGAGGAGGACGAGACAACCGCAACGGGGTCGCCCCGATGCGCG

AspThrGluValAlaAlaSerCysGlyGlyValValLeuValGlyLeuMetAlaLeuThr
1081 TGGACACGGAGGTGGCCGCTCGTGGCGGTGTGTCTCGTCGGGTTGATGGCGCTAA
ACCTGTGCCTCCACCGCGCAGCACACCGCCACAACAAGAGCAGCCCAACTACCGCGATT

LeuSerProTyrTyrLysArgTyrIleSerTrpCysLeuTrpTrpLeuGlnTyrPheLeu
1141 CTCTGTCACCATATTACAAGCGCTATATCAGCTGGTGTGTGGTTCAGTATTTC
GAGACAGTGGTATAATGTTCGCGATATAGTCGACCACCAACACCAGTCATAAAAG

ThrArgValGluAlaGlnLeuHisValTrpIleProProLeuAsnValArgGlyGlyArg
1201 TGACCAGAGTGGAAAGCGCAACTGCACGTGTGGATTCCCCCTCAACGTCCGAGGGGGC
ACTGGTCTCACCTTCGCGTTGACGTGCACACCTAAGGGGGGAGTTGCAGGCTCCCCCG

AspAlaValIleLeuLeuMetCysAlaValHisProThrLeuValPheAspIleThrLys
1261 GCGACGCTGTCACTTACTCATGTGTGTGTACACCCGACTCTGTATTGACATCACCA
CGCTGCGACAGTAGAATGAGTACACACGACATGTGGCTGAGACCATAAACTGTAGTGGT

LeuLeuLeuAlaValPheGlyProLeuTrpIleLeuGlnAla
1321 AATTGCTGCTGGCCGTCTTCGGACCCCTTTGGATTCTTCAAGCCAG
TTAACGACGACCGGCAGAGCCTGGGGAAACCTAAGAGTTCGGTC

SUBSTITUTE SHEET

FIG. 3 Translation of DNA 15e

GlyAlaGlyLysArgValTyrLeuThrArgAspProThrProLeuAlaArgAla
CGCGCTGGAAGAGGTCTACTACCTCACCCGTGACCTACAACCCCTCGGAGAGC
GCCGCGACCTTCTCCAGATGATGGAGTGGCACTGGGATGTGGGGGGAGCGCTCTCG

Overlap with 26g-

AlaTrpGluThrAlaArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMetPhe
TGCGTGGAGACAGACACACTCCAGTCAATTCTTGCTAGGCAACATAATCATGTT
ACGCACCCCTCTGTCGTTCTGTGTGAGGTCAGTTAAGGACCGATCCGTTGTATTAGTACAA

AlaProThrLeuTrpAlaArgMetIleLeuMetThrHisPhePheSerValLeuIleAla
TGCCCCCACACTGTGGCGAGGATGATGATGACCCATTCTTAGCGTCCCTATAGC
ACGGGGTGTGACACCCGCTCTACTATGACTACTGGGTAAAGAAATCGCAGGAATATCG

ArgAspGlnLeuGluGlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIleGlu
CAGGGACCAGCTTGAAACAGGCCCTCGATTCCGAGATCTACGGGCCCTGCTACTCCATAGA
GTCCCTGGTCGAAC TTGTCCGGGAGCTAACGCTCTAGATGCCCCGGACGATGAGGTATCT

ProLeuAspLeuProProIleIleGlnArgLeu
ACCACTTGATCTACCTCCAATCATTCAAAGACTC
TGGTGAACTAGATGGAGGTTAGTAAGTTTCTGAG

7/86

FIG.4 Translation of DNA 131

```

ProSerProValValGlyThrThrAspArgSerGlyAlaProThrTyrSerTrpGly
1 CTCCAGCCCCGTGGTGGAAACGACCGACAGTCGGCGCGCTACCTACAGCTGGG
  GAGGTCGGGGCACCAACCCTTGCTGGCTGTCCAGCCCGCGGATGATGTCGACCC
  GluAsnAspThrAspValPheValLeuAsnAsnThrArgProProLeuGlyAsnTrpPhe
61 GTGAAATGATACGGACGTCTTCGTCCTTAACAATACCAAGCCACCGCTGGCAATTGGT
  CACTTTACTATGCTGCAGAACGAGGAATTGTATGGTCCGGTGGCACCCTTAACCA
  GlyCysThrTrpMetAsnSerThrGlyPheThrLysValCysGlyAlaProProCysVal
121 TCGGTTGTACCTGGATGAACCTCAACTGGATTCAACCAAGTGTGGAGCGCTCCTTGTG
  AGCCAACATGGACCTACTTGAGTTGACCTAAGTGTTCACACGCTCGCGGAGGAACAC
  IleGlyGlyAlaGlyAsnAsnThrLeuHisCysProThrAspCysPheArgLysHisPro
181 TCATCGGAGGGCGGCAACACACCTGCACTGCCCACTGATTGCTCCGCAAGCATC
  AGTAGCCTCCCCCGCTTGTGGGACGTGACGGGTGACTAACGAAGCGTTCGTAG
  AspAlaThrTyrSerArgCysGlySerGlyProTrpLeuThrProArgCysLeuValAsp
241 CGGACGCCACATACTCTCGGTGGCTCCGGTCCCTGGCTCACACCCAGGTGCTGTGTCG
  GCCTCGGGTATGAGAGCCACGCCGAGGCCAGGACCGAGTGTGGTCCACGGACCAGC
  -----
TyrProTyrArgLeuTrpHisTyrProCysThrIleAsnTyrThrIlePheLysIleArg
301 ACTACCCGTATAGGCTTTGGCATATATCTTGATCCATCAACTACCATATTTAAATCA
  TGATGGGCATATCCGAAACCGTAATAGGAACATGGTAGTGTGATATAAATTTAGT
  -----
MetTyrValGlyGlyValGluHisArgLeuGluAlaAlaCysAsnTrpThrArgGlyGlu
361 GGATGTACGTGGGAGGGTCGAGCACAGGCTGGAGCTGCCTGCAACTGGACGGGGCG
  CCTACATGCCACCTCCCGAGCTCGTGTCCGACCTTCGACGGACGTTGACCTGCGCCCCCG
  -----Overlap with 12f-----
ArgCysAspLeuGluAspArgAspArgSerGluLeuSerProLeuLeuThrThrThr
421 AACGTTGCGATCTGGAAGACAGGACAGGTCGAGCTCAGCCCGTTACTGCTGACCACTA
  TTGCAACGCTAGACCTTCTGTCCCTGTCAGGCTCGAGTCGGGCAATGACGACTGGTGAT
  -----
GlnTrpGlnValLeuProCysSerPheThrThrLeuProAlaLeuSerThrGlyLeu
481 CACAGTGGCAGGTCTCCCGTGTTCCTTCACAACCCCTGCCAGCCTTGTCACCGCCCTCA
  GTGTACCGTCCAGGAGGCACAAAGGAAGTGTGGGACGGTCGGAACAGGTGGCCGGAGT

```

SUBSTITUTE SHEET

8 / 86

Translation of DNA 26j FIG. 5

LeuPheTyrHisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArg
1 GCCTTTCTATCACCAAGTTCAACTCTTCAGGCTGTCCTGAGAGGCTAGCCAGCTGCCG
CGAAAAGATAGTGGTTCAGTTGAGAGTCCGACAGGACTCTCCGATCGGTCGACGGC

ProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyPro
61 ACCCCTTACCGATTGTGACCAAGGCTGGGGCCCTATCAGTTATGCCAACGGAAGCGCCC
TGGGGAATGGCTAAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCCTTCGCCGGG

AspGlnArgProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLys
121 CGACCAGCGCCCTACTGCTGGCACTACCCCCCAAACCTTGCGGTATTGTGCCCGCGAA
GCTGGTCGGGGGATGACGACCGTGTATGGGGGTTTGGAAACGCCATAACACGGGCGCTT

---Overlap with 13i---

SerValCysGlyProValTyrCysPheThrProSerProValValVal
181 GAGTGTGTGGTCCGGTATATTGCTTCACTCCAGCCCCCGTGGTGGTGGG
CTCACACACACAGGCCATATAACGAAGTGAGGTCGGGGCACCACCC

SUBSTITUTE SHEET

9 / 86

Translation of DNA CA59a FIG. 6

```

LeuValMetAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIleAlaGlyAla
1 TTGGTAATGGCTCAGCTGCTCCGGATCCCAAGCCATCTTGGACATGATCGTGGTGCT
AACCATTAACGAGTCGACGAGCCCTAGGGTGTTCGGTAGAACCTGTACTAGCGACCCACGA

HisTrpGlyValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrpAlaLysVal
61 CACTGGGGAGTCCCTGGCGGCATAGCGTATTTCTCCATGGTGGGAACTGGCGGAAGGTC
GTGACCCCTCAGGACCGCCCGTATCGCATAAAGAGGTACCAACCCCTTGACCCGCTTCCAG

LeuValValLeuLeuLeuPheAlaGlyValAspAlaGluThrHisValThrGlyGlySer
121 CTGGTAGTGTGCTGCTATTTCGCGCGTCGACGCGGAACCCACGTCACCGGGGAAGT
GACCATCACGACGACGATAAACGGCCGCGAGCTGGCCCTTTGGGTGCAGTGGCCCCCTTCA

AlaGlyHisThrValSerGlyPheValSerLeuLeuAlaProGlyAlaLysGlnAsnVal
181 GCCGGCCACACTGTGTCTGGATTGTGTAGCTCTCCGACCAAGCGGCCCAAGCAGAACGTC
CGCCCGGTGTGACACAGACCTAAACAATCGGAGGAGCGTGGTCCGCGGTTCGCTTGCAG

GlnLeuIleAsnThrAsnGlySerTrpHisLeuAsnSerThrAlaLeuAsnCysAsnAsp
241 CAGCTGATCAACACCAACGGCAGTTGGCACCTCAATAGCACGGCCCTGAACGCAATGAT
GTCGACTAGTTGTGGTTGCCGTCAACCGTGGAGTTATCGTGCCGGGACTTGACGTTACTA
-----

SerLeuAsnThrGlyTrpLeuAlaGlyLeuPheTyrHisHisLysPheAsnSerSerGly
301 AGCCCTCAACACCGGCTGGTTGGCAGGGCTTTTCTATCACCACAAGTTCAACTCTTCAGGC
TCGGAGTTGTGGCCGACCAACCGTCCCGAAAGATAGTGGTGTTCAGTTGAGAAGTCCG
-----Overlap with 26j-----

-----Overlap with K9-1-----
CysProGluArgLeuAlaSerCysArgPro
361 TGTCCCTGAGAGGCTAGCCAGCTGCCGACCCC
ACAGGACTCTCCGATCGGTCGACGGCTGGG
-----

```

SUBSTITUTE SHEET

10 / 86

FIG. 7 Translation of DNA CA84a

GlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrpAsp
 1CGCAAGGTTGCAATTGCTCTATATCCCGGCCATATAACGGGTACCCGATGGCATGGG
 GCGTTCCCAACGTTAACGAGATAGAGCGCGGTATATTGCCCAGTGGCGTACCGTACCC

MetMetMetAsnTrpSerProThrAlaLeuValMetAlaGlnLeuLeuArgIlePro
 61ATATGATGATGAACCTGGTCCCTACGACGGCGTTGGTAATGGCTCAGCTCCGGATCC
 TATACTACTACTTGACCAAGGATGCTGCCGCAACCATTACCGAGTCGACGAGGCCTAGG

GlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyr
 121CACAAAGCCATCTTGGACATGATCGCTGGTGCTCACTGGGAGTCTCGGGCATAGCGT
 GTGTTCCGTAGAACCTGTACTAGCGACCACGAGTGACCCCTCAGGACCCCGTATCGCA

-----Overlap with CA59a-----
 PheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGlyVal
 181ATTCTCCATGGTGGGAAGTGGGCAAGTCTCGGTAGTGTGCTGCTATTGCCGGCG
 TAAAGAGTACCACCCCTTGACCCGCTTCCAGGACCATCAGACGACGATAAACGGCCCG

 AspAlaGluThrHisValThrGly
 241TCGACGCGGAACCCACGTACCCGGG
 AGCTGCGCCCTTGGGTGCAGTGGCCCC

SUBSTITUTE SHEET

11 / 86

FIG. 8 Translation of DNA CA156e

CysTrpValAlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThrGln
 1 GTGTGGTGGCGATGACCCCTACGGTGCCACCAAGGATGGCAAACTCCCGCGACGCA
 CACAACCCACCGCTACTGGGATGCCACCGTGGTCCCTACCGTTTGAGGGGCGCTGCGT

LeuArgArgHisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyrVal
 61 GCTTCGACGTCACATCGATCTGCTTGTCGGAGCGCCACCTCTGTTCGGCCCTCTACGT
 CGAAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGGGAGATGCA

GlyAspLeuCysGlySerValPheLeuValGlyGlnLeupheThrPheSerProArgArg
 121 GGGGACCTATGCGGCTCTGCTTCTTGTCGGCCAACCTGTTCACCTTCTCTCCAGGCG
 CCCCCTGGATACGCCCCAGACAGAAAGAACAGCCGTTGACAAGTGGAGAGAGGGTCCGC

 HisTrpThrThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArg
 181 CCACTGGACGACGCAAGTTGCAATTGCTCTATCTATCCCGCCATATAACGGTCAACCG
 GTGACCTGCTGCGTTCCAACGTTAACGAGATAGATAGGCCCGGTATATTGCCCCAGTGCG

-----Overlap with CA84a-----
 MetAlaTrpAspMetMetMetAsnTrpSerProThrThrAlaLeuValValAlaGlnLeu
 241 CATGGCATGGGATATGATGATGAACCTGGTCCCCCTACGACGGCGTTGGTAGTGCTCAGCT
 GTACCGTACCCCTATACTACTACTTGACCAAGGGGATGCTGCCGCAACCATCACCAGTCCA

 LeuArgIleProGlnAla
 301 GCTCCGGATCCCAAGCC
 CGAGGCCTAGGGTGTTCGG

SUBSTITUTE SHEET

12 / 86

FIG. 9 Translation of DNA CA167b

SerThrGlyLeuTyrHisValThrAsnAspCysProAsnSerSerIleValTyrGluAla
1 CTCCACGGGGCTTTACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGC
GAGGTGCCCCGGAATGGTGCAGTGGTTACTAACGGGATTGAGCTCATACACATGCTCCG

AlaAspAlaIleLeuHisThrProGlyCysValProCysValArgGluGlyAsnAlaSer
61 GGCCGATGCCATCCTGCACACTCCGGGTGCGTCCCTTGCGTTCGTGAGGGCAACGCCCTC
CCGGCTACGGTAGGACGTGTGAGGCCCCACGCAGGAAACGCAAGCACTCCCGTTGCGGAG

ArgCysTrpValAlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThr
121 GAGGTGTTGGTGGCGATGACCCCTACGGTGGCCACCAGGATGGCAAACTCCCCCGGAC
CTCCACAACCCACCGCTACTGGGGATGCCACCGGTGGTCCCTACCGTTTGAGGGGCGCTG

-----Overlap with CA156-----

GlnLeuArgArgHisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyr
181 GCAGCTTCGACGTCACATCGATCTGCTTGTCTGGGAGCGCTACCCCTCTGTTCGGCCCTCTA
CGTCGAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGATGGGAGACAAGCCGGGAGAT

ValGlyAspLeuCysGlySerValPheLeu
241 CGTGGGGGACTTGTGCGGGTCTGTCTTTCTTG
GCACCCCTGAACACGCCACAGACAGAAAGAAC

FIG.10 Translation of DNA ssCA216a

1 ArgArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCysGlyPheAlaAsp
 CCCGGCGTAGGTCGCGCAATTGGTAAGTCATCGATACCCCTTACGTGCGCTTCGCCG
 GGGCCGATCCAGCGCGTTAAACCCATTCCAGTAGCTATGGGAATGCACGCCGAAGCGGC
 LeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAla
 61 ACCTCATGGGGTACATACCGCTCGTCGGCGCCCTCTTGAGGCGCTGCCAGGCCCTGG
 TGGAGTACCCCATGTATGGCGAGCAGCCGGGGAGAACCTCCGCGACGGTCCCGGACC
 HisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsnLeuProGlyCys
 121 CGCATGGCGTCCGGTTCTGGGAAGACGGCGTGAACTATGCAACAGGGAACCTTCCTGGTT
 GCGTACCGCAGGCCCAAGACCTTCTGCCGCACTTGATACGTTGTCTCCCTTGAAGACCAA
 SerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValProAlaSerAlaTyr
 181 GCTCTTCTCTATCTTCTCTGCCCCCTGCTCTCTTGCTTGACTGTGCCCGCTTCGGCCT
 CGAGAAAGAGATAGAAGGAAGACCGGGACGAGAGAACGAACTGACACGGGCGAAGCCGGA

 GlnValArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysProAsnSerSerIle
 241 ACCAAGTGCAGCAACTCCACGGGGCTTTACCAACGTCAACCAATGATTGCCCTAACTCGAGTA
 TGGTTACGCGTTGAGGTGCCCCCGAAATGGTGTCAGTGGTTACTAAACGGGATTGAGCTCAT
 -----overlap with CA167b-----
 ValTyrGluAlaAlaAspAlaIleLeuHisThrProGlyCysValProCysValArgGlu
 301 TTGTGTACGAAGCGCCGATGCCATCCTGCACACTCCGGGTGCGTCCCTTGCGTTCTGTG
 AACACATGCTTCGCCGGCTACGGTAGGACGTGTGAGGCCCCACGACGAGGAACGCAAGCAC

 GlyAsnAlaSerArgCysTrpValAlaMetThrProThrValAla
 361 AGGGCAACGCTCGAGGTGTTGGTGGCGATGACCCCTACGGTGGCC
 TCCCGTTGCGGAGCTCCACAACCCACCGCTACTGGGGATGCCACCGG

FIG. 11

Translation of DNA ssCA290a

1 LysLysAsnLysArgAsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGly
 AAAAAAAAAACAAACGTAACACCAACCGTCGCCACAGGACGTCAAGTTCCCGGTGGCG
 TTTTTTTTTTTGTGTCATTGTGTTGGCAGCGGTCTCTGCAGTTCAAGGGCCACCGC
 GlnIleValGlyGlyValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAla
 61 GTCAGATCGTTGGTGGAGTTTACTTGTTCGCCGCGCAGGGCCCTAGATTGGGTGTGGCG
 CAGTCTAGCAACACCTCAATGAACAACGCGCTCCCGGATCTAACCACACGCGC
 ThrArgLysThrSerGluArgSerGlnProArgGlyArgGlnProIleProLysAla
 121 CGACGAGAAAGACTTCGAGCGGTCCGAACCTCGAGGTAGACGCCAGCCTATCCCCAAGG
 GCTGCTCTTCTGAAGGCTCGCCAGCGTTGGAGCTCCATCTGCGTCCGATAGGGTTCC
 ArgArgProGluGlyArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsn
 181 CTCGTGCGCCGAGGCGAGGACCTGGGCTCAGCCCGGTACCTTGGCCCTCTATGGCA
 GAGCAGCCGGCTCCCGTCTGGACCCGAGTCGGGCCCATGGGAACCGGGAGATACCGT
 GluGlyCysGlyTrpAlaGlyTrpLeuLeuSerProArgGlySerArgProSerTrpGly
 241 ATGAGGCTGCGGTGGCGGATGGTCTCTCTCCCGTGGCTCTCGGCCTAGCTGGG
 TACTCCGACGCCACCCGCCCTACCGAGACAGAGGGCACCGAGCCGGATCGACCC

 ProThrAspProArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCys
 301 GCCCCACAGACCCCGCGTAGGTCCGCCAATTGGGTAAAGTCAATCGATACCTTACGT
 CGGGGTGTCTGGGGCCGCATCCAGCGGTTAAACCCATTCCAGTAGCTATGGGAATGCA

 GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAla
 361 GCGGCTTCGCCGACCTCATGGGTACATACCGTCTGTCGGCGCCCTCTTGGAGCGCTG
 CGCCGAAGCGGTGGAGTACCCCATGTATGCGAGCAGCCGCGGGAGAACCTCCCGGAC
 -----overlap with CA216a-----
 ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
 421 CCAGGGCCCTGGCGCATGGGTCGGGTCTGGAAGACGCGGTGAACATATGCAACAGGA
 GTCCCGGACCGGTACCGAGGCCCAAGACCTTCTGCCGCACTTGATACGTTGTCCCT

 LeuProGlyCysSerPheSerThrPhe
 481 ACCTTCCTGGTTGCTCTTTCTCTACCTTC
 TGGAAGGACCAACGAGAAAGAGATGGAAG

FIG.12-1 Translation of DNA ag30a

#MetSerValValGlnProProGlyProProLeu

#MetAlaLeuValOP

1 CGCAGAAAGCGTCTAGCCATGGCGTTAGTATGAGTGTCTGTCAGCCTCCAGGACCCCCC
 GCGTCTTTTCGCAGATCGGTACCGCAATCATACTACAGCACGTCGGAGGTCCCTGGGGGGG

ProGlyGluProAM

61 TCCCCGGGAGAGCCATAGTGGTCTGCCGAACCGGTGAGTACACCGGAATTGCCAGGACGAC
 AGGGCCCTCTCGGTATCACCCAGACGCCCTTGCCACTCATGTGCCCTTAACGGTCTGCTG

#MetProGlyAspLeuGlyValProProGlnAsp

121 CGGGTCCCTTTCTTGATCAACCCGCTCAATGCCCTGGAGATTGGGCGTGCCCCCGCAAGA
 GCCAGGAAAGAACCTAGTTGGGCGAGTTACGGACCTCTAAACCCGCACGGGGCGTTCT

CysAM OP AM GlyAlaCys
 *

181 CTGCTAGCCGAGTAGTGTGGTCCGGAAGGCCCTTGTGGTACTGCCCTGATAGGTGCTT
 GACGATCGGCTCATCACAAACCCAGCGCTTCCGGAACACCATGACGGACTATCCCACGAA

GluCysProGlyArgSerArgProCysThrMetSerThrAsnProLysProGlnLys
 |

16 / 86

241 GCGAGTCCCCGGAGGTCTCGTAGACCGTGCACCATGAGCACGAATCCTAAACCTCAAA
CGCTACGGGGCCCTCCAGAGCATCTGGCACGTGCTACTCTGCTTAGGATTGGAGTTT
LysAsnLysArgAsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGlyGln

301 AAAAAACAAACGTAACACCAACCGTCGCCACAGGACGTCAAGTTCCCGGTGGCGGTC
TTTTTTTGTGTGCAATTGTGTTGGCAGCGGGTGTCTCTGCAGTTCAAGGCCACCGCCAG
IleValGlyGlyValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAlaThr

361 AGATCGTTGGTGGAGTTTACTTGTTCGCCGCGCAGGGGCCCTAGATTGGGTGTGCGCGGA
TCTAGCAACCACCTCAAATGAACAACGGCGCTCCCCGGGATCTAAACCCACACGCGCGCT
ArgLysThrSerGluArgSerGlnProArgGlyArgArgGlnProIleProLysAlaArg

421 CGAGAAAGACTTCCGAGCGGTCGCAACCTCGAGGTAGACGTCAGCCTATCCCCAAGGCTC
GCTCTTTCTGAAGGCTCGCCAGCGTTGGAGCTCCATCTGCAGTCGGATAGGGGTTCCCGAG
ArgProGluGlyArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsnGlu

FIG.12-2

17 / 86

-----overlap with CA290a-----
 481 GTCGGCCGAGGGCAGGACCTGGGCTCAGCCCGGTACCCCTTGCCCTCTATGGCAATG
 CAGCCGGGCTCCCGTCCGTGACCCGAGTCGGGCCCATGGGAACCGGGAGATACCGTTAC
 GlyCysGlyTrpAlaGlyTrpLeuLeuSerProArgGlySerArgProSerTrpGlyPro

 541 AGGCTGCGGGTGGCGGATGGCTCCTGTCTCCCGTGGCTCTCGGCCCTAGCTGGGGCC
 TCCCGACGCCACCCGCCCTACCGAGGACAGAGGGGCACCGAGAGCCGGATCGACCCCGG
 ThrAspProArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCysGly

 601 CCACAGACCCCGGTAGTCGCGCAATTGGGTAAGGTATCGATACCCCTTACGTGCG
 GGTGTCTGGGGCCGCATCCAGCGCGTTAAACCCATTCCAGTAGCTATGGGAATGCACGC
 Phe

 661 GCTTC
 CGAAG

* = Start of long HCV ORF
 | = Putative first amino acid of large HCV polyprotein
 # = Putative small encoded peptides (that may play a
 translational regulatory role)

FIG.12-3

FIG. 13

Translation of DNA CA205a

1 ValLeuGlyArgGluArgProCysGlyThrAlaOP AM GlyAlaCysGluCysProGly
 GTCTGGGTCGCGAAAGCCCTTGTTACTGCTGATAGGTGCTTGCGAGTCCCCCGG
 CAGAACCCAGCGCTTCCGGAAACACCATGACGGACTATCCACGAACGCTACGGGGCCC

*

61 ArgSerArgArgProCysThrMetSerThrAsnProLysProGlnArgLysThrLysArg
 AGGTCTCGTAGACCGTGACCATGAGCACGAATCCTAAACCTCAAAGAAACCAACGCT
 TCCAGAGCATCTGGCACGTGTTACTCTCGTGCTTAGGATTGGAGTTCTTTTGGTTTGCA

121

AsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGlnIleValGlyGly
 AACACCAACCGTCGCCACAGGACGTCAAGTCCCGGGTGGCGGTGAGATCGTTGGTGGA
 TTGTGTTGGCAGCGGGTGTCTCTGCAGTTCAAGGGCCACCGCCAGTCTAGCAACACCT

181

ValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAlaThrArgLysThrSer
 GTTTACTTGTGTCGCGCAGGGCCCTAGATTGGTGTGCGCGCAGACGAGAAAGACTTCC
 CAAATGAACAACGGCGCTCCCCGGGATCTAACCACACGCGCGCTGCTCTTTCTGAAGG

-----overlap with CA290a-----

241

GluArgSerGlnProArgGlyArgArgGlnProIleProLysAlaArgArgProGluGly
 GAGCGGTGCGCAACCTCGAGGTAGACGTGAGCCCTATCCCAAGGCTCGTCGCGCCGAGGGC
 CTCGCCAGCGTTGGAGCTCCATCTGCAGTCGGATAGGGGTTCCGAGCAGCCGGGCTCCCCG

301

ArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsnGluGlyCys
 AGGACCTGGGCTCAGCCCGGTACCTTGCCCTCTATGGCAATGAGGGCTGCG
 TCCTGGACCCGAGTCGGGCCCATGGGAACCGGGGAGATACCGTTACTCCCCGACGC

* = putative initiator methionine codon

19/86

FIG.14 Translation of DNA 18g

```

#ProProOP
#SerThrMetAsnHisSerProValArgAsnTyrCysLeuHisAlaGluSerValAM Pro
#LeuHisHisGluSerLeuProCysGluGluLeuSerSerArgArgLysArgLeuAla
1  CTCACCATGAATCACTCCCTGTGAGGAACACTGTCTTCACGCAGAAAGCGTCTAGCC
   GAGGTGGTACTTAGTGAGGGACACTCCTTGATGACAGAAAGTGCGTCTTTCGCAGATCGG
-----
#MetSerValValGlnProProGlyProProLeuProGlyGluProAM
MetAlaLeuValOP
61  ATGGCGTTAGTATGAGTGTGCGTGCAGCCTCCAGGACCCCTCCGGGAGAGCCATAGT
   TACCGCAATCATACTACAGCACGTCGGAGGTCTGGGGGAGGCCCTCTCGGTATCA
-----
121 GGTCTGCGGAACGGTGAGTACACCGGAATTGCCAGGACGCGGTCTTCTTGGATC
   CCAGACGCCCTTGGCCACTCATGTGGCCTTAACGGTCTCTGTGCCCCAGGAAAGAACCTAG
-----overlap with ag30a-----
#MetProGlyAspLeuGlyValProProGlnAspCysAM
181  AACCCGCTCAATGCCCTGGAGATTGGGGCGTGCCCCCGCAAGACTGCTAGCCGAGTAGTGT
   TTGGCGGAGTTACGGACCTCTAAACCCGACGGGGCGTTCTGACGATCGGCTCATCACA
-----
241  TGGGTCGCGAAAGCCCTTGTGGTACTGCCCTGATAGGTTGCTTGCAGTGCCCCGGGAGGT
   ACCCAGCGCTTTCGGAACACCATGACGGACTATCCCCACGAACTCACGGGGCCCTCCA
-----
* = Start of long HCV ORF
# = Putative small encoded peptides (that may
   play a translational regulatory role)
-----
ArgArg
301  CTCGTAGA
   GAGCATCT

```

20/86

FIG. 15 Translation of DNA 16jh

-----Overlap with 15e -----
 GlyAlaCysTyrSerIleGluProLeuAspLeuProProIleIleGlnArgLeuHisGly
 1 GGGGCCCTGCTACTCCATAGAACCACTGGATCTACCTCCAATCATTCAAAGACTCCATGGC
 CCCCCGACGATGAGGTATCTTGGTGACCTAGATGGAGGTTAGTAAGTTTCTGAGGTACCG

 LeuSerAlaPheSerLeuHisSerTyrSerProGlyGluIleAsnArgValAlaAlaCys
 61 CTCAGCGCATTTTCACTCCACAGTTACTCTCCAGGTGAAATTAATAGGGTGCCCATGC
 GAGTCGCGTAAAGTGAGGTGTCAATGAGAGGTCCACTTTAATTATCCACCGCGGTACG

 Gly*
 G
 LeuArgLysLeuGlyValProProLeuArgAlaTrpArgHisArgAlaArgSerValArg
 121 CTCAGAAAACTTGGGTACCGCCCTTGGAGCTTGGAGACACCGGGCCCGAGCGTCCGC
 GAGTCTTTTGAACCCCATGGCGGAACGCTCGAACCTCTGTGGCCCCGGGCTCGCAGGCG

 AlaArgLeuLeuAlaArgGlyGlyArgAlaAlaIleCysGlyLysTyrLeuPheAsnTrp
 181 GCTAGGCTTCTGGCCAGAGAGGAGGCTGCCATATGTGGCAAGTACCTCTTCAACTGG
 CGATCCGAAGACCGGTCTCCCTCCGTCCCGACGGTATACACCGTTTCATGGAGAGTTGACC

 AlaValArgThrLysLeuLys
 241 GCAGTAAGAACAAAGCTCAAAC
 CGTCATTCTTGTTCGAGTTTG

* = nucleotide heterogeneity

21 / 86

FIG. 16 Translation of DNA 6k

```

-----Overlap with 16jh-----
1  GlyArgAlaAlaIleCysGlyLysTyrLeuPheAsnTrpAlaValArgThrLysLeuLys
   GGCAGGGCTGCCATATGTGGCAAGTACCTCTTCAACTGGGCAGTAAGAACAAAGCTCAAA
   CCGTCCCCGACGGTATACACCGTTTCATGGAGAAAGTTGACCCCGTCATTCTTGTTCGAGTTT
   -
61  LeuThrProIleAlaAlaAlaGlyGlnLeuAspLeuSerGlyTrpPheThrAlaGlyTyr
   CTCACTCCAATAGCGCGCGCTGGCCAGCTGGACTTGTCCGGCTGGTTCACGGCTGGCTAC
   GAGTGAGGTTATCGCCGGGACCGGTGACCTGAACAGGCCGACCAAGTGCCGACCGATG

121 SerGlyGlyAspIleTyrHisSerValSerHisAlaArgProArgTrpIleTrpPheCys
   AGCGGGGAGACATTATCACAGCGTGTCTCATGCCGCGCCCGCTGGATCTGGTTTTC
   TCGCCCCCTCTGTAAATAGTGTCCGACAGAGTACGGGGCGGGGACCTAGACCAAAACG

181 CC
   GG

```

22 / 86

Translation of DNA pl31jh

-----Overlap with 6k-----
1 TyrHisSerValSerHisAlaArgProArgTrpIleTrpPheCysLeuLeuLeuAla
TTATCACAGCGTGTCTCATGCCCGCCCGCTGGATCTGGTTTGGCTACTCCTGCTTGC
AATAGTGTGCGCACAGAGTACGGGGCGGGGACCTAGACCAAAACGGATGAGGACGAAACG
61 AlaGlyValGlyIleTyrLeuLeuProAsnArgOP
TGCAGGGGTAGGCATCTACCTCCTCCCAACCGATGAAGGTGGGGTAAACACTCCGGCC
ACGTCCCATCCGTAGATGGAGGAGGGTTGGCTACTTCCAACCCCATTTGTGAGGCCGG
121 T A

FIG.17

SUBSTITUTE SHEET

23 / 86

FIG. 18-1

-341 GCCAGCCCCCTGATGGGGCGA
CGGTGGGGGACTACCCCCGCT

-319 CACTCCACCATGAATCACTCCCCCTGTGAGGAACACTACTGTCTTCACGCAGAAAGCGTCTAG
GTGAGGTGGTACTTAGTGAGGGGACACTCCTTGATGACAGAAAGTGCGTCTTTCGCAGATC

-259 CCATGGCGTTAGTATGAGTGTGTCGTCAGCCTCCAGGACCCCCCTCCCGGAGAGCCATA
GGTACCGCAATCACTACTCACAGCACGTCTGGAGGTCTCTGGGGGAGGCCCTCTCTCGGTAT

-199 GTGGTCTGCGGAACCGGTGAGTACACCGGAATTGCCAGGACGACCGGGTCTTCTTTGGA
CACCAGACGCCCTTGCCACTCATGTGGCCTTAACGGTCTCTGTGCCCCAGGAAAGAACCT

-139 TCAACCCGCTCAATGCCCTGGAGATTTGGCGGTGCCCGCAAGACTGCTAGCCGAGTAGT
AGTTGGCGGAGTTACGGACCTCTAAACCCGACGGGGCGTTCTGACGATCGGCTCATCA

- 79 GTGGGTCGCGAAAGCCCTTGTTGTTACTGCCCTGATAGGGTGTCTTGCAGTCCCCGGGAG
CAACCCAGCGCTTCCGGAAACACCATGACGGACTATCCACGAAACGCTACGGGGCCCTC

- 19 GTCTCGTAGACCGTGACCC
CAGAGCATCTGGCACGTGG

Arg Thr

MetSerThrAsnProLysProGlnLysLysAsnLysArgAsnThrAsnArgArgProGln
1 ATGAGCACGAAATCCTAAACCTCAAAAAAACAACGTAACACCAACCGTCGCCACAG
TACTCGTGCTTAGGATTGGAGTTTCTTTTGTGTCATTGTGTTGGCAGCGGGTGTCTC

AspValLysPheProGlyGlyGlyGlnIleValGlyGlyValTyrLeuLeuProArgArg
61 GACGTCAAGTTCCCGGTGGCGGTTCAGATCGTTGGTGGAGTTTACTTGTGCCGCGCAGG
CTGCAGTTCAAGGGCCACCGCCAGTCTAGCAACCACTCAATGAACAACGGCGCGTCC

SUBSTITUTE SHEET

24 / 86

FIG. 18-2

121 GlyProArgLeuGlyValArgAlaThrArgLysThrSerGluArgSerGlnProArgGly
GGCCCTAGATTGGGTGTGCGCGACGAGAAAGACTTCCGAGCGGTGCGAACCTCGAGGT
CCGGATCTAACCACACGCGCGTCTTCTGAAGGCTCGCCAGCGTTGGAGCTCCA

181 ArgArgGlnProIleProLysAlaArgArgProGluGlyArgThrTrpAlaGlnProGly
AGACGTCAGCCTATCCCCAAGGCTCGTCGCCCGAGGGCAGGACCTGGGCTCAGCCCCGG
TCTGCAGTCGGATAGGGTTCCGAGACCGGGCTCCCGTCTGGACCCGAGTCGGGCC

241 TyrProTrpProLeuTyrGlyAsnGluGlyCysGlyTrpAlaGlyTrpLeuLeuSerPro
TACCCCTTGGCCCTCTATGGCAATGAGGCTGCGGGTGGCGGATGGCTCCTGCTCCC
ATGGGAACCGGGAGATACCGTTACTCCCGACGCCACCCGCCCTACCGAGGACAGAGG

301 ArgGlySerArgProSerTrpGlyProThrAspProArgArgArgSerArgAsnLeuGly
CGTGGCTCTCGGCTAGCTGGGCCCCACAGACCCCGCGTAGGTGCGCAATTGGGT
GCACCGAGAGCCGATCGACCCCGGGGTGTCTGGGGCCGCATCCAGCGGTTAAACCCA

361 LysValIleAspThrLeuThrCysGlyPheAlaAspLeuMetGlyTyrIleProLeuVal
AAGTCAATCGATACCCCTTACGTGCGCTTCGCCGACCTCATGGGTACATACCGCTCGTC
TTCCAGTAGCTATGGGAATGCACGCCGAGCGGTGGAGTACCCCATGTATGGCGAGCAG

421 GlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAlaHisGlyValArgValLeuGluAsp
GGCGCCCTCTTGAGGCGCTGCCAGGCCCTGGCGCATGGCGTCCGGTTCTGGAAGAC
CCGCGGGAGAACCTCCGCGACGGTCCCGGACCGCGTACCGCAGGCCCAAGACCTTCTG

SUBSTITUTE SHEET

25 / 86

FIG. 18-3

481 GlyValAsnTyrAlaThrGlyAsnLeuProGlyCysSerPheSerIlePheLeuLeuAla
 GGCGTGAACTATGCAACAGGAACTTCTGGTTGCTCTTCTCTATCTTCTCTTGCC
 CCGCACTTGATACGTTGTCTCCCTTGGAAGGACCAACGAGATAGAGAGAACCGG
 Thr
 541 LeuLeuSerCysLeuThrValProAlaSerAlaTyrGlnValArgAsnSerThrGlyLeu
 CTGCTCTCTTGCTTGACTGTGCCCCCTTCGGCTTACCAAGTGGCAACTCCACGGGCTT
 GACGAGAGAACGAACTGACACGGCGGCGGATGGTTACCGCTTGAGGTGCCCCGAA
 601 TyrHisValThrAsnAspCysProAsnSerSerIleValTyrGluAlaAlaAspAlaIle
 TACCACGTACCAATGATGCCCCCTAAGTGGCTGAGTATTGTGTACGAGCGGCGGATGCCATC
 ATGGTGCAGTGTACTAACGGGATTGAGCTCATAACACATGTCCCGGCTACGGTAG
 661 LeuHisThrProGlyCysValProCysValArgGluGlyAsnAlaSerArgCysTrpVal
 CTGCACACTCCGGGTGCGTCCCTTGGCTTCGTGAGGGCAACGCCCTCGAGGTGTGGGTG
 GACGTGTAGGCCCCACGCAGGGAACGCAAGCACTCCCGTTGCGGAGCTCCACAACCCAC
 721 AlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThrGlnLeuArgArg
 GCGATGACCCCTACGGTGGCCACACGAGGATGGCAAACTCCCGCGACGACGCTTCGACGT
 CGCTACTGGGATGCCACCGGTGCTCCCTACCGTTTGAGGGGCGCTGCGTCAAGCTGCA
 781 HisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyrValGlyAspLeu
 CACATCGATCTGCTTGTGCGGAGCGCCACCTCTGTTCGGCCCTCTACGTGGGGACCTA
 GTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGGGAGATGACCCCTGGAT
 841 CysGlySerValPheLeuValGlyGlnLeuPheThrPheSerProArgArgHisTrpThr
 TGCGGGTCTGTCTTCTTGTGCGGCAACTGTTCACCTTCTCTCCAGGCGCCACTGGACG
 ACGCCAGACAGAAAGAACAGCCGGTTGACAAAGTGGAGAGAGGTCGCGGTGACCTGC

SUBSTITUTE SHEET

26 / 86

FIG. 18-4

901 ThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrp
ACGCAAGGTTGCCAATTGCTCTATCTATCCCGCCATATAACGGGTACCGCATGGCATGG
TGCGTCCAACGTTAACGAGATAGATAGGGCCGGTATATTGCCCAAGTGGCGTACCGTACC

961 AspMetMetMetAsnTrpSerProThrAlaLeuValMetAlaGlnLeuLeuArgIle
Val
GATATGATGATGAACCTGGTCCCCCTACGACGGCGTTGGTAATGGCTCAGCTGCTCCGGATC
CTATACTACTTGTACCAAGGGGATGCTGCCCAACCATTACCAGTCGACGAGGCCCTAG

1021 ProGlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAla
CCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTGGGGAGTCTGGCGGCATAGCG
GGTGTTCGGTAGAACCTGTACTAGCGACCAAGAGTGACCCCTCAGGACCGCCCGTATCGC

1081 TyrPheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGly
TATTTCTCCATGGTGGGAACCTGGCGAAGGTCCTGGTAGTGTGCTGCTATTTGCCGCGC
ATAAAGAGGTACCAACCCCTTGACCCGCTTCCAGGACCATCACGACGACGATAAACGGCCG

1141 ValAspAlaGluThrHisValThrGlyGlySerAlaGlyHisThrValSerGlyPheVal
GTCGACGCGGAACCCACGTCAACCGGGGAAGTCCCGGCCACACTGTGTCTGGATTGTGT
CAGCTGCCCTTTGGGTGCAGTGCGCCCTTACGGCCGGTGTGACACAGACCTAAACAA

1201 SerLeuLeuAlaProGlyAlaLysGlnAsnValGlnLeuIleAsnThrAsnGlySerTrp
AGCCTCCTCGCACACGCGCCAAAGCAGAACGTCCAGCTGATCAACACCAACGGCAGTTGG
TCGGAGGAGCGTGGTCCGGGTTTCGTCTTGCAGGTCGACTAGTTGTGGTTGCCGTC AAC

SUBSTITUTE SHEET

27/86

FIG. 18-5

1261 HisLeuAsnSerThrAlaLeuAsnCysAsnAspSerLeuAsnThrGlyTrpLeuAlaGly
CACCTCAATAGCACGGCCCTGAACCTGCAATGATAGCTCAACACCGGCTGGTGGCAGGG
GTGGAGTTATCGTGCCGGGACTTGACGTTACTATCGGAGTTGTGGCCGACCAACCGTCCC

1321 LeuPheTyrHisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArg
CTTTTCATCACCAAGTTCAACTCTTCAGGCTGTCTCAGAGGCTAGCCAGCTGCCGA
GAAAGATAGTGGTGTTCAGTTGAGAAGTCCGACAGGACTCTCCGATCGGTCGACGGCT

1381 ProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyPro
CCCCTTACCGATTGTGACCAGGGCTGGGGCCCTATCAGTTATGCCAACGGAAGCGCCCC
GGGAAATGGCTAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCCTTCGCCGGGG

1441 AspGlnArgProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLys
GACCAGCGCCCTACTGCTGGCACTACCCCCCAAACCTTGCGGTATTGTGCCCGGAAG
CTGGTCGCGGGATGACGACCGTGATGGGGGTTTGGAACGCCATAACACGGCGCTTC

1501 SerValCysGlyProValTyrCysPheThrProSerProValValGlyThrThrAsp
AGTGTGTGTGTCGGTATATTGCTTCACTCCAGCCCCGTGGTGGTGGAAACGACCGAC
TCACACACACAGGCCATATAACGAAGTGAGGTCGGGGCACCAACCTTGCTGGCTG

1561 ArgSerGlyAlaProThrTyrSerTrpGlyGluAsnAspThrAspValPheValLeuAsn
AGGTGGGGCGGCCACCTACAGCTGGGTGAAATGATACGACGCTCTTCGTCCTTAAC
TCCAGCCCCGCGGTGGATGTCGACCCCACTTTTACTATGCCTGCAGAACGAGGAATTG

AsnThrArgProProLeuGlyAsnTrpPheGlyCysThrTrpMetAsnSerThrGlyPhe

SUBSTITUTE SHEET

28/86

FIG. 18-6

1621 AATACCAGGCCACCGCTGGGCAATTGGTTGTTACCTGGATGAACCTCAACTGGATTC
TTATGGTCCGGTGGCGACCCGTTAACCAAGCCAACATGGACCTACTTGAGTTGACCTAAG

1681 ThrLysValCysGlyAlaProProCysValIleGlyGlyAlaGlyAsnAsnThrLeuHis
ACCAAAGTGTGGGAGCGCTCCTTGTGTCTATCGAGGGCGGCAACAACACCTGCAC
TGGTTTCACACGCCCTCGCGGAGGAACACAGTAGCCTCCCCCGCTTGTGTGGACGTG

1741 CysProThrAspCysPheArgLysHisProAspAlaThrTyrSerArgCysGlySerGly
TGCCCCACTGATTGCTTCGCAAGCATCCGGACGCCACATACCTCTCGGTGCGGCTCCGGT
ACGGGGTGACTAAACGAAGCGTTCGTAGGCCCTGCGGTGTATGAGAGCCACGCCGAGGCCA

Ile

1801 ProTrpLeuThrProArgCysLeuValAspTyrProTyrArgLeuTrpHisTyrProCys
CCCTGGATCACACCCAGGTGCTGTCGACTACCCGTATAGGCTTTGGCATATATCCTTGT
GGGACCTAGTGTGGTCCACGGACCAGCTGATGGGCATATCCGAAACCGTAATAGGAACA

1861 ThrIleAsnTyrThrIlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeu
ACCATCAACTACACCATATTTAAATCAGGATGTACGTGGGAGGGTCTGAACACACAGGCTG
TGGTAGTTGATGTGGTATAAATTTTAGTCCACATGCACCCCTCCCGAGCTTGTGTCCGAC

1921 GluAlaAlaCysAsnTrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSer
GAAGCTGCCCTGCAACTGGACGCGGGCGAACGTTGCCGATCTGGAAGACAGGACAGGTCC
CTTCGACGGACGTTGACCTGCGCCCCCGCTTGCAACGCTAGACCTTCTGTCCCTGTCCAGG

1981 GluLeuSerProLeuLeuLeuThrThrGlnTrpGlnValLeuProCysSerPheThr
GAGCTCAGCCCGTTACTGCTGACCACTACACAGTGGCAGGTCTCCCGTGTCTCTTCACA
CTCGAGTCGGGCAATGACGACTGGTGTATGTGTACCGTCCAGGAGGCACAGGAAGTGT

SUBSTITUTE SHEET

29 / 86

FIG. 18-7

ThrLeuProAlaLeuSerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGln
2041 ACCCTACCAGCCTTGTCACCGCCTCATCCACCTCCACAGAACATTTGTGGACGTGCAG
TGGGATGGTCGGAACAGGTGGCCGGAGTAGGTGGAGTGGTCTTGTAAACACCTGCACGTC
TyrLeuTyrGlyValGlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValVal
2101 TACTTGTAACGGGTGGGTCAAGCATCGCTCCTGGGCCATTAAAGTGGAGTACGTCGTT
ATGAACATGCCCCACCCAGTTCGTAGCGCAGGACCCGGTAATTACCCCTCATGCAGCAA
LeuLeuPheLeuLeuAlaAlaArgValCysSerCysLeuTrpMetMetLeuLeu
2161 CTCCTGTTCTCTGCTTGCAAGCGCGCTCTGCTCTGCTTGTGGATGATGCTACTC
GAGGACAAGGAAGACGAACGTCTGCGCGCAGACGAGGACGAACACCTACTACGATGAG
IleSerGlnAlaGluAlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAla
2221 ATATCCCAAGCGGAGCGGCTTTGGAGAACCTCGTAATACTTAATGCAGCATCCCTGGCC
TATAGGTTGCGCTCCGCCGAAACCTCTTGGAGCATTAATGAATTACGTCGTAGGGACCCGG
GlyThrHisGlyLeuValSerPheLeuValPhePheCysPheAlaTrpTyrLeuLysGly
2281 GGGACGCACGGTCTTGTAATCCTTCCCTCGTGTCTCTGCTTGCATGGTATTGAAGGGT
CCCTGCGTGCCAGAACATAGGAAGGAGCACAAAGACGAAACGTACCATAAACTTCCCA
LysTrpValProGlyAlaValTyrThrPheTyrGlyMetTrpProLeuLeuLeuLeu
2341 AAGTGGTGGCCCGGAGCGGTCTACACCTTCTACGGGATGTGGCCTCTCCCTGCTCCTG
TTCACCCACGGGCTCGCCAGATGTGGAAGATGCCCTACACCGGAGAGGACGAGGAC

SUBSTITUTE SHEET

30/86

FIG. 18-8

2401 LeuAlaLeuProGlnArgAlaTyrAlaLeuAspThrGluValAlaAlaSerCysGlyGly
TTGGCGTTGCCCGCCAGCGGGCTACGCGTGGACACGGAGTGCCCGCTCGTGTGGCGGT
AACCGCCAACGGGTCCGCCCGCATGCCGACCTGTGCTCCACCGCGCAGCACACCGCCA

2461 ValValLeuValGlyLeuMetAlaLeuThrLeuSerProTyrTyrLysArgTyrIleSer
GTTGTTCTCGTCGGTGTGATGGCGCTGACTCTGTCAACCATATTACAAGCGCTATATCAGC
CAACAAGAGCAGCCCCAACTACCGGCTGAGACAGTGGTATAATGTTCCGATATAGTCG

(Asn)

2521 TrpCysLeuTrpTrpLeuGlnTyrPheLeuThrArgValGluAlaGlnLeuHisValTrp
TGGTGCTTGTGGTGGCTTCAGTATTTTCTGACCAAGAGTGGAAGCGCAACTGCACGTGTGG
ACCACGAACACCAACCGAAGTCATAAAGACTGGTCTCACCTTCGCGTTGACGTGCACACC

IleProProLeuAsnValArgGlyGlyArgAspAlaValIleLeuLeuMetCysAlaVal

2581 ATTCCCCCTCAACGTCCGAGGGGGCGCGCGCTCATCTTACTCATGTGTGTGTGTA
TAAGGGGGGAGTTGCAGGCTCCCCCGCGCTGCGCAGTAGAATGATACACACGACAT

2641 HisProThrLeuValPheAspIleThrLysLeuLeuAlaValPheGlyProLeuTrp
CACCCGACTCTGGTATTGACATCACCAAAATTGCTGTGGCCGTCTTCGGACCCCTTTGG
GTGGGCTGAGACCATAAACTGTAGTGGTTTAACGACGACCGGCAGAACCTGGGGAAACC

2701 IleLeuGlnAlaSerLeuLeuLysValProTyrPheValArgValGlnGlyLeuLeuArg
ATTCTTCAAGCCAGTTTGCTTAAAGTACCCTACTTTGTGCGCGTCCAAAGCCTTCTCCGG
TAAGAAGTTCGGTCAACGAATTTCATGGGATGAACACACGCGCAGGTTCGGGAAGAGGCC

SUBSTITUTE SHEET

31 / 86

FIG. 18-9

2761 PheCysAlaLeuAlaArgLysMetIleGlyGlyHisTyrValGlnMetValIleIleLys
 TTCCTGCGCGTTAGCGCGAAGATGATCGGAGGCCATTACGTGCAAAATGGTCATCATTAAG
 AAGACGCGCAATCGCGCCTTCTACTAGCCTCCGGTAATGCACGTTTACCAGTAGTAATTC

2821 LeuGlyAlaLeuThrGlyThrTyrValTyrAsnHisLeuThrProLeuArgAspTrpAla
 TTAGGGGCGCTTACTGGCACCTATGTTTATAAACCATCTCACTCCTCTTCGGGACTGGGCG
 AATCCCCCGGAATGACCGTGTGATACAAATATTGTTAGAGTGAGGAGAGCCCTGACCCGC

2881 HisAsnGlyLeuArgAspLeuAlaValAlaValGluProValValPheSerGlnMetGlu
 CACAACGGCTTTCGAGATCTGGCCGTGGCTGTAGAGCCAGTCGTCTTCTCCCAAATGGAG
 GTGTTGCCGAACGCTCTAGACCGGCACCGACATCTCGGTACGACAGAGGGTTTACCTC

2941 ThrLysLeuIleThrTrpGlyAlaAspThrAlaAlaCysGlyAspIleIleAsnGlyLeu
 ACCAAGCTCATCACGTGGGGGCAGATACCGCCGTCGCGTGACATCATCAACGGCTTG
 TGGTTCGAGTAGTGCAACCCCGCTCTATGGCGGCACGCCACTGTAGTAGTTGCCGAAC

3001 ProValSerAlaArgArgGlyArgGluIleLeuLeuGlyProAlaAspGlyMetValSer
 CCTGTTCGCCCGCAGGGCCGGAGATACTGCTCGGGCCAGCCGATGGAATGGTCTCC
 GGACAAAGCGGGCGTCCCCGCCCTCTATGACGAGCCCGGTCTACCTTACCAGAGG

3061 LysGlyTrpArgLeuLeuAlaProIleThrAlaTyrAlaGlnGlnThrArgGlyLeuLeu
 AAGGGTGGAGGTGCTGGCGCCCATCACGGCGTACGCCCCAGCAGACAAAGGGCCTCCTA
 TTCCCCACCTCCAAACGACCGGGTAGTGCCGCGCATGCGGTCTGTTCCTCCGGAGGAT

3121 GlyCysIleIleThrSerLeuThrGlyArgAspLysAsnGlnValGluGlyGluValGln
 GGGTGCAATAATCACAGCCTAACTGGCCGGACAAAACCAAGTGGAGGGTGAGGTCCAG
 CCCACGTATTAGTGGTCGGATTGACCGGCCCTGTTTTGGTTTCACTCCACTCCAGGTC

SUBSTITUTE SHEET

32 / 86

FIG. 18-10

IleValSerThrAlaAlaGlnThrPheLeuAlaThrCysIleAsnGlyValCysTrpThr
 3181 ATTGTGTCAACTGCTGCCCAAACTTCTCTGGCAACGTGCATCAATGGGTGTGCTGGACT
 TAACACAGTTGACGACGGGTTTGGAAAGGACCGTTGCACGTAGTTACCCACACGACCTGA

 ValTyrHisGlyAlaGlyThrArgThrIleAlaSerProLysGlyProValIleGlnMet
 3241 GTCTACACGGGGCGGAACGAGGACCATCGCGTCACCCAAAGGTCTCTGTCAATCCAGATG
 CAGATGGTCCCCCGGCTTGTCTCTGTAGCGCAGTGGGTTCCAGGACAGTAGGTCTAC

 TyrThrAsnValAspGlnAspLeuValGlyTrpProAlaProGlnGlySerArgSerLeu
 3301 TATACCAATGTAGACCAAGACCTTGTGGGCTGGCCCGCTCCGCAAGGTAGCCGCTCATTTG
 ATATGGTTACATCTGGTTCTGTGAACACCCGACCGGGCGAGCGTTCCATCGGCGAGTAAC

 ThrProCysThrCysGlySerSerAspLeuTyrLeuValThrArgHisAlaaspValIle
 3361 ACACCTGCACCTGCGGCTCCTCGGACCTTTACCTGGTACGAGGACGACCGCATGTCAATT
 TGTGGACGTGAACGCCGAGGAGCCTGGAAATGGACCAGTGTCCGTCCGGGCTACAGTAA

 ProValArgArgGlyAspSerArgGlySerLeuLeuSerProArgProIleSerTyr
 3421 CCGGTGCGCGCGGGGTGATAGCAGGGCAGCCTGTCTGTGCCCCCGGCCCATTTCCCTAC
 GGGCACGCGCGCGCCCACTATCGTCCCCGTGGACGACAGCGGGCGGGGTAAAGGATG

 LeuLysGlySerSerGlyGlyProLeuLeuCysProAlaGlyHisAlaValGlyIlePhe
 3481 TTGAAAGGCTCCTCGGGGTCCGCTGTGTGCCCCCGGGGCGACCGCGTGGGCATATTT
 AACTTTCCGAGGAGCCCCCAGGCGGACACACGGGGCGCCCCGTGGCACCCCGTATAAA

 ArgAlaAlaValCysThrArgGlyValAlaLysAlaValAspPheIleProValGluAsn
 3541 AGGGCCGGGTGTGCACCCGTGGAGTGGCTAAGCGGTGGACTTTATCCCTGTGGAGAAC

SUBSTITUTE SHEET

33 / 86

FIG. 18-11

TCCCGGCGCCACACGTGGGCACCTCACCGATTCCGCCACCTGAAATAGGGACACCTCTTG

3601 LeuGluThrThrMetArgSerProValPheThrAspAsnSerSerProProValValPro
CTAGAGACAACCATGAGGTCCCGGTGTTACGGATAACTCTCTCCACCAGTAGTGCCC
GATCTGTGTGTTACTCCAGGGGCCACAAGTGCCCTATTGAGGAGAGGTGTCATCACGGG

3661 GlnSerPheGlnValAlaHisLeuHisAlaProThrGlySerGlyLysSerThrLysVal
CAGAGCTTCCAGGTGGCTCACCTCCATGCTCCACAGGCAGCGGCAAAGCAACCAAGGTC
GTCTCGAAGGTCCACCGAGTGGAGGTACGAGGGTGTCCTCGCCGTTTTCGTGGTTCCAG

3721 ProAlaAlaTyrAlaAlaGlnGlyTyrLysValLeuValLeuAsnProSerValAlaAla
CCGGCTGCATATGCAGCTCAGGGCTATAAGGTGCTAGTACTCAACCCCTCTGTGTGCTGCA
GGCCGACGTATACGTCCGAGTCCCGATATTCACGATCATGAGTTGGGGAGACAACGACGT

Leu

3781 ThrLeuGlyPheGlyAlaTyrMetSerLysAlaHisGlyIleAspProAsnIleArgThr
ACACTGGGCTTTGGTGCTTACATGTCCAAGGCTCATGGGATCGATCCTAACATCAGGACC
TGTGACCCGAAACCACGAATGTACAGTTCCGAGTACCCTAGCTAGGATTGTAGTCCCTGG

3841 GlyValArgThrIleThrThrGlySerProIleThrTyrSerThrTyrGlyLysPheLeu
GGGGTGAGAACAAATTACCACCTGGCAGCCCCCATCACGTACTCCACCTACGGCAAGTTCTT
CCCCACTCTTGTAAATGTTGACCGTCGGGTAGTGCATGAGGTGGATGCCGTTCAAGGAA

3901 AlaAspGlyGlyCysSerGlyGlyAlaTyrAspIleIleIleCysAspGluCysHisSer
GCCGACGGCGGTGCTCGGGGGCGCTTATGACATAATAATTGTGACGAGTGCCACTCC
CGGCTGCCGCCACGAGCCCCCGGAATACTGTATTATTAAACACTGCTCACGGTGAGG

SUBSTITUTE SHEET

34 / 86

FIG. 18-12

(Val)

3961 ThrAspAlaThrSerIleLeuGlyIleGlyThrValLeuAspGlnAlaGluThrAlaGly
ACGGATGCCACATCCATCTTGGCATCGCACTGTCTTGACCAAGCAGAGACTGCGGGG
TGCCCTACGGTGTAGTAGAACCCGTAGCCGTGACAGGAAGTGGTTCGTCTCTGACGCCCC

4021 AlaArgLeuValValLeuAlaThrAlaThrProProGlySerValThrValProHisPro
GCGAGACTGGTTGTGCTCGCCACCGCCACCCCTCCGGGCTCCGTCACTGTGCCCATCCC
CGCTCTGACCAACACGAGCGGTGGCGGTGGGAGGCCCGAGGCAGTGACACGGGGTAGGG

4081 AsnIleGluGluValAlaLeuSerThrThrGlyGluIleProPheTyrGlyLysAlaIle
AACATCGAGGAGGTGTCTCTGTCCACCAACCGGAGAGATCCCTTTTACGGCAAGGCTATC
TTGTAGCTCCTCCAAACGAGACAGGTGGTGGCTCTCTAGGGAAAAATGCCGTTCCGATAG

4141 ProLeuGluValIleLysGlyGlyArgHisLeuIlePheCysHisSerLysLysLysCys
CCCCTCGAAGTAATCAAGGGGGGAGACATCTCATCTTCTGTCAATTCAAAGAAGAGTGC
GGGAGCCTTCATTAGTTCCCCCTCTGTAGAGTAGAAGACAGTAAGTTTCTTCTTCACG

4201 AspGluLeuAlaAlaLysLeuValAlaLeuGlyIleAsnAlaValAlaTyrTyrArgGly
GACGAACCTGCCCGCAAGCTGGTCGCAATGGGCATCAATGCCGTGCCCTACTACCGCGGT
CTGCTTGAGCGCGGTTTCGACCAAGCGTAACCCGTAGTTACGGCACCGGATGATGGCGCCA

4261 LeuAspValSerValIleProThrSerGlyAspValValValAlaThrAspAlaLeu
CTTGACGTGTCCGTCAATCCGACCAAGCGCATGTGTCTCGTGGCAACCGATGCCCTC
GAACTGCACAGGCAGTAGGGCTGGTCGCCGCTACAAACAGCAGCACCGTTGGCTACGGGAG

SUBSTITUTE SHEET

35 / 86

FIG. 18-13

Tyr

MetThrGlyTyrThrGlyAspPheAspSerValIleAspCysAsnThrCysValThrGln
 ATGACCGGTATACCGCGACTTCGACTCGGTGATAGACTGCAATACGTGTGTACCCAG
 TACTGGCCGATATGGCCGCTGAAGCTGAGCCACTATCTGACGTTATGCACACAGTGGGTC

4321

(Ser)

ThrValAspPheSerLeuAspProThrPheThrIleGluThrIleThrLeuProGlnAsp
 ACAGTCGATTTCAGCCTTGACCCCTACCTTCACCAATTGAGACAAATCACGCTCCCCAGGAT
 TGTACGCTAAAGTCGGAACTGGGATGGAGTGTTAACTCTGTAGTGCAGGGGGTCCCTA

4381

AlaValSerArgThrGlnArgArgGlyArgThrGlyArgGlyLysProGlyIleTyrArg
 GCTGTCTCCCGCACTCAACGTCGGGCGAGGACTGGCAGGGGAGCCAGGCATCTACAGA
 CGACAGAGGGCGTGAGTTGCAGCCCCCGTCCCTGACCGTCCCCCTTCGGTCCGTAGATGTCT

4441

PheValAlaProGlyGluArgProSerGlyMetPheAspSerSerValLeuCysGluCys
 TTTGTGGCACCGGGGAGCGCCCTCCGGCATGTTTCGACTCGTCCGTCTCTGTGAGTGC
 AAACACCGTGGCCCCCTCGGGGGAGGCCGTACAAGCTGAGCAGGACGAGACACTCAGC

4501

TyrAspAlaGlyCysAlaTrpTyrGluLeuThrProAlaGluThrThrValArgLeuArg
 TATGACGACAGGCTGTGCTTGGTATGAGCTACGCCCGCCGAGACTACAGTTAGGCTACGA
 ATACTGCGTCCGACACGAACCATACTCGAGTGCGGGCGGCTCTGATGTCAATCCGATGCT

4561

AlaTyrMetAsnThrProGlyLeuProValCysGlnAspHisLeuGluPheTrpGluGly
 GCGTACATGAACACCCCGGGCTTCCCGTGTGCCAGGACCATCTTGAATTTTGGAGGGC
 CGCATGTACTTGTGGGGCCCCGAAGGGCACACGGTCTCTGGTAGAACTTAAACCCCTCCCG

4621

ValPheThrGlyLeuThrHisIleAspAlaHisPheLeuSerGlnThrLysGlnSerGly
 GTCTTTACAGGCCCTCACTCATATAGATGCCCACTTCTATCCAGACAAAGCAGAGTGGG
 CAGAAATGTCCGGAGTGAGTATATCTACGGGTGAAAGATAGGGTCTGTTCGTCTCACCC

4681

SUBSTITUTE SHEET

36 / 86

FIG. 18-14

GluAsnLeuProTyrLeuValAlaTyrGlnAlaThrValCysAlaArgAlaGlnAlaPro
 4741 GAGAACCTTCCCTTACCTGGTAGCGTACCAAGCCACCGTGTCGGCTAGGGCTCAAGCCCCCT
 CTCTTGGAAAGGAATGGACCATCGCATGGTTCCGTGGCACACGCCGATCCCAGTTCGGGGGA

 ProProSerTrpAspGlnMetTrpLysCysLeuIleArgLeuLysProThrLeuHisGly
 4801 CCCCATCGTGGGACCAGATGTGGAAGTGTGTGATTGCTCGCTCAAGCCACCCCTCCATGGG
 GGGGTAGCACCCCTGTCTACACCTTCACAACTAAGCGGAGTTCGGGTGGGAGGTACCC

 ProThrProLeuTyrArgLeuGlyAlaValGlnAsnGluIleThrLeuThrHisPro
 4861 CCAACACCCCTGCTATACAGACTGGCGCTGTTTCAGAAATGAATCACCCCTGACGCACCCCA
 GGTGTGGGACGATATGCTGACCCCGCACAAGTCTTACTTAGTGGGACTGCGTGGGT

 ValThrLysTyrIleMetThrCysMetSerAlaAspLeuGluValThrSerThrTrp
 4921 GTCACCAATAATACATCATGACATGTCGCGCCGACCTGGAGGTCTGCACGAGCACCTGG
 CAGTGGTTTATGTAGTACTGTACGTACAGCCGGCTGGACCTCCAGCAGTGTCTCGTGGACC

 ValLeuValGlyGlyValLeuAlaAlaLeuAlaAlaTyrCysLeuSerThrGlyCysVal
 4981 GTGCTCGTTGGCGGCGTCTGGCTGCTTGGCCCGCTATTGCCCTGTCAACAGGCTGCGTG
 CACGAGCAACCGCCGACGACCGACGAAACCGCGGCATAACGGACAGTTGTCCGACGCAC

 ValIleValGlyArgValValLeuSerGlyLysProAlaIleProAspArgGluVal
 5041 GTCATAGTGGCAGGGTCGTCTTGTCCGGGAAGCCGCAATCATACCTGACAGGGAAGTC
 CAGTATCACCCGTCCTCCAGCAGAACAGGCCCTTCGGCCGTTAGTATGGACTGTCCCTTCAG

 LeuTyrArgGluPheAspGluMetGluGluCysSerGlnHisLeuProTyrIleGluGln
 5101 CTCTACCGAGAGTTCGATGAGATGGAAGAGTGTCTCTCAGCAGCTTACCGTACATCGAGCAA
 GAGATGGCTCTCAAGCTACTCTACCTTCTCAGCAGAGTCTGTGAATGGCATGTAGCTCGTT

SUBSTITUTE SHEET

37/86

FIG. 18-15

5161 GlyMetMetLeuAlaGluGlnPheLysGlnLysAlaLeuGlyLeuLeuGlnThrAlaSer
GGGATGATGCTCGCCGAGCAGTTCAAGCAGAGGCCCTCGGCCTCCTGCAGACCGCGTCC
CCCTACTACGAGCGGCTCGTCAAGTTCGTCTCCGGGAGCCGGAGGACGTCTGGCGCAGG

5221 ArgGlnAlaGluValIleAlaProAlaValGlnThrAsnTrpGlnLysLeuGluThrPhe
CGTCAGGCAGAGGTTATCGCCCCCTGCTGTCCAGACCAACTGGCAAAACTCGAGACCTTC
GCAGTCCGTCTCCAATAGCGGGACGACAGTCTGTTGACCGTTTTTTGAGCTCTGGAAG

5281 TrpAlaLysHisMetTrpAsnPheIleSerGlyIleGlnTyrLeuAlaGlyLeuSerThr
TGGGCGAAGCATATGTGGAACCTTCATCAGTGGGATACAAATACTTGGCGGGCTTGTCAACG
ACCCGCTTCGTATACACCTTGAAGTAGTCACCCCTATGTATGAACCGCCCGAACAGTTGC

5341 LeuProGlyAsnProAlaIleAlaSerLeuMetAlaPheThrAlaAlaValThrSerPro
CTGCCCTGGTAACCCCGCCATTGCTTCATTGATGGCTTTTACAGCTGCTGTCAACGCCCCA
GACGGACCATTTGGGGCGGTAAACGAAGTAATACTACCGAAATGTGACGACAGTGTCCGGGT

5401 LeuThrThrSerGlnThrLeuLeuPheAsnIleLeuGlyGlyTrpValAlaAlaGlnLeu
CTAACCCACTAGCCAAACCCCTCCTCTTCAACATATTGGGGGGTGGTGGCTGCCAGCTC
GATTGGTGATCGGTTTGGGAGGAGAGTTGTATATAACCCCCCACCACCGGCTCGAG

5461 AlaAlaProGlyAlaAlaThrAlaPheValGlyAlaGlyLeuAlaGlyAlaIleGly
GCCGCCCCCGGTGCGGCTACTGCCCTTTGTGGCGCTGGCTTAGCTGGCGCCGCCATCGGC
CGCGGGGGCCACGGCGATGACGGAAACACCCGCGACCGAATCGACCGCGGTAGCCG

SUBSTITUTE SHEET

38 / 86

FIG. 18-16

5521 SerValGlyLeuGlyLysValLeuIleAspIleLeuAlaGlyTyrGlyAlaGlyValAla
 AGTGTGGACTGGGAAGGTCTCATAGACATCTTGAGGGTATGCGCGGCGTGCGG
 TCACAACCTGACCCCTTCCAGGAGTATCTGTAGGAACGTCCCATACCGGCCCCGACCGC

5581 GlyAlaLeuValAlaPheLysIleMetSerGlyGluValProSerThrGluAspLeuVal
 (Gly)
 GGAGCTCTTGCGCATTCAGATCATGAGCGGTGAGGTCCCTCCACGAGGACCTGGTC
 CCTCGAGAACACCGTAAGTTCTAGTACTCGCCACTCCAGGGAGGTGCTCTGGACCAG

5641 AsnLeuLeuProAlaIleLeuSerProGlyAlaLeuValGlyValValCysAlaAla
 AATCTACTGCCCCGCATCTCTCGCCCGAGCCCTCGTAGTCGGCTGTGTGCAGCA
 TTAGATGACGGCGGTAGGAGAGCGGCCCTCGGAGCATCAGCCGCACAGACACGTCGT

5701 IleLeuArgArgHisValGlyProGlyGluGlyAlaValGlnTrpMetAsnArgLeuIle
 ATACTGCGCCGGCACGTTGGCCCGGGCGAGGGGCGAGTGCAGTGGATGAACCGGCTGATA
 TATGACGGGCGCGTGCAACCGGCCCGCTCCCGTCACGTACCTACTTGGCCGACTAT

5761 AlaPheAlaSerArgGlyAsnHisValSerProThrHisTyrValProGluSerAspAla
 GCCTTCGCCCTCCCGGGGAACCATGTCTTCCCCACGCACCTACGTCCCGAGAGCGATGCA
 CGGAAGCGGAGGCCCCCTTGGTACAAAGGGGTGCGTGATGCACGGCCTCTCGCTACGT

5821 AlaAlaArgValThrAlaIleLeuSerSerLeuThrValThrGlnLeuLeuArgArgLeu
 (HisCys)
 GGTGCCCGCGTCACTGCCATACCTACGAGCCCTCACTGTAAACCCAGCTCTGAGGCGACTG
 CGACGGGCGCAGTGACGGTATGAGTCGTGCGAGTGACATTGGGTGCGAGGACTCCGCTGAC

SUBSTITUTE SHEET

39/86

FIG. 18-17

5881 HisGlnTrpIleSerSerGluCysThrThrProCysSerGlySerTrpLeuArgAspIle
 CACCAAGTGGATAAGCTCGAGTGTACCACTCCATGCTCCGGTTCCTGGCTAAGGGACATC
 GTGGTCACCTATTTCGAGCCTCACATGGTGAGGTACGAGGCCAAGGACCGATTCCCTGTAG

5941 TrpAspTrpIleCysGluValLeuSerAspPheLysThrTrpLeuLysAlaLysLeuMet
 TGGGACTGGATATGCGAGGTGTGAGCGACTTTAAGACCTGGCTAAAGCTAAGCTCATG
 ACCCTGACCTATACGCTCCACAACCTCGCTGAAATTCTGGACCGATTTCGATTTCGAGTAC

6001 ProGlnLeuProGlyIleProPheValSerCysGlnArgGlyTyrLysGlyValTrpArg
 CCACAGCTGCCCTGGGATCCCCCTTTGTCTCCTGCCAGCGGGGTATAAGGGGTCTGGCGGA
 GGTGTCGACGGACCCCTAGGGGAAACACAGGACGGTCGCGGCCCATATTCCCCCAGACCGCT

(Val)
 6061 GlyAspGlyIleMetHisThrArgCysHisCysGlyAlaGluIleThrGlyHisValLys
 GTGGACGGCATCATGCACACTCGCTGCCACTGTGGAGCTGAGATCACTGGACATGTCAAA
 CACCTGCCGTAGTACGTGTGAGCGACGGTGACACCTCGACTCTAGTGACCTGTACAGTTT

6121 AsnGlyThrMetArgIleValGlyProArgThrCysArgAsnMetTrpSerGlyThrPhe
 AACGGGACGATGAGGATCGTCGGTCCCTAGGACCTGCAGGAACATGTGGAGTGGACCTTC
 TTGCCCTGCTACTCCTAGCAGCCAGGATCCTGGACGTCTTGTACACCTCACCCCTGGAAG

6181 ProIleAsnAlaTyrThrThrGlyProCysThrProLeuProAlaProAsnTyrThrPhe
 CCCATTAAATGCCCTACACCAACGGGCCCTGTACCCCCCTTCCTGCGCCGAACACACGTTC
 GGGTAATTACGGATGTGGTGTCCCCGGGACATGGGGGGAAGGACGCGGCTTGATGTGCAAG

SUBSTITUTE SHEET

40/86

FIG. 18-18

6241 AlaLeuTrpArgValSerAlaGluGluTyrValGluIleArgGlnValGlyAspPheHis
 GCGCTATGGAGGGTGTCTGCAGAGGAATATGTGGAGATAAGGCAGGTGGGACCTCCAC
 CGCGATACCTCCACAGACGTCTCCTTATACACCTCTATTCCGTCCACCCCTGAAGGTG

6301 TyrValThrGlyMetThrThrAspAsnLeuLysCysProCysGlnValProSerProGlu
 TACGTGACGGGTATGACTACTGACAATCTCAATGCCCGTGCCAGGTCCCATCGCCCGAA
 ATGCACTGCCCATACTGATGACTGTAGAGTTTACGGGCACGGTCCAGGTAGCGGGCTT

6361 PhePheThrGluLeuAspGlyValArgLeuHisArgPheAlaProProCysLysProLeu
 TTTTTCACAGAAATTGGACGGGTGCGCTACATAGGTTTGCGCCCCCTGCAAGCCCTTG
 AAAAGTGCTTAACTTGCCCCACGGCGATGTATCCAAACGCGGGGGACGTTTCGGGAAC

LeuArgGluGluValSerPheArgValGlyLeuHisGluTyrProValGlySerGlnLeu

6421 CTGCGGGAGGAGGTATCATTCAGAGTAGGACTCCACGAATACCCGGTAGGGTCGCAATTA
 GACGCCCTCCTCCATAGTAAGTCTCATCTGAGGTGCTTATGGGCCCATCCCCAGCGTTAAT

6481 ProCysGluProGluProAspValAlaValLeuThrSerMetLeuThrAspProSerHis
 CCTTGCGAGCCCGAACCGGACGTGGCCGTGTGACGTCCATGCTCACTGATCCTCCCAT
 GGAAACGCTCGGGCTTGGCCTGCACCCGGCACAACTGCAGGTACGAGTGACTAGGGAGGTA

6541 IleThrAlaGluAlaAlaGlyArgArgLeuAlaArgGlySerProProSerValAlaSer
 ATAACAGCAGAGGCGGCCGCGAAGTTGGCGAGGGGATCACCCCCCTCTGTGGCCAGC
 TATTGTCTCTCCGCCGCCCGCTTCCAAACCGCTCCCCCTAGTGGGGGAGACACCGGTCTG

SUBSTITUTE SHEET

41 / 86

FIG. 18-19

6601 SerSerAlaSerGlnLeuSerAlaProSerLeuLysAlaThrCysThrAlaAsnHisAsp
TCCTCGGCTAGCCAGCTATCCGCTCCTCACTCTCAAGCAACTTGACCGCTAACCATGAC
AGGAGCCGATCGGTCGATAGCGGAGTAGAGAGTTCCGTTGAACGTGGCGATTGGTACTG

6661 SerProAspAlaGluLeuIleGluAlaAsnLeuLeuTrpArgGlnGluMetGlyGlyAsn
TCCCCCTGATGCTGAGCTCATAGAGGCCAACCTCCTATGGAGGACAGAGATGGCGGCAAC
AGGGACTACGACTCGAGTATCTCCGGTTGGAGGATACCTCCGTCCTCTACCCCGCCGTTG

6721 IleThrArgValGluSerGluAsnLysValIleLeuAspSerPheAspProLeuVal
ATCACCAAGGTTGAGTCAGAAACAAAGTGGTGATTTCTGGACTCCTTCGATCCGCTTGTG
TAGTGTCCCCAACTCAGTCTTTTGTTCACCACTAAGACCTGAGGAAGCTAGGCGCAACAC

6781 AlaGluGluAspGluArgGluIleSerValProAlaGluIleLeuArgLysSerArgArg
GCGGAGGAGGACGAGCGGGAGATCTCCGTACCCGCAGAAATCCTGCGGAAGTCTCGGAGA
CGCCTCCTCCTGCTCGCCCTCTAGAGGCATGGCGCTCTTAGGACGCCCTTCAGAGCCTCT

6841 PheAlaGlnAlaLeuProValTrpAlaArgProAspTyrAsnProProLeuValGluThr
TTCGCCAGGCCCTGCCCGTTTGGCGCGCGGACTATAACCCCGCTAGTGGAGACG
AAGCGGTCCGGGACGGGCAACCCGCGCGCTGATATTGGGGGCGGATCACCTCTGC

6901 TrpLysLysProAspTyrGluProProValValHisGlyCysProLeuProProLys
TGGAATAAGCCCGACTACGAACCACTGTGGTCCATGGCTGTCCGCTTCCACCTCCAAAG
ACCTTTTTCGGGCTGATGCTTGGTGGACACCAAGTACCCGACAGCGGAAGGTGGAGTTTC

6961 SerProProValProProArgLysLysArgThrValValLeuThrGluSerThrLeu
TCCCCCTCCTGTGCTCCGCCCTCGGAAGAACGGACGGTGGTCTCCTCACTGAATCAACCCCTA
AGGGAGGACACGGAGCGGAGCCTTCTTCGCCCTGCCACCAAGGAGTGACTTAGTTGGGAT

SUBSTITUTE SHEET

42/86

FIG. 18-20

(Ser.)

SerThrAlaLeuAlaGluLeuAlaThrArgSerPheGlySerSerSerThrSerGlyIle
 TCTACTGCCCTTGCCGAGCTCGCCACACAGAAAGCTTTGGCAGCTCCTCAACTTCCGGCATTT
 AGATGACGGAACCGGCTCGAGCGGTGCTTCGAAACCGTCGAGGAGTTGAAGGCCCGTAA

7021

ThrGlyAspAsnThrThrSerSerGluProAlaProSerGlyCysProProAspSer
 ACGGGCACAATACGACAACATCCTCTGAGCCCGCCCTTCTGGCTGCCCCCGACTCC
 TGCCCCGCTGTATGCTGTGTAGGAGACTCGGGCGGGGAAGACCGACGGGGGGCTGAGG

7081

(PheAla)

AspAlaGluSerTyrSerSerMetProProLeuGluGlyGluProGlyAspProAspLeu
 GACGCTGAGTCCTATTCTCTCCATGCCCCCTGGAGGGGAGCCTGGGATCCGGATCTT
 CTGCGACTCAGGATAAGGAGGTACGGGGGGACCTCCCCCTCGGACCCCTAGGCCCTAGAA

7141

SerAspGlySerTrpSerThrValSerSerGluAlaAsnAlaGluAspValValCysCys
 AGCGACGGGTCAATGTCACGGTCAGTAGTGAGGCCAACCGGAGGATGTCGTGTGCTGC
 TCGCTGCCCCAGTACCAGTTGCCAGTCATCCTCCGGTTGGCCCTCTACAGCACACGACG

7201

SerMetSerTyrSerTrpThrGlyAlaLeuValThrProCysAlaAlaGluGluGlnLys
 TCAATGTCCTTACTCTTGGACAGGCGCAGTCGTACCCCGTGCCTCGCGGAAGAACAGAAA
 AGTTACAGAAATGAGAACCTGTCCGGTGAGCAGTGGGCGCACGGCGCCTTCTTGTCTTT

7261

LeuProIleAsnAlaLeuSerAsnSerLeuLeuArgHisAsnLeuValTyrSerThr
 CTGCCCCATCAATGCACATAAGCAACTCGTTGCTACGTACCCACCAATTTGGTGATTCACCC
 GACGGGTAGTTACGTGATTCTGTTGAGCAACGATGCAGTGGTGTAAACCATAGGTGG

7321

ThrSerArgSerAlaCysGlnArgGlnLysLysValThrPheAspArgLeuGlnValLeu

SUBSTITUTE SHEET

43 / 86

FIG. 18-21

7381 ACCTCAGCAGTGCTTGCCAAAGGCAGAAAGTCACATTTGACAGACTGCAAGTTCTG
TGGAGTGGTCAACGAACGGTTTCCGTCTTCTTTCAGTGTAACCTGTCTGACGTTCAAGAC

7441 AspSerHisTyrGlnAspValLeuLysGluValLysAlaAlaSerLysValLysAla
GACAGCCATTACCAAGGACGTACTCAAGGAGGTTAAAGCAGCGGTCAAAGTGAAGGCT
CTGTCCGGTAATGGTCCCTGCATGAGTTCCTCCTCCAATTTCGTCCGCCGAGTTTTCACCTTCCGA

(Phe)

7501 AsnLeuLeuSerValGluGluAlaCysSerLeuThrProHisSerAlaLysSerLys
AACTTGCTATCCGTAGAGGAAGCTTGACGCTGACGCCCCCACACTCAGCCAAATCCAAG
TTGAACGATAGGCATCTCCTTCGAACGTCGGACTGCGGGGTGTGAGTCGGTTTAGGTTT

7561 PheGlyTyrGlyAlaLysAspValArgCysHisAlaArgLysAlaValThrHisIleAsn
TTTGGTTATGGGGCAAAGACGTCCTGCTGCCATGCCAGAAAGGCCGTAAACCCACATCAAC
AAACCAATACCCCGTTTCTGCAGGCAACGGTACGGTCTTTCCGGCATTTGGGTGTAGTTG

7621 SerValTrpLysAspLeuLeuGluAspAsnValThrProIleaspThrIleMetAla
TCCGTGTGGAAAGACCTTCTGGAGAGACAAATGTAAACACCAATAGACACTACCATGGCT
AGGCACACCTTTCTGGAAGACCTTCTGTACATTGTGGTTATCTGTGTATGGTAGTACCGA

7681 LysAsnGluValPheCysValGlnProGluLysGlyGlyArgLysProAlaArgLeuIle
AAGAACGAGGTTTCTGCGTTTCAGCCTGAGAAAGGGGTGCGTAAGCCAGCTCGTCTCATC
TTCCTTGCTCCAAAGACGCAAGTCGGACTCTTCCCCCCCCAGCATTCGGTCCGAGCAGTAG

7741 ValPheProAspLeuGlyValArgValCysGluLysMetAlaLeuTyrAspValValThr
GTGTTCCCCGATCTGGGGGTGCGGTGCGAAAGATGGCTTTGTACGACGTGGTTACA
CACAAAGGGCTAGACCCGCGCACACGCTTTTCTACCGAAACATGCTGCACCAATGT

SUBSTITUTE SHEET

44/86

FIG. 18-22

7801 LysLeuProLeuAlaValMetGlySerSerTyrGlyPheGlnTyrSerProGlyGlnArg
AAGCTCCCTTGCCGTGATGGGAAGCTCCTACGGATTCCAATACTCACCAGGACAGCGG
TTCGAGGGGAACCGCACTACCCCTTCGAGGATGCCCTAAGGTTATGAGTGGTCTCTCGCGCC

7861 ValGluPheLeuValGlnAlaTrpLysSerLysLysThrProMetGlyPheSerTyrAsp
GTTGAATTCCCTCGTGCAAGCGTGAAGTCCAAGAAACCCCAATGGGGTTCTCGTATGAT
CAACTTAAGGAGCACGTTTCGCACCTTCAGGTTCTTTTGGGGTTACCCCAAGAGCATACTA

7921 ThrArgCysPheAspSerThrValThrGluSerAspIleArgThrGluGluAlaIleTyr
ACCCGCTGCTTTGACTCCACAGTCACTGAGAGCGACATCCGTACGGAGGAGGCAATCTAC
TGGCGACGAAACTGAGGTGTCAGTCACTCTCGCTGTAGGCATGCCTCCTCCGTTAGATG

7981 GlnCysCysAspLeuAspProGlnAlaArgValAlaIleLysSerLeuThrGluArgLeu
CAATGTTGTGACCTCGACCCCCAAGCCCGCGTGCCCATCAAGTCCCTCACCGAGAGGCTT
GTTACAACACTGGAGCTGGGGGTTTCGGGGCGCACCGGTAGTTCAGGGAGTGGCTCTCCGAA

8041 TyrValGlyGlyProLeuThrAsnSerArgGlyGluAsnCysGlyTyrArgArgCysArg
(Gly)
TATGTTGGGGCCCTCTTACCAATTCAAGGGGGAGAACTGCGGCTATCGCAGGTGCCGC
ATACAACCCCGGAGAAATGTTAAGTTCCCCCTCTTGACGCCCGATAGCGTCCACGGCG

8101 AlaSerGlyValLeuThrThrSerCysGlyAsnThrLeuThrCysTyrIleLysAlaArg
GCGAGCGCGGTACTGACAACCTAGCTGTGGTAACACCCCTCACTTGCTACATCAAGGCCCCG
CGCTCGCCGCATGACTGTTGATCGACACCATTTGTGGGAGTGAACCATGATGTTCCGGCC

SUBSTITUTE SHEET

45 / 86

FIG. 18-23

8161 AlaAlaCysArgAlaAlaGlyLeuGlnAspCysThrMetLeuValCysGlyAspAspLeu
 GCAGCCTGTCGAGCCGAGGGCTCCAGGACTGCACCATGCTCGTGTGTGGCGACGACTTA
 CGTCGGACAGCTCGGCTCCGAGGTCCCTGACGTGTACGTACGAGCACACACCGCTGCTGAAT

8221 ValValIleCysGluSerAlaGlyValGlnGluAspAlaAlaSerLeuArgAlaPheThr
 GTCGTTATCTGTGAAGCGGGGTCCAGGAGACGCGGCGAGCCTGAGAGCCTTCACG
 CAGCAATAGACACTTTCGCGCCCCCAGGTCTCTGCGCCGCTCGGACTCTCGGAAGTGC

8281 GluAlaMetThrArgTyrSerAlaProProGlyAspProProGlnProGluTyrAspLeu
 GAGGCTATGACCCAGGTACTCCGCCCCCTGGGACCCCCACAAACCAGAAATACGACTTG
 CTCCGATACTGTCATGAGCGGGGGACCCCTGGGGGTGTGTGCTTATGCTGAAC

8341 GluLeuIleThrSerCysSerSerAsnValSerValAlaHisAspGlyAlaGlyLysArg
 GAGCTCATAACATCATGCTCCTCCAACGTGTCAAGTCCGCCACGACGGCGCTGGAAAGAGG
 CTCGAGTATTGTAGTACGAGGAGGTTCACAGTCAGCGGGTGTGCTGCCGACCTTTCTCC

8401 ValTyrTyrLeuThrArgAspProThrThrProLeuAlaArgAlaAlaTrpGluThrAla
 GTCTACTACCTCACCCGTGACCCCTACAAACCCCTCGCGAGAGCTGCGTGGGAGACAGCA
 CAGATGATGGAGTGGCACTGGGATGTGTGGGGGAGCGCTCTCGACGACCCCTCTGTCGT

8461 ArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMetPheAlaProThrLeuTrp
 AGACACACTCCAGTCAATTCTCTGGCTAGGCAACATAATCATGTTGCCCCCACCACACTGTGG
 TCTGTGTGAGGTCAAGTAAAGGACCGATCCGTTGTATTAGTACAAACGGGGTGTGACACC

SUBSTITUTE SHEET

46 / 86

FIG. 18-24

8521 AlaArgMetIleLeuMetThrHisPhePheSerValLeuIleAlaArgAspGlnLeuGlu
GCGAGGATGATACTGATGACCCATTCTTTAGCGTCCTTATAGCCAGGACCGACTTGAA
CGCTCCTACTATGACTACTGGGTAAAGAAATCGCAGGAATATCGGTCCCTGGTCGAACCTT

8581 GlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIleGluProLeuAspLeuPro
CAGGCCCTCGATTGCGAGATCTACGGGCCCTGCTACTCCATAGAACCACTTGATCTACCT
GTCCGGGAGCTAACGCTCTAGATGCCCGGACGATGAGGTATCTTGGTGAAGTATGGA

8641 ProIleIleGlnArgLeuHisGlyLeuSerAlaPheSerLeuHisSerTyrSerProGly
CCAATCATTCAAAGACTCCATGGCCTCAGCGCATTTTCACTCCACAGTTACTCTCCAGGT
GGTTAGTAAAGTTTCTGAGGTACCGGAGTCGCGTAAAGTGAGGTGTCAATGAGAGGTCCA

8701 GluIleAsnArgValAlaAlaCysLeuArgLysLeuGlyValProProLeuArgAlaTrp
GAAATTAATAGGGTGGCCGATGCCCTCAGAAACTTGGGGTACCGCCCTTGCGAGCTTGG
CTTTAATTATCCCAACCGCGGTACGGAGTCTTTTGAAACCCCATGGCGGGAACGCTCGAACC

Gly

8761 ArgHisArgAlaArgSerValArgAlaArgLeuLeuAlaArgGlyGlyArgAlaIle
AGACACCGGGCCGGAGCGTCCGCGCTAGGCTTCTGGCCAGAGGAGCGAGGCTGCCATA
TCTGTGGCCCGGCCCTCGCAGGCGCGATCCGAAGACCGGTCTCTCCGTCCCGACGGTAT

8821 CysGlyLysTyrLeuPheAsnTrpAlaValArgThrLysLeuLysLeuThrProIleAla
TGTGGCAAGTACCTCTTCAACTGGGCGAGTAAGAACAAGCTCAAACTCACTCCCAATAGCG
ACACCGTTTCATGGAGAGAAGTTGACCCGTCATTCTTGTTCGAGTTTGAGTGAGGTTATCGC

SUBSTITUTE SHEET

47/86

FIG. 18-25

8881 AlaAlaGlyGlnLeuAspLeuSerGlyTrpPheThrAlaGlyTyrSerGlyGlyAspIle
GCCGCTGGCCAGCTGGACTTGTCCGGCTGGTTCACGGCTGGCTACAGCGGGGAGACATT
CGCGACCGGTCCGACCTGAACAGGCCGACCAAGTGCCGACCGATGTCGCCCCCTCTGTAA

8941 TyrHisSerValSerHisAlaArgProArgTrpIleTrpPheCysLeuLeuLeuAla
(Pro)
TATCACAGCGTGTCATGCCCGGCCCGCTGGATCTGGTTTTCCTACTCCTGCTTGCT
ATAGTGTCGACAGAGTACGGGGCGGACCTAGACCAAAACGGATGAGGACGAACGA

9001 AlaGlyValGlyIleTyrLeuLeuProAsnArgOP
GCAGGGGTAGGCATCTACCTCCTCCCAACCGATGAAGTTGGGTAAACACTCCGGCCT
CGTCCCCATCCGTAGATGGAGGAGGGTTGGCTACTTCCCAACCCCAATTGTGAGGCCGGA

SUBSTITUTE SHEET

48/86

:42.16.XT1
GGTAGGGTCAAGGCTGAAATCGACTGTCTGCTTCTTTGGAGAAAGTGGTG

:42.17.XT1
ATCCTGGGGGAGCGTGATTGTCTCAATGGTCTTCTTTGGAGAAAGTGGTG

:42.18.XT1
AGTCCTGCCCCGACGTTGAGTGCGGGAGACCTTCTTTGGAGAAAGTGGTG

:42.19.XT1
CACAAATCTGTAGATGCCTGGCTTCCCCCTCTTCTTTGGAGAAAGTGGTG

:42.20.XT1
GTCGAACATGCCGGAGGGGCGCTCCCCCGGCTTCTTTGGAGAAAGTGGTG

:42.21.LLA2C
GCCTGCGTCATAGCACTCACAGAGGACGGATTAGGCATAGGACCCGTGTC

:42.22.LLA2C
AGTCTCGGCGGGCGTGAGCTCATACCAAGCTTAGGCATAGGACCCGTGTC

:42.23.LLA2C
CGGGGTGTTTCATGTACGCTCGTAGCCTAACTTAGGCATAGGACCCGTGTC

:42.24.LLA2C
AAATTCAAGATGGTCCTGGCACACGGGAAGTTAGGCATAGGACCCGTGTC

:42.25.LLA2C
TATATGAGTGAGGCCTGTAAAGACGCCCTCTTAGGCATAGGACCCGTGTC

:42.26.LLA2C
ACTCTGCTTTGTCTGGGATAGAAAGTGGGCTTAGGCATAGGACCCGTGTC

:42.27.LLA2C
TTGGTACGCTACCAGGTAAGGAAGGTTCTCTTAGGCATAGGACCCGTGTC

:42.28.LLA2C
GGGAGGGGCTTGAGCCCTAGCGCACACGGTTTAGGCATAGGACCCGTGTC

:42.29.LLA2C
AATCAAACACTTCCACATCTGGTCCCACGATTAGGCATAGGACCCGTGTC

:42.30.LLA2C
GGGTGTTGGCCCATGGAGGGTGGGCTTGAGTTAGGCATAGGACCCGTGTC

:42.31.LLA2C
TTCATTCTGAACAGCGCCAGTCTGTATAGTTAGGCATAGGACCCGTGTC

FIG. 19-1

SUBSTITUTE SHEET

49/86

:42.XT1.1
TCCTCACAGGGGAGTGATTCATGGTGGAGTCTTCTTTGGAGAAAGTGGTG

:42.XT1.2
ATGGCTAGACGCTTTCTGCGTGAAGACAGTCTTCTTTGGAGAAAGTGGTG

:42.XT1.3
TCCTGGAGGCTGCACGACACTCATACTAACCTTCTTTGGAGAAAGTGGTG

:42.XT1.4
CGCAGACCACTATGGCTCTCCCGGGAGGGGCTTCTTTGGAGAAAGTGGTG

:42.XT1.5
TCGTCCTGGCAATTCCGGTGTACTCACCGGCTTCTTTGGAGAAAGTGGTG

:42.LLA2C.6
GCATTGAGCGGGTTGATCCAAGAAAGGACCTTAGGCATAGGACCCGTGTC

:42.LLA2C.7
AGCAGTCTTGCGGGGGCACGCCCAAATCTCTTAGGCATAGGACCCGTGTC

:42.LLA2C.8
ACAAGGCCTTTCGCGACCCAACTACTCGTTAGGCATAGGACCCGTGTC

:42.LLA2C.9
GGGGCACTCGCAAGCACCCCTATCAGGCAGTTTAGGCATAGGACCCGTGTC

:42.LLA2.10
CGTGCTCATGGTGCACGGTCTACGAGACCTTTAGGCATAGGACCCGTGTC

:42.LLA2C.11
GTTACGTTTGTTTTTTTTTTGAGGTTTAGGTTAGGCATAGGACCCGTGTC

:42.LLA2C.12
CGGGAACCTTGACGTCCTGTGGGCGACGGTTTAGGCATAGGACCCGTGTC

:42.LLA2C.13
CAAGTAAACTCCACCAACGATCTGACCGCCTTAGGCATAGGACCCGTGTC

:42.LLA2C.14
GCGCACACCCAATCTAGGGCCCCCTGCGCGGTTAGGCATAGGACCCGTGTC

:42.LLA2C.15
AGGTTGCGACCGCTCGGAAGTCTTCTCGTTTAGGCATAGGACCCGTGTC

FIG. 19-2

SUBSTITUTE SHEET

50/86

:42.32.XT1
ATGTTGGGATGGGGCACAGTGACGGAGCCCCCTTCTTTGGAGAAAGTGGTG

:42.33.XT1
ATCTCTCCGGTGGTGGACAGAGCAACCTCCCTTCTTTGGAGAAAGTGGTG

:42.34.XT1
ACTTCGAGGGGGATAGCCTTGCCGTAAAACTTCTTTGGAGAAAGTGGTG

:42.35.XT1
TGACAGAAGATGAGATGTCTCCCCCCTTGCTTCTTTGGAGAAAGTGGTG

:42.36.LLA2C
TTTGCGGCGAGTTCGTCGCACTTCTTCTTTTAGGCATAGGACCCGTGTC

:42.37.LLA2C
TAGGCCACGGCATTGATGCCCAATGCGACCTTAGGCATAGGACCCGTGTC

:42.38.LLA2C
GTCGGGATGACGGACACGTCAAGACCGCGGTTAGGCATAGGACCCGTGTC

:42.39.LLA2C
GCATCGGTTGCCACGACGACAACATCGCCGTTAGGCATAGGACCCGTGTC

:42.40.LLA2C
GAGTCGAAGTCGCCGGTATAGCCGGTCATGTTAGGCATAGGACCCGTGTC

:42.41.LLA2C
GTCTGGGTGACACACGTATTGCAGTCTATCTTAGGCATAGGACCCGTGTC


:42.42.LLA2C
ATGGTGAAGGTAGGGTCAAGGCTGAAATCGTTAGGCATAGGACCCGTGTC

:42.43.LLA2C
GAGACAGCATCCTGGGGGAGCGTGATTGTCTTAGGCATAGGACCCGTGTC

FIG. 19-3

SUBSTITUTE SHEET

51/86



5-1-1

C100

FIG. 20

SUBSTITUTE SHEET

FIG. 22 Translation of DNA 81

SerGlyLysProAlaIleIleProAspArgGluValLeuTyrArgGluPheAspGluMet
 1 GTCCGGGAAGCCGCAATCATACCTGACAGGAAGTCTCTACCGAGAGTTCGATGAGAT
 CAGGCCCTTCGGCCGTAGTAGTATGGACTGTCCCTTCAGGAGATGGCTCTCAAGCTACTCTA

GluGluCysSerGlnHisLeuProTyrIleGluGlnGlyMetMetLeuAlaGluGlnPhe
 61 GGAAGAGTGTCTCTCAGCACTTACCGTACATCGAGCAAGGATGATGTCGCCGAGCAGTT
 CCTTCTCAGAGAGTCGTGAATGGCATGTAGCTCGTTCCTACTACGAGCGGCTCGTCAA

LysGlnLysAlaLeuGlyLeuLeuGlnThrAlaSerArgGlnAlaGluValIleAlaPro
 121 CAAGCAGAAGGCCCTCGGCTCTCTGCAGACCGCTCCCGTCAGGCAGAGGTTATCGCCCC
 GTTCGTCTTCCGGGAGCCGGAGACGTCTGGCGCAGGCGAGTCCGTCTCCAATAGCGGGG

AlaValGlnThrAsnTrpGlnLysLeuGluThrPheTrpAlaLysHisMetTrpAsnPhe
 181 TGCTGTCCAGACCAACTGGCAAAACTCGAGACCTTCTGGCGGAAGCATATGTGGAACCTT
 ACGACAGTCTGGTTGACCGTTTTTGAGCTCTGGAAGACCCGCTTCGTATACACCTTGAA

IleSerGlyIleGlnTyrLeuAlaGlyLeuSerThrLeuProGlyAsnProAlaIleAla
 241 CATCAGTGGGATACAAATACTTGGCGGGCTTGTCAACGCTGCCCTGGTAACCCGCCATTGC
 GTAGTCACCCCTATGTTATGAACCGCCGAACAGTTGCGACGACCATTTGGGGCGGTAAACG

SerLeuMetAlaPheThrAlaAlaValThrSerProLeuThrThrSerGln
 301 TTCATTGATGGCTTTTACAGCTGCTGTCAACGACCCACTAACCACTAGCCAAA
 AAGTAACTACCGAAAATGTCGACGACAGTGGTCCGGTGATTGGTGATCGGTTT

52 / 86

SUBSTITUTE SHEET

FIG. 23 Translation of DNA 36

AspAlaHisPheLeuSerGlnThrLysGlnSerGlyGluAsnLeuProTyrLeuValAla
 1 GATGCCACTTCTATCCAGACAAAGCAGAGTGGGAGAACCTTCTTACCTGGTAGCG
 CTACGGGTGAAGATAGGTCTGTTTCGTCTCACCCCTCTTGAAGGAATGGACCATCGC
 TyrGlnAlaThrValCysAlaArgAlaGlnAlaProProSerTrpAspGlnMetTrp
 61 TACCAAGCCACCGTGTGCGCTAGGGCTCAAGCCCCCTCCCCCATCGTGGACCAGATGTGG
 ATGGTTCGGTGGCACACGCGATCCCGAGTTCGGGAGGGGTAGCACCTGGTCTACACC
 LysCysLeuIleArgLeuLysProThrLeuHisGlyProThrProLeuLeuTyrArgLeu
 121 AAGTGTTCGATTCGCTCAAGCCCAACCTCCATGGGCCAACACCCCTGCTATACAGACTG
 TTCACAACCTAAGCGGAGTTCGGGTGGAGGTACCCGGTTGTGGGACGATATGTCTGAC
 GlyAlaValGlnAsnGluIleThrLeuThrHisProValThrLysTyrIleMetThrCys
 181 GCGCGCTGTCAGAAATGAATCACCTGACGCACCCAGTCACCAATAACATCATGACATGC
 CCGGACAAAGTCTTACTTAGTGGGACTGCGTGGTCACTGTTATGTAGTACTGTACG
 MetSerAlaAspLeuGluValValThrSerThrTrpValLeuValGlyValLeuAla
 241 ATGTCGGCCGACCTGGAGGTCTGTCACGAGCACCTGGGTGCTCGTTGGCGGCTCCTGGCT
 TACAGCCGGCTGGACCTCCAGCAGTCTCGTGGACCCACGAGCAACCCGCCGAGGACCGA
 AlaLeuAlaAlaTyrCysLeuSerThrGlyCysValValIleValGlyArgValValLeu
 301 GCTTTGGCCGCGTATTGCTGTCAACAGGCTGCGTGTGTCATAGTGGCAGGTCGTCTTG
 CGAAACCGGCGCATAACGGACAGTTGTCCGACGCCACCATCATCACCCTCCAGCAGAAC
 -----Overlap with 81-----
 SerGlyLysProAlaIleIleProAspArgGluValLeuTyrArg
 361 TCCGGGAAGCCGGCAATCATACCTGACAGGGAAGTCCCTCTACCCGAG
 AGGCCCTTCGGCCGTTAGTATGACTGTCCCTTCAGGAGATGGCTC

SUBSTITUTE SHEET

54 / 86

FIG. 24 Translation of DNA 37b

LeuAlaAlaLysLeuValAlaLeuGlyIleAsnAlaValAlaTyrTyrArgGlyLeuAsp
 1 CTCGCCGCAAGCTGGTCGCAATGGGCATCAATGCCGTGGCCTACTACCGCGTCTTGAC
 GAGCGGCGTTTCGACCAGCGTAACCCGTAGTTACGGCACCGGATGATGGCGCCAGAACTG

ValSerValIleProThrSerGlyAspValValValAlaIleThrAspAlaLeuMetThr
 61 GTGTCCGTCATCCCGACCGAGCGGCGATGTTGTCGTCGTGGCAACCGATGCCCTCATGACC
 CACAGGCAGTAGGGCTGGTCGCCGCTACAACAGCAGCACCGTTGGCTACGGGAGTACTGG

GlyTyrThrGlyAspPheAspSerValIleAspTyrAsnThrCysValThrGlnThrVal
 121 GGCTATACCGCGACTTCGACTCGGTGATAGACTACAATACGTGTGTACCCAGACAGTC
 CCGATATGGCCGCTGAAGCTGAGCCACTATCTGATGTTATGCACACAGTGGGTCTGTCTAG

AspPheSerLeuAspProThrPheThrIleGluThrIleThrLeuProGlnAspAlaVal
 181 GATTTCAGCCTTGACCCCTACCTTCACCATTTGAGACAATCACGCTCCCCAGGATGCTGC
 CTAAAGTCGGAACTGGGATGGAAGTGTAACCTCTGTAGTGCAGGGGGTCTTACGACAG

clone 35-----Overlap with
 SerArgThrGlnArgArgGlyArgThr
 241 TCCCGCACTCAACGTCGGGCGAGGACTG
 AGGGCGTGAGTTGCAGCCCCCTCTGAC

SUBSTITUTE SHEET

55 / 86

FIG. 25A

1 2 3 4 5 6 7



FIG. 25B

+ + - + + + + - + - + - - +
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17



SUBSTITUTE SHEET

FIG. 26A

ALT(mU/ml)





FIG. 26B

ALT(mU/ml)



1 2 3 4 a b c

FIG. 26B'

57 / 86

Translation of DNA CA84a

GlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrpAsp
CGCAAGGTTGCAATTGCTCTATCTATCCCGGCATATAACGGGTACCGCATGGCATGGG
GCGTTCCAACGTTAACGAGATAGATAGGCGCGGTATATTGCCAGTGCGGTACCGTACCC

1

MetMetMetAsnTrpSerProThrAlaLeuValMetAlaGlnLeuLeuArgIlePro
ATATGATGATGAACCTGGTCCCCCTACGACGGCGTTGGTAATGGCTCAGCTGCCGATCC
TATACTACTACTTGACCAAGGGGATGCTGCCCAACCATTTACCGAGTCGACGAGGCCCTAGG

61

GlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyr
CACAAAGCCATCTTGACATGATCGCTGGTGCTCACTGGGAGTCTTGGCGGCATAGCGT
GTGTTCCGTTAGAACCTGTACTAGCGACCAACGAGTGACCCCTCAGGACCGCCCGTATCGCA

121

-----Overlap with CA59a-----

PheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGlyVal
ATTTCCTCCATGGTGGGAACCTGGGCGAAGTCCCTGGTAGTGTCTGTCTATTGCCGGCG
TAAAGAGTACCACCCCTTGACCCGCTTCCAGGACCATCAGCAGCAGATAAACGGCCGC

181

AspAlaGluThrHisValThrGly
TCGACGCGGAAACCCACGTACCCGGG
AGCTGCGCCCTTTGGGTGCAGTGCCCCC

241

FIG. 27

SUBSTITUTE SHEET

58 / 86

1 AlaTyrMetSerLysAlaHisGlyIleAspProAsnIleArgThrGlyValArgThrIle
 GGCTTACATGTCCTCAAGGCTCATGGGATCGATCCTAACATCAGGACCGGGTGAGAACAAAT
 CCGAATGTACAGGTTCCGAGTACCCTAGCTAGGATTGTAGTCTGCCCCCCTCTTTGTTA
 61 ThrThrGlySerProIleThrTyrSerThrTyrGlyLysPheLeuAlaAspGlyGlyCys
 TACCACCTGGCAGCCCCATCAGTACTCCACCTACGGCAAGTTCTTCCGACGCGGGTG
 ATGGTGACCGTCGGGTAGTGCATGAGGTGATGCCGTTCAAGGAACGGCTGCCGCCAC
 121 SerGlyGlyAlaTyrAspIleIleIleCysAspGluCysHisSerThrAspAlaThrSer
 CTCGGGGGCGCTTATGACATAATAATTGTGACGAGTGCCACTCCACGGATGCCACATC
 GAGCCCCCGCGAATACTGTATTATTAAACACTGCTCACGGTGAGGTGCCGTACGGTGTAG
 181 IleLeuGlyIleGlyThrValLeuAspGlnAlaGluThrAlaGlyAlaArgLeuValVal
 CATCTTGGGCATCGGCACCTGTCTTGACCAAGCAGAGACTGCGGGGCGAGACTGGTTGT
 GTAGAACCCGTAGCCGTGACAGGAACCTGTTCTGCTCTGACGCCCCCGCTCTGACCAACA
 241 LeuAlaThrAlaThrProGlySerValThrValProHisProAsnIleGluGluVal
 GCTCGCCACCGCCACCCCTCCGGGCTCCGTCACTGTGCCCCCATCCCAACATCGAGGAGGT
 CGAGCGGTGGCGGTGGAGGCCCGAGGCAGTGACACGGGTAGGGTTGTAGCTCCTCCA
 301 AlaLeuSerThrThrGlyGluIleProPheTyrGlyLysAlaIleProLeuGluValIle
 TGCTCTGTCCACCAACCGGAGAGATCCCTTTTACGGCAAGGCTATCCCCCTCGAAGTAAT
 ACGAGACAGGTGTGGCCTCTCTAGGGAAAAATGCCGTTCGGATAGGGGAGCTTCATTA
 -----Overlap with 37b-----
 361 LysGlyGlyArgHisLeuIlePheCysHisSerLysLysLysCysAspGluLeuAlaAla
 CAAGGGGGGAGACATCTCATCTTCTGTCTCATTCAAAGAGAAGTGCAGCAACTCGCCGC
 GTTCCCCCTCTGTAGAGTAGAAGACAGTAAGTTTCTTCTCACGCTGCTTGAGCGCGC
 421 LysLeuValAlaLeuGlyIleAsnAlaValAlaTyrTyrArgGlyLeuAspValSerVal
 AAAGCTGGTCGCATGGGCATCAATGCCGTGGCCTACTACCGCGGTCTTGACGTGTCCTG
 TTTCGACACGCGTAACCCGTAGTTACGGCACCGGATGATGGCGCCAGAACTGCACAGGCA

 481 IleProThr
 CATCCCGACCCAG
 GTAGGGCTGGTC

Translation of DNA 40b FIG. 28

SUBSTITUTE SHEET

59/86

1 2 3



FIG. 29

SUBSTITUTE SHEET

60 / 86

HCV cDNA OF CLONE 40a

1 GluPheGlyAlaIleProLeuGluValIleLysGlyGlyArgHisLeuIlePheCysHis
GAATTCGGGGCTATCCCCCTCGAAGTAATCAAGGGGGAGACATCTCATCTTCTGTGTCAT
CTTAAGCCCCCGATAGGGGAGCTTCAATTAGTTCCCCCTCTGTAGAGTAGAAGACAGTA

61 TCAAAGAAGAAGTGCGACGAACTCGCCGAAAGCTGTCGCATTGGCATCAATGCCGTG
AGTTTCTTCTTCAAGCTGCTTGAGCGGCGTTTCGACCAGCGTAACCCGTAGTTACGGCAC

121 GCCTACTACCGCGGTCTTGACGTGTCCGTCAATCCCGACCAAGCGGTGATGTTGTCGTGTCG
CGGATGATGGCGCCAGAACTGCACAGGCAGTAGGGCTGGTCGCCACTACAACAGCAGCAC

181 GCAACCGATGCCCTCATGACCGGCTATACCGCGACTTCGACTCGGTGATAGACTGCAAT
CGTTGGCTACGGGAGTACTGGCCGATATGCGCCGCTGAAGCTGAGCCCACTATCTGACGTTA

241 ACGTGTGTCAACCCAGACAGTCGATTTTCAGCCCTTGACCCCTACCTTCACTATTGAGACAATC
TGCACACAGTGGGTCTGTCAGCTAAAGTCGGAACTGGGATGGAGTGATAACTCTGTAG


301 ACGCTCCCCCAAGATGCTCCGAATTC
TCCGAGGGGGTTCTACGAGGCTTAAG

FIG. 32

SUBSTITUTE SHEET

61/86

FIG. 33



SUBSTITUTE SHEET

62 / 86

FIG. 34-1 Translation of DNA 35

SerIleGluThrIleThrLeuProGlnAspAlaValSerArgThrGlnArgArgGlyArg
 1 TCCATTGAGACAAATCACGCTCCCCAGGATGCTGTCTCCGCACTCAACGTCTGGGGCAGG
 AGGTAACCTCTGTAGTGCAGGGGTCCTACGACAGAGGGCGTGAGTTGCAGCCCCGTCC

ThrGlyArgGlyLysProGlyIleTyrArgPheValAlaProGlyGluArgProSerGly
 61 ACTGGCAGGGGAAGCCAGGCATCTACAGATTGTGTGCCACCGGGGAGCGCCCTCCGGC
 TGACCGTCCCCCTTCGGTCCGTAGATGTCTAAACACCGTGGCCCCCTCGCGGGAGGCCG

MetPheAspSerSerValLeuCysGluCysTyrAspAlaGlyCysAlaTrpTyrGluLeu
 121 ATGTTCCGACTCGTCCGTCCTCTGTGAGTGCTATGACGCAGGCTGTGCTTGGTATGAGCTC
 TACAAGCTGAGCAGGCAGGAGACACTCAGGATACGCTCCGACACGACCACTACTCGAG

ThrProAlaGluThrThrValArgLeuArgAlaTyrMetAsnThrProGlyLeuProVal
 181 ACGCCCGCGAGACTACAGTTAGGCTACGAGCGTACATGAACACCCCCGGGCTTCCCGTG
 TCGGGGGGGCTCTGATGTCAATCCGATGCTCGCATGTACTTGTGGGGCCCCGAAGGGCAC

SUBSTITUTE SHEET

63 / 86

FIG. 34-2

CysGlnAspHisLeuGluPheTrpGluGlyValPheThrGlyLeuThrHisIleAspAla
241 TGCCAGGACCATCTTGAAATTTGGAGGGCGTCTTACAGGCCCTCACTCATATAGATGCC
ACGGTCCTGGTAGAACTTAAACCCCTCCCGCAGAAATGTCCGGAGTGAGTATATCTACGG

HisPheLeuSerGlnThrLysGlnSerGlyGluAsnLeuProTyrLeuValAlaTyrGln
301 CACTTCTATCCAGACAAAGCAGAGTGGGAGAACCTTCCTTACCTGGTAGCGTACCAA
GTGAAAGATAGGGTCTGTTCGTCTCACCCCTCTTGGAAGGAATGGACCATCGCATGGTT

-----Overlap with 36-----

AlaThrValCysAlaArgAlaGlnAlaProProSerTrpAspGlnMetTrpLysCys
361 GCCACCGTGTGGCTAGGGCTCAAGCCCTCCCCATCGTGGACCAAGATGTGGAAGTGT
CGGTGGCACACGCCGATCCCGAGTTCGGGGAGGGGTAGCACCCCTGGTCTACACCTTCACA

LeuIleArgLeuLysProThrLeuHisGlyProThrProLeuLeuTyrArgLeuGlyAla
421 TTGATTCGCCCTCAAGCCCAACCTCCATGGGCCAACACCCCTGCTATACAGACTGGCGCT
AACTAAGCGGAGTTCGGGTGGGAGGTACCCGGTTGTGGGACGATATGTCTGACCCCGCA


SUBSTITUTE SHEET

NOT FURNISHED UPON FILING

NOT FURNISHED UPON FILING

PCR/HCV ENV Region

FIG. 37




100 bp

— = probes CA216a, CA84a
 - - = primers for probes

67 / 86

FIG. 38



SUBSTITUTE SHEET

68/86

FIG. 39-1

1. human 27 2. HCV 1 3. human 23

1 CGGCTTCGCCGACCTCATGGGTACATCCGCTCGTCGGCGCCTCCTCTTGGGGCGCTGCCAGGGCCCTGGC

1 CGGCTTCGCCGACCTCATGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGGCGCTGCCAGGGCCCTGGC

1 CGGCTTCGCCGACCTCATGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGGCGCTGCCAGGGCCCTGGC

73 GCATGGCGTCCGGTTCGGAAGACGGCGTGAACTATGCAACAGGGAACCTTCCTGGTTGCTCTTCTCTAT

73 GCATGGCGTCCGGTTCGGAAGACGGCGTGAACTATGCAACAGGGAACCTTCCTGGTTGCTCTTCTCTAT

73 GCACGGCGTCCGGTTTGGGAAGACGGCGTGAACTATGCAACAGGGAACCTTCCTGGTTGCTCTTCTCTAT

145 CTTCCTTCTGGCtCTGCTCTTGCCTGACCGTGCCCGCaTCGGCCTACCAAGTaCGCAACTCCTCGGGCaT

145 CTTCCTTCTGGCCCTGCTCTTGCtTGACTGTGCCCGCTTCGGCCTACCAAGTGCGCAACTCCACGGGGCT

145 CTTCCTTCTGGCCCTaCTCTCTTGCTGCTGACCGTGCCCGCTTCaGCCCTACCAAGTGCGCAACTCTACGGGGCT

217 TTACCAtGTCACCAATGATTGCCCTAAATCGAGTATTGTGTACGAGaCGGCCGAcCaCCATCCTaCActCTCC

217 TTACCAcGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGCGGCCGATGCCATCCTGCACaCTCC

217 TTACCAtGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGCGGCCGATGCCATCCTGCACgCTCC

289 GGGGTGtGTCCCTTGCGTTCGcGAGGGtAACGCCCTCGAaaTGTTGGTGcCGgTagCCCCcACaGTGGCCAC

289 GGGGTGcGTCCCTTGCGTTCGtGAGGGcAACGCCCTCGAGgTGTGGGTGGCGaTGACCCCTACGGTGGCCAC

289 GGGGTGtGTCCCTTGCGTTCGcGAGGataACGtCTCGAGaTGTTGGGTGGCGgTGACCCcACGGTGGCCAC

SUBSTITUTE SHEET

70/86

793 tGcCaggACcaccGcagGcgCTcaccAGtTtTtTCagcCCAGGCGCCAAGCAGgAtaTCCAGCTGATCAACAC
* * * * *
793 CGgCCaCACtgtGtCTGGAtTTGtTAGcCTCcTCgCACAGGCGCCAAGCAGAAcGTCCAGCTGATCAACAC
* * * * *
793 CGcCCgCAgcacGgCTGGAgTTGcTAGtCTCtTcCaCACcAGGCGCTAgGCAGAAcATCCAGCTGATCAACAC

865 CAACGGCAGTTGGCACaTCAATcGCACGGCCtTGAACtGtAATGcgAGCCTCgACACtGGCTGGgTaGCgGG
* * * * *
865 CAACGGCAGTTGGCACcCTCAATAGCACGGCCcTGAACtGCAATGAtAGCCTCAACACCGGCTGGTTgGCaGG
* * * * *
865 CAACGGCAGTTGGCACaTCAATAGtACGGCCtTGAACtGCAATGAcAGCCTtAcCACCGGCTGGTTaGCgGG

937 GCTcTTCTATtACCACAAaTTCAACTCTTCAGGCTGcCCcGAGAGGaTgGCCAGCTGtaGgCCCCTTgCCGA
* * * * *
937 GCTTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCCcTGAGAGGcTaGCCAGCTGCCGACCCCTTACCga
* * * * *
937 GCTTTTCTATCACCAtAAaTTCAACTCTTCAGGCTGTCCcGAGAGGtTgGCCAGCTGCCGACCCCTcACCCGA

1009 TTTcGACCAGG
* * * * *
1009 TTTTGACCAGG
* * * * *
1009 TTTTGCCcAGG

FIG. 39-3

SUBSTITUTE SHEET

FIG. 40

1 GFADLMGYIPLVGAPLGGAAARALAHGVRVLEEDGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSSGi

1 GFADLMGYIPLVGAPLGGAAARALAHGVRVLEEDGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSTGL

1 GFADLMGYIPLVGAPLGGIARALAHGVRVLEEDGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSTGL
73 YHVTNDCPNSSIVYEtADtILHSPGCVPCVREGNASKCWVpvaPTVATRDNGLPATQLRRHIDLIVGSATLC

73 YHVTNDCPNSSIVYEADAAILHtPGCVPCVREGNASRCWVAmTPTVATRDNGLPATQLRRHIDLIVGSATLC

73 YHVTNDCPNSSIVYEADAAILHaPGCVPCVREDnVSRCWVAVTPTVATkDGKLPtTQLRRHIDLIVGSATLC⁷
145 SALYVGDLGGSVFLVGQLFTFSRRHWTtQdCNCSiYPGHITGHRMAWDMMNWSPtaALVMAQLLRIPQAI⁶

145 SALYVGDLGGSVFLVGQLFTFSRRHWTtQdCNCSiYPGHITGHRMAWDMMNWSPtTaALVMAQLLRIPQAI

145 SALYVGDLGGSiFLVGQLFTFSRRHWTtQdCNCSiYPGHITGHRMAWDMMNWSPtaALVVAQLLRIPQAI
217 LDMIAGAHWGVLAGIAyFSMVGNWAKVLVLLLFAGVDAtTyttGGnAarTtqaltSffsPGAKQdiQLINT

217 LDMIAGAHWGVLAGIAyFSMVGNWAKVLVLLLFAGVDAETHVTGGSAGhTvsGfvSLlapGAKQNVQLINT

217 LDMIAGAHWGVLAGmAYFSMVGNWAKVLVLLLFAGVDAETHrTGGSAarstaGvaSLftPGARoniQLINT
289 NGSWHInrTALNCNaSLdTGwVAGLFyYHKFNSSGCPERMASCRPLaDFDQ

289 NGSWHInSTALNCNDSLnTGwLAGLFyYHKFNSSGCPERLAsCRPLTDfDQ

289 NGSWHInSTALNCNDSLttGwLAGLFyYHKFNSSGCPERLAsCRPLTDfAQ

1. human 27
2. HCV 1
3. human 23

SUBSTITUTE SHEET

72/ 86

1 2 3 4




FIG. 41

SUBSTITUTE SHEET




FIG. 43




FIG. 44A




FIG. 44B

74 / 86

FIG. 46-1

Human 23

GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyArgAla
 1 GGCTTCGCCGACCTCATGGGTACATACCGCTCGTCGGCCCTCTTGGAGGCCGTGCC
 ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
 61 AGGGCCCTGGCGCACGGCTCCGGTTTGGAGACGGCGTGAACCTATGCAACAGGGAAC
 CG A
 LeuProGlyCysSerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValPro
 121 CTTCCCTGGTTGCTCCTTTCTATCTTCTCTTCTGGCCCTACTCTCTTGCCCTGACCGTGCCC
 GA
 AlaSerAlaTyrGlnValArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysPro
 181 GCTTCAGCCTACCAAGTGGCAACTCTACGGGGCTTTACCATGTCAACCAATGATTGCCCT
 AsnSerSerIleValTyrGluAlaAlaAspAlaIleLeuHisAlaProGlyCysValPro
 241 AACTCGAGTATTGTGTACGAGGCGCGGATGCCATCCTGCACGCTCCGGGTGTGTCCCT
 T
 CysValArgGluAspAsnValSerArgCysTrpValAlaValThrProThrValAlaThr
 301 TCGGTTCCGCGAGGATAACGTCTCGAGATGTTGGGTGGGTGACCCACCGGTGCCACC
 G
 LysAspGlyLysLeuProThrThrGlnLeuArgArgHisIleAspLeuLeuValGlySer
 361 AAGGACGGCAAACTCCCCACAACGACGCTTCGACGTACATCGATCTGCTTGTCTGGGAGC
 C
 AlaThrLeuCysSerAlaLeuTyrValGlyAspLeuCysGlySerIlePheLeuValGly
 421 GCCACCTCTGCTCGCCCTCTACGTGGGGACCTTTGCGGGTCCATCTTCTTGTCTCGGT
 T
 GlnLeuPheThrPheSerProArgArgHisTrpThrThrGlnAspCysAsnCysSerIle
 481 CAACTGTTTACCTTCTCTCCAGGCGCCACTGGACGACGACGAGGACTGCAACTGTTCTATC
 C

SUBSTITUTE SHEET

75/ 86

FIG. 46-2

TyrProGlyHisIleThrGlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerPro
 541 TATCCCGCCATATAACGGTCACCGCATGGCATGATGATGATGAAGTGGTCCCT
 ThrAlaAlaLeuValValAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIle
 601 ACGCGGCATTGGTAGTCTAGCTCAGCTGCTCCGATCCCAAGCCATCTTGGACATGATC
 AlaGlyAlaHisTrpGlyValLeuAlaGlyMetAlaTyrPheSerMetValGlyAsnTrp
 661 GCTGGTCTCACTGGGAGTCTCTGGCGGCATGGCGTATTCTCTCCATGGTGGGAACCTGG
 AlaLysValLeuValValLeuLeuLeuPheAlaGlyValAspAlaGluThrHisArgThr
 721 GCGAAGGTCTCTGGTAGTGCTGCTTCTATTGCGCGTCTCGACGCGGAACCCACCGTACC
 GlyGlySerAlaAlaArgSerThrAlaGlyValAlaSerLeuPheThrProGlyAlaArg
 781 GGGGAAGTGCCGCCGACGACGGCTGGAGTGTCTCTCTTCACACACGCGCTAGG
 GlnAsnIleGlnLeuIleAsnThrAsnGlySerTrpHisIleAsnSerThrAlaLeuAsn
 841 CAGAACATCCAGCTGATCAACACCAACGGCAGTTGGCACATCAATAGTACGGCCTTGAAC
 CysAsnAspSerLeuThrThrGlyTrpLeuAlaGlyLeuPheTyrHisHisLysPheAsn
 901 TGCAATGACAGCCTTACCACCGGCTGGTGTAGCGGGCTTTTCTATCACCATAAATTCAAC
 SerSerGlyCysProGluArgLeuAlaSerCysArgProLeuThrAspPheAlaGln
 961 TCTTCAGGCTGTCCGAGAGGTGGCCAGCTGCCGACCCCTCACCGATTGTGCCCAGG

SUBSTITUTE SHEET

76/86

FIG. 47-1

Human 27

1 GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAla
 GGCTTCGCCGACCTCATGGGTACATCCGCTCGTCGGCTCCTCTTGGGGCGCTGCC
 61 ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
 AGGCCCTGGCGCATGGCGTCCGGTTCTGGAAGACGGCGTGAACATATGCAACAGGAAC
 121 LeuProGlyCysSerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValPro
 CTCCTGGTTGCTCTTTCTCTATCTTCCTCTGGCTCTGCTCTCTTGCCCTGACCCGTGCC
 181 AlaSerAlaTyrGlnValArgAsnSerSerGlyIleTyrHisValThrAsnAspCysPro
 GCATCGGCCTACCAAGTACGCAACTCCTCGGGCATTTACCATGTCAACCAATGATTGCCCT
 241 AsnSerSerIleValTyrGluThrAlaAspThrIleLeuHisSerProGlyCysValPro
 AATTCGAGTATTGTGTACGAGACGGCCGACACCATCTACACTCTCCGGGTGTGTCCCT
 301 CysValArgGluGlyAsnAlaSerLysCysTrpValProValAlaProThrValAlaThr
 TGCGTTCGGAGGGTAACGCCCTCGAAATGTTGGTGGCCGTAGCCCCCACAGTGGCCACC
 361 ArgAspGlyAsnLeuProAlaThrGlnLeuArgArgHisIleAspLeuLeuValGlySer
 AGGGACGGCAACCTCCCCGCAACGCAGCTTCGACGTCACATCGATCTGCTTGTGGGAGT
 421 AlaThrLeuCysSerAlaLeuTyrValGlyAspLeuCysGlySerValPheLeuValGly
 GCCACCCCTTGCTCGGCCCTCTATGTGGGGACTGTGCGGGTCTGTCTTTCTTGTTCGGT
 481 GlnLeuPheThrPheSerProArgArgHisTrpThrThrGlnAspCysAsnCysSerIle
 CAACTGTTCACTTTCTCCCCCAGGGCCACTGGACAAACGCAAGATTGCAACTGCTCTATC

SUBSTITUTE SHEET

FIG. 47-2

541 TyrProGlyHisIleThrGlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerPro
TACCCGGCCATATAACGGGACACCGCATGGCATGGGATATGATGATGAAGTGTCCCT

601 ThrAlaAlaLeuValMetAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIle
ACAGCAGCGCTGGTAATGGCTCAGCTGCTCAGGATCCGCAAGCCATCTTGGACATGATC
G

661 AlaGlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrp
GCTGGTGCTCACTGGGAGTCCTAGCGGCATAGCGTATTTCTCCATGTTGGGGAACCTGG

721 AlaLysValLeuValLeuLeuLeuPheAlaGlyValAspAlaThrThrTyrThrThr
GCGAAGTGCTGCTGGTGCTGTTGCTGTTGCGCGCTCGATGCGACAACCTATACCACC

781 GlyGlyAsnAlaAlaArgThrThrGlnAlaLeuThrSerPhePheSerProGlyAlaLys
GGGGGAATGCTGCCAGGACCACGCGCGCTCACCAGTTTTCAGCCCCAGGCGCCAAG

841 GlnAspIleGlnLeuIleAsnThrAsnGlySerTrpHisIleAsnArgThrAlaLeuAsn
CAGGATATCCAGCTGATCAACACCAACGGCAGTTGGCACATCAATCGCACGGCCTTGAAC
G T

901 CysAsnAlaSerLeuAspThrGlyTrpValAlaGlyLeuPheTyrTyrHisLysPheAsn
TGTAATGCGAGCCTCGACACTGGCTGGGTAGCGGGGCTCTTCTATTACCACAAATTCAAC
T G

961 SerSerGlyCysProGluArgMetAlaSerCysArgProLeuAlaAspPheAspGln
TCTTCAGGCTGCCCCGAGAGGATGGCCAGCTGTAGGCCCTTGCCGATTTCGACCAGG
C

FIG. 48




FIG. 49-1

CLUSTERED PAIR-WISE 'REGION' ALIGNMENT
in 'identity (no translation)' alphabet of:

1. ssThorn#8.r (1-587)
2. ssEC1#2.r (1-587)
3. ssHCT18#7.r (1-587)
4. env1.hcv (1-1657)

$$\text{GA} \parallel \text{GA} \parallel \text{GA}.$$

289 gggtaggggatggctcctgtctcccgtagctgggcccacagacccccggcgtaGg

3 ATTCGCAATTGGGTAAGGTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC
3 ATTCGCAATTGGGTAAGGTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC
3 ATTCGCAATTGGGTAAGGTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTATATACCGCTC
361 tgcgCGCAATTGGGTAAGGTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC

75 GTCGGCGCCCCCTCTTGGGGCGCTGCCAGGGCCCCCTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAAC
75 GTCGGCGCCCCCTCTTGGAGCGCTGCCAGGGCCCCCTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAAC
75 GTCGGCGCCCCCTCTTGGAGCGCTGCCAGGGCCCCCTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAAC
433 GTCGGCGCCCCCTCTTGGAGCGCTGCCAGGGCCCCCTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAAC

80/86

FIG. 49-2

147 TATGCAACAGGAACCTTCCGTGCTCTTCTcTcTCTTCCCTTCTGGCCcTGCTCTCTTGTcTGACcGTG
147 TATGCAACAGGAACCTTCCGTGCTCTTCTcTcTCTTCCCTTCTGGCCcTGCTCTCTTGTcTGACcGTG
147 TATGC CAGGGAACCTTCCGTGCTCTTCTcTcTCTTCCCTTCTGGCCcTGCTCTCTTGTcTGACcGTG
505 TATGCAACAGGAACCTTCCGTGCTCTTCTcTcTCTTCCCTTCTGGCCcTGCTCTCTTGTcTGACcGTG
219 CCCGCTTCAGCCTACCAAGTGGCAACTCcaCGGGGCTTTACCATGTCAACAAcGATTGCCCCcAACTCGAGt
219 CCCGCTTCAGCCTACCAAGTGGCAACTCCTcCGGGGCTTTACCATGTCAACAAcGATTGCCCCcAACTCGAGc
219 CCCGCTTCAGCCcACCAAGTGGCAACTCCACGGGGCTTTACCATGTCAACAAcGATTGCCCCcAACTCGAGT
577 CCCGCTTCgGCCTACCAAGTGGCAACTCCACGGGGCTTTACCAcGTCAACAAcGATTGCCCCcAACTCGAGT
291 ATTGTGTACGAGGGGGCCGATGCTcATCCTGCACgCTCCGGGGTGTGTCCCTTGCGTTcGCGAGGtAACGcc
291 ATTGTGTACGAGGGGGCCGATGCCATCCTGCACACTCCGGGGTGTGTCCCTTGCGTTcACGAGGGCAACGTC
291 ATTGTaTACGAaGCGGCCGACGCCATCCTGCACACTCCGGGGTGTGTCCCTTGCGTTcACGAGGGCAACGTC
649 ATTGTgTACGAgGCGGCCGAtGCCATCCTGCACACTCCGGGGTGTGTCCCTTGCGTTcGtGAGGGCAACGcc
363 TCGAGGTGTGGGTGGCGATGACCCcCACGGTGGCCcCCAGGGgCCAGGAcCGGCAgACTCCCCACAACGAGCTgCGA
363 TCGAGGTGTGGGTGGCGATGACCCcCACGGTGGCCcCCAGGGgCCGGCAAACTCCCCACAACGAGCTTCGA
363 TCGAGGTGTGGGTGGCGgTGACCCcCACGGTGGCCcCCAGGATGGCAAACTCCCCACAACGAGCTTCGA
721 TCGAGGTGTGGGTGGCGaTGACCCcCACGGTGGCCcCCAGGATGGCAAACTCCCCCgCgACGAGCTTCGA

435 CGTCACATCGATCTGCTTGTTCGGGAGCGCCACCCCTCTGCTCGGCCCTCTACGTGGGGGACCTGTGCGGGTCC
435 CGTCACATCGATCTGCTTGTTCGGGAGCGCCACCCCTCTGCTCGGCCCTCTACGTGGGGGACCTGTGCGGGTCT
435 CGTCACATCGATCTGCTTGTTCGGGAGCGCCACCCCTCTGCTCGGCCCTCTATGTGGGGGACCTGTGCGGGTCT
793 CGTCACATCGATCTGCTTGTTCGGGAGCGCCACCCCTCTGCTCGGCCCTCTACGTGGGGGACCTaTGCGGGTCT

507 aTCTTtCTTGTCGGTCAACTGTTcACCTTCTCTCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCT 81/86
507 GTCTTcCTTGTCGGTCAACTGTTTACCTTCTCTCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCT
507 GTCTTCTTGTCGGCCAACGTGTTTACCTTCTCTCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCT
865 GTCTTCTTGTCGGCCAACGTGTTcACCTTCTCTCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCT

579 ATCGAATTC
579 ATCGAATTC
579 ATCGAATTC
937 ATCTAtccc

FIG. 49-3

EC10
HCV1

10 20 30 40
GAATTCGGACGACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATAT
X::

550 560 570 580 590 600
CTCTCCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATAT

50 60 70 80 A 90 100
AACAGGTCACCGCATGGCATGGGATATGATGATGAAGTGGTCCCCTACGACGGCGTTAGT
::: ::: ::
AACGGGTCACCGCATGGCATGGGATATGATGATGAAGTGGTCCCCTACGACGGCGTTGGT
610 620 630 640 650 660

110 120 130 140 150 160
GGTAGCTCAGCTGCTCCGGATCCCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTG
: ::: ::
AATGGCTCAGCTGCTCCGGATCCCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTG
670 680 690 700 710 720

170 180 190 200 210 220
GGGAGTCCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAACTGGGCGAAGGTCTTGGC
::: ::
GGGAGTCCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAACTGGGCGAAGGTCTTGGT
730 740 750 760 770 780

230 240 250 260 270 280
AGTGCTGCTGCTATTTGCCGGCGTCGACGCGGAAACCCACGTCACTGGGGGGATCGCCGC
::: :::: : ::
AGTGCTGCTGCTATTTGCCGGCGTCGACGCGGAAACCCACGTCAACGGGGGAAGTGCCGG
790 800 810 820 830 840

290 300 310 320 330 340
CAAACTACGGCTAGCCTTACTGGTCTCTTCAATTTAGGTGCCAAGCAGAACATCCAGCT
: : :: : :: : ::
CCACACTGTGTCTGGATTGTGTAGCCTCCTCGCACCAGGCGCCAAGCAGAACGTCCAGCT
850 860 870 880 890 900

350 360 370 380 390 400
GATCAACACCAACGGCAGTTGGCACATCAACAGGACGGCCTTGAAGTGAATGATAGCCT
::::::::::::::::::::::::: :::: : : : : : : : : : : : : : : : : :
GATCAACACCAACGGCAGTTGGCACCTCAATAGCACGGCCTTGAAGTGAATGATAGCCT
910 920 930 940 950 960

410 420
CAACACCGGCTGGAATTC
::::::::::::X
CAACACCGGCTGGTTGGCAGGGCTTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCC
970 980 990 1000 1010 1020

FIG. 50-1

SUBSTITUTE SHEET

NOT FURNISHED UPON FILING

84 / 86

AA #117-308 (putative envelope region)

FIG. 51

- | | |
|-----------------------|--------------------|
| 1) HCT #18 (USA) | 3 clones sequenced |
| 2) JH23 (USA) | ? |
| 3) JH 27 (USA) | ? |
| 4) PBL-Th (USA) | 2 clones sequenced |
| 5) EC1 (Italy) | 3 clones sequenced |
| 6) HCV-1 (chimpanzee) | multiple |

C/M ← T → S

- 1) (P)
- 2)
- 3)
- 4)
- 5)
- 6) RNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARALAHGVRVLEDGVNYATGNL

- 1) H
- 2)
- 3) S T T
- 4) L
- 5) (F) S
- 6) PGCSFSIFLLALLSCLTVPASAYQVRNSTGLYHVTNDCPNSSIVYEAADAILH

- 1) Y (H) V V T
- 2) A D V V K T
- 3) S PVA N
- 4) A A R T
- 5) H V T
- 6) TPGCVPCVREGNASRCWVAMTPTVATRDGKLPATQLRRHIDLLVGSATLCS

- 1)
- 2) I D
- 3) D
- 4)
- 5) I
- 6) ALYVGDLGSGVFLVGQLFTFSPRRHWTTQGCNCSI

SUMMARY: "S" AA117-308 (93%)

HCT#18, PBL-Th, EC1(Italy) have 97% homology with HCV-1

JH23 and JH 27 have 96% and 95% homology with HCV-1, respectively

SUBSTITUTE SHEET

85 / 86

AA#300-438 (C-terminal region of the putative envelope region and amino ~1/3 of NSI)

- 1) JH23 ?
 2) JH27 ?
 3) Japanese Isolate (T. Miyamura) ?
 4) EC10 (Italy) 2 clones sequenced
 (one nt difference, which did not
 result in an amino acid change)
 multiple

5) HCV-1 (chimpanzee)

S ← → NS I

1) D

A V

2) D

A

3)

V S

V M V

4)

5) TTQGCNCSIYPGHITGHRMAWDMMMWNWSPPTALVMAQLLRIPQAILDMIAGA

1) M

R

A R S T A V A

2)

T Y T

N A R T Q A L T F

3) L Y

I M

G H R

V Q V T T L T

4)

A

I A K T A S L T A

5) HWGVLAGIAYFSMVGNWAKVLVLLLFAGVDAETHVTGGSAGHTVSGFVSL

1) F S R I I T V

2) F T D I I R A D

3) F R S K I V I R Q F

4) F N L I I R N

5) L A P G A K Q N V Q L I N T N G S W H L N S T A L N C N D S L N T G W L

SUMMARY: NS 1 AA 330-660

| "Isolate" | %Homology (AA330-438) | %Homology (AA383-405) |
|--------------|-----------------------|-----------------------|
| JH23 | 83 | 57 |
| JH27 | 80 | 39 |
| Japanese | 73 | 48 |
| EC10 (Italy) | 84 | 48 |

FIG. 52

SUBSTITUTE SHEET

86 / 86

FIG. 53

| <u>Name</u> | <u>Common Sequence</u> | <u>Variable Sequence</u> |
|-------------|------------------------|--------------------------|
| 5'-3-1 | AAGCTTGATCGAATTC | CGATCTTGC |
| -2 | | CGATCCTGC |
| -3 | | CGATCATGC |
| -4 | | CGATCGTGC |
| -5 | | CGAAGTTGC |
| -6 | | CGAAGCTGC |
| -7 | | AGATCTTGC |
| -8 | | AGATCCTGC |
| -9 | | AGATCATGC |
| -10 | | AGATCGTGC |
| -11 | | AGAAGTTGC |
| -12 | | AGAAGCTGC |
| -13 | | CGATCTTGT |
| -14 | | CGATCCTGT |
| -15 | | CGATCATGT |
| -16 | | CGATCGTGT |
| -17 | | CGAAGTTGT |
| -18 | | CGAAGCTGT |
| -19 | | AGATCTTGT |
| -20 | | AGATCCTGT |
| -21 | | AGATCATGT |
| -22 | | AGATCGTGT |
| -23 | | AGAAGTTGT |
| -24 | | AGAAGCTGT |
| -25 | | CGCTCTTGC |
| -26 | | CGCTCCTGC |
| -27 | | CGCTCATGC |
| -28 | | CGCTCGTGC |
| -29 | | CGCAGTTGC |
| -30 | | CGCAGCTGC |
| -31 | | CGCTCTTGT |
| -32 | | CGCTCCTGT |
| -33 | | CGCTCATGT |
| -34 | | CGCTCGTGT |
| -35 | | CGCAGTTGT |
| -36 | | CGCAGCTGT |

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No **PCT/US90/02853**

| | | | | | | |
|--|--|---|------------------------------------|------------------------|------|-------|
| I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ³
According to International Patent Classification (IPC) or to both National Classification and IPC
IPC (5): C12Q 1/20
U.S. CL.: 435/5 | | | | | | |
| II. FIELDS SEARCHED
<div style="text-align: right; margin-right: 100px;">Minimum Documentation Searched ⁴</div> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;">Classification System ¹</td> <td style="width: 50%; border: none;">Classification Symbols</td> </tr> <tr> <td style="border: none; text-align: center; padding-top: 20px;">U.S.</td> <td style="border: none; text-align: center; padding-top: 20px;">435/5</td> </tr> </table> <div style="text-align: center; margin-top: 10px;">Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched ⁵</div> <div style="padding: 10px; margin-top: 10px;"> Databases: USPTO Automated Patent System (File U.S. Pat. 1925-90)
 Genebank UEMBL </div> | | | Classification System ¹ | Classification Symbols | U.S. | 435/5 |
| Classification System ¹ | Classification Symbols | | | | | |
| U.S. | 435/5 | | | | | |
| III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴ | | | | | | |
| Category ⁶ | Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷ | Relevant to Claim No. ¹⁸ | | | | |
| X | US, A, 4,683,195 (MULLIS ET AL) 28 July 1987
See the entire document. | 16, 17, 19 | | | | |
| X | US, A, 4,683,202 (MULLIS ET AL) 28 July 1987
See the entire document. | 16, 17, 19 | | | | |
| <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁵ * Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p> </div> </div> | | | | | | |
| IV. CERTIFICATION | | | | | | |
| Date of the Actual Completion of the International Search ⁹

<div style="text-align: center; font-size: 1.2em;">22 AUGUST 1990</div> | | Date of Mailing of this International Search Report ⁸

<div style="text-align: center; font-size: 1.5em; font-weight: bold;">28 SEP 1990</div> | | | | |
| International Searching Authority ¹

<div style="text-align: center;">ISA/US</div> | | Signature of Authorized Officer ¹⁰
<div style="text-align: center;"> BRADLEY L. SISSON </div> | | | | |